



# Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA


## Physics Division

Submitted to Physical Review Letters

### A Geometry of the Generations

L.J. Hall and H. Murayama

August 1995



REFERENCE COPY  
Does Not  
Circulate

81dg. 50 Library.

Copy 1

LBL-37627

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

August 14, 1995

LBL-37627  
UCB-PTH-95/29

## A Geometry of the Generations<sup>1</sup>

Lawrence J. Hall and Hitoshi Murayama  
*Department of Physics, University of California  
Berkeley, California 94720*  
and  
*Theoretical Physics Group  
Ernest Orlando Lawrence Berkeley National Laboratory  
University of California, Berkeley, California 94720*

### Abstract

We propose a geometric theory of flavor based on the discrete group  $(S_3)^3$ , in the context of the minimal supersymmetric standard model. The group treats three objects symmetrically, while making fundamental distinctions between the generations. The top quark is the only heavy quark in the symmetry limit, and the first and second generation squarks are degenerate. The hierarchical nature of Yukawa matrices is a consequence of a sequential breaking of  $(S_3)^3$ .

---

<sup>1</sup>This work was supported in part by the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, Division of High Energy Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098 and in part by the National Science Foundation under grant PHY-90-21139.

The smallness of the electroweak symmetry breaking scale and the hierarchical nature of the Yukawa matrices provide two of the most important problems of particle physics. Weak scale supersymmetry may well play a crucial role in the former, since it is the only symmetry which can protect the mass of an elementary scalar. However, weak scale supersymmetry widens the scope of flavor physics: any supersymmetric extension of the standard model possesses eleven flavor matrices rather than the three Yukawa matrices of the standard model. The additional eight flavor matrices all involve couplings to squarks and sleptons, and have therefore not been directly probed experimentally. However, rare processes, such as the  $K_L$ - $K_S$  mass difference provide experimental constraints on these flavor mixing matrices [1]. Hence the problem of flavor symmetries is greatly affected by the inclusion of weak scale supersymmetry.

It is frequently remarked that the most striking feature of the observed flavor physics is that the top quark is the only fermion with a mass of order the weak scale. In the context of the standard model this implies that only one entry of the three Yukawa matrices is of order unity, while all other entries are numerically small. In the context of supersymmetric standard model, we find that there are now *two* features of flavor physics which must be considered at the zeroth order level: (1) the large mass of the top quark, (2) the near absence of flavor-changing neutral currents strongly suggest that scalars of a given charge of the light two generations are degenerate [1, 2]. In this paper we explore the consequences of assuming that both of these salient features arise from a common origin – a flavor symmetry group  $G_f$ .

The existence of an exact flavor symmetry group at high energies is very plausible – it is suggested by the replication of generations. However, in many supersymmetric theories it becomes a necessity. Presumably the ultimate theory of flavor will involve no small parameters: all the dimensionless couplings will be of order unity and small mass ratios will result from hierarchies of dynamically generated mass scales, or perhaps from loop factors. If the supersymmetry breaking squark masses appear as hard interactions in such theories, as they do in supergravity models, then the couplings of order unity will lead to large radiative contributions to the squark masses [4]. The degeneracy between first two generation scalars can then only be maintained if the dimensionless couplings of the theory possess a non-Abelian flavor symmetry  $G_f$ .

What should we take for  $G_f$ ? In the context of supergravity theories it

was suggested that a  $U(N)$  invariance of the Kähler potential, where  $N$  is the total number of chiral superfields of the theory, be used to protect the squark degeneracy [5]. However, in this paper we require that  $G_f$  also acts on the superpotential interactions which generate the fermion masses, so this  $U(N)$  invariance is not possible. Flavor symmetries which have been considered to date fall into two categories:

(1) **Unified** The group is such that in the symmetry limit there is no distinction whatever between generations. This occurs if the three generations are assigned to an irreducible representation which has three indistinguishable components – such as a triplet of  $SU(3)$ .

(2) **Asymmetric** The action of the group is such that there is no symmetrical treatment of  $N$  objects, where  $N = 3$  is the number of generations. There are many examples with  $G_f$  taken to be  $U(1)^n$  [6] or  $SU(2)$  [7].

A unified  $G_f$  has the advantage of providing a more complete theory of flavor, whereas an asymmetric  $G_f$  does not provide an understanding of the difference between the generations. On the other hand, a unified  $G_f$  must be broken by couplings of order unity to obtain  $m_t$ , whereas an asymmetric  $G_f$ , such as  $SU(2)$ , can provide an understanding of the salient flavor features even in the absence of symmetry breaking. In this paper we propose to combine the advantages of a unified  $G_f$  with those of an asymmetric  $G_f$  by introducing a third category of flavor symmetry:

(3) **Symmetric** The group has an action which is identical on three objects, yet has a representation structure which treats the generations differently.

In searching for such a group we are guided by three principles:

(a) The fields of the theory are those of the minimal supersymmetric standard model: three generations and two Higgs doublets  $H_u$  and  $H_d$ .

(b) The group should be a local discrete symmetry [8]. Continuous global symmetries are broken by quantum gravity [9] and should therefore be gauged. However, flavor symmetries must be broken to generate Yukawa matrices, and the breakdown of gauged flavor symmetries splits masses of different families due to the  $D$ -term contribution [10].

(c) The representation structure of the three generations should be  $(1 + 2)$ , such that, in the  $G_f$  symmetry limit, the top quark Yukawa coupling is allowed, and the non-Abelian nature of the group maintains degeneracy between the scalars of the lighter two generations.

The discrete non-Abelian group with fewest group elements is the symmetric group  $S_3$ . By its very definition it acts symmetrically on three objects.

Remarkably it has two singlets and a doublet as irreducible representations, and therefore offers an excellent match to the flavor problem of supersymmetric theories. The action of  $S_3$  has a geometrical interpretation as all possible rotations in three dimensions which leave an equilateral triangle invariant. The three vectors representing the vertices of the triangle,  $e_1$ ,  $e_2$  and  $e_3$  in Figure 1, are treated identically by the group. Yet the sums and differences of these vectors form a singlet representation ( $v_3$ ) and a doublet representation ( $v_1, v_2$ ), whose two components have different group properties. Despite a geometrical symmetry amongst three objects, there is also a geometrical understanding of the differences between the generations.

The group  $S_3$  has six elements:

$$S_3 = \{e, (12), (13), (23), (123), (132)\}, \quad (1)$$

where  $e$  is the identity element. The two elements (123) and (132) are  $120^\circ$  rotation of the triangle around the axis  $v_3 = (1, 1, 1)/\sqrt{3}$ , which form  $Z_3$  subgroup of even permutations in  $S_3$ . The (12), (13) and (23) elements rotate the triangle by  $180^\circ$  around one of its symmetry axes, which are odd permutations. The vector  $v_3$  flips its sign under odd permutations but does not under even permutations. This is a non-trivial singlet representation which we call  $\mathbf{1}_A$ , and will be identified with the third generation later. Two other orthogonal vectors  $v_1 = (1, 1, -2)/\sqrt{6}$  and  $v_2 = (-1, 1, 0)/\sqrt{2}$  form a doublet representation  $\mathbf{2}$  of  $S_3$ . We identify them later with first and second generation fields, respectively. Any  $\mathbf{2}$  representation can be written as a two-vector in  $(v_1, v_2)$  space. There are only three irreducible representations of  $S_3$ :  $\mathbf{1}_A$  and  $\mathbf{2}$  above and another singlet  $\mathbf{1}_S$  which is a trivial representation (invariant). The  $\mathbf{1}_S$  representation can be obtained as a symmetric product  ${}^t\mathbf{x}\mathbf{y}$  of two  $\mathbf{2}$ 's,  $\mathbf{x}_i$  and  $\mathbf{y}_i$ , while  $\mathbf{1}_A$  is an anti-symmetric product  ${}^t\mathbf{x}\sigma_2\mathbf{y}$ . The other combinations form a  $\mathbf{2}$  such as  ${}^t\mathbf{x}\sigma_3\mathbf{y} \sim v_1$  and  ${}^t\mathbf{x}\sigma_1\mathbf{y} \sim -v_2$ , and  $\mathbf{2}^3$  contains a totally symmetric invariant  $({}^t\mathbf{x}\sigma_3\mathbf{y})z_1 - ({}^t\mathbf{x}\sigma_1\mathbf{y})z_2$ . Decomposition of tensor products is shown in Table 1.

The group  $S_3$  has been used before in the context of flavor physics, but from a different perspective. The democratic ansatz for quark mass matrices [11], which leads to a heavy top quark, is known to possess the symmetry group  $S_3 \times S_3 \times SU(2) \times SU(2)$  [12]. However, in this work the fundamental origin of the flavor structure was assumed to come from other dynamics, perhaps BCS-like, and  $S_3$  simply appeared as an accidental consequence of this

democratic dynamics. In contrast, in this paper we argue that the supersymmetric flavor puzzle suggests uniquely that  $S_3$  is the fundamental origin of flavor.

The work [13] is somewhat closer to our philosophy. They advocate  $G_f = Q_{2n}$ , the dicyclic dihedral groups, rather than  $D_n$  or  $S_n$ , based on an anomaly freedom constraint which we find to be too restrictive. They also need “Q-leptons” to cancel anomalies. Furthermore, although  $Q_{2n}$  possesses only singlet and doublet representations and therefore allows a large  $m_t$ , this is clearly also possible with Abelian groups. In this paper we combine the supersymmetric motivation for some non-Abelian nature to  $G_f$  with the aesthetic desire for a symmetric flavor group.

Despite the encouraging features of  $S_3$ , it is not possible to satisfy the few guiding principles (a), (b), (c) above using a single  $S_3$  as  $G_f$ . For  $\bar{d}$  and  $\bar{s}$  to be degenerate,  $(d, s)_L$  and  $(d, s)_R$  should both transform as  $\mathbf{2}$ . Since  $\mathbf{2} \times \mathbf{2} = \mathbf{1}_A + \mathbf{1}_S + \mathbf{2}$ ,  $m_d$  and  $m_s$  are allowed by  $S_3$ , no matter whether  $H_d$  is assigned to  $\mathbf{1}_A$  or to  $\mathbf{1}_S$ . An enlargement of the group is thus necessary. One possibility is to search for interesting structures in larger discrete groups, such as  $S_n$ ,  $D_n$ ,  $Q_{2n}$ , and  $\Delta(3n^2)$  [14, 13]. We find the geometric picture of the three generations arising from the symmetric action of  $S_3$  to be sufficiently compelling that we prefer to replicate  $S_3$  factors. Hence we consider a group  $S_3^Q \times S_3^U \times S_3^D$  with each of  $Q$ ,  $U$ ,  $D$  transform as  $\mathbf{1} + \mathbf{2}$  under its own  $S_3$ , while transforming trivially under other factors.

We identify the third generation with  $\mathbf{1}_A$  rather than  $\mathbf{1}_S$ , because we would like to consider the discrete flavor group as an anomaly-free gauge symmetry. The only anomaly one can discuss with the low-energy particle content alone is  $S_3 \times H^2$  where  $H$  is either  $SU(2)$  or  $SU(3)$  in the standard model [15]. Consider the element (12), which leaves  $\mathbf{1}_S$  and  $v_1$  in  $\mathbf{2}$  invariant but changes sign of  $\mathbf{1}_A$  and  $v_2$  in  $\mathbf{2}$ . To avoid an anomaly, the total number of  $\mathbf{1}_A$  and  $\mathbf{2}$  with a given quantum number has to be even. In our context, this requirement uniquely selects  $\mathbf{1}_A + \mathbf{2}$ . The anomaly freedom of this choice can be easily understood by noting a vector  $\mathbf{3}$  in an anomaly free group  $SO(3)$  decomposes to  $\mathbf{1}_A + \mathbf{2}$ . Furthermore, this choice is precisely the one which allows a geometric interpretation of families in terms of rotations. It is interesting to see that the three generations, although in a reducible representation  $\mathbf{1}_A + \mathbf{2}$ , require each other to render the theory consistent quantum mechanically.

The flavor transformation properties of the quarks are shown in Table 2.

The quantum number of  $H_u$  is fixed to allow  $m_t$  by  $G_f$ . Anomaly freedom then dictates an identical transformation for  $H_d$ . Because of this charge assignment, only the top quark is heavy in the  $(S_3)^3$  symmetric limit. The top-bottom asymmetry, or more generally the up-down asymmetry, is built into the representation structure of the Higgs. On the other hand, squark mass matrices all have the form  $\text{diag}(M_1^2, M_1^2, M_3^2)$ . The lepton sector will be discussed elsewhere.

Now we consider the breaking of  $(S_3)^3$  symmetry and discuss its consequence on the Yukawa and squark mass matrices. In order to keep the number of breaking parameters as small as possible, we take the following “minimal” form of the Yukawa matrices [16],

$$Y_u = \left( \begin{array}{cc|c} h_u & O(\sqrt{h_u h_c}) & -h_t \lambda^3 A(\rho + i\eta) \\ O(\sqrt{h_u h_c}) & h_c & -h_t \lambda^2 A \\ \hline 0 & 0 & h_t \end{array} \right), \quad (2)$$

$$Y_d = \left( \begin{array}{cc|c} h_d & h_s \lambda & 0 \\ O(h_s \lambda) & h_s & 0 \\ \hline 0 & 0 & h_b \end{array} \right), \quad (3)$$

which correctly reproduce Cabbibo–Kobayashi–Maskawa (CKM) matrix in Wolfenstein parametrization. The quark masses are related to the Yukawa couplings by  $m_{u,c,t} = h_{u,c,t} \langle H_u \rangle$  and  $m_{d,s,b} = h_{d,s,b} \langle H_d \rangle$ . We assumed  $(Y_u)_{12}$  and  $(Y_u)_{21}$  to be  $O(\sqrt{h_u h_c})$ , because larger off-diagonal elements need a fine-tuning in the determinant. We actually do not need these elements and can set them vanishing, but we kept them to make the discussion more general. The same comment applies to  $(Y_d)_{21}$ . The Cabbibo angle originates in the down sector and it may be possible to keep the famous relation  $\lambda \sim \sqrt{m_d/m_s}$ .

The largest breaking parameters in the Yukawa matrices are  $h_b$  which transforms as  $(\mathbf{1}_S, \mathbf{1}_A, \mathbf{1}_A)$  and  $h_t \lambda^2 A$  as  $(\mathbf{2}, \mathbf{1}_S, \mathbf{1}_S)$ .  $h_b$  breaks  $S_3^U \times S_3^D$  down to a subgroup  $S_3^U \times S_3^D / \mathbf{Z}_2$ , where (even, odd) and (odd, even) elements are removed. Note that the diagonal subgroup  $S_3^{U,D}$  is a subgroup of the unbroken symmetry.  $h_t \lambda^2 A$  is a  $\mathbf{v}_1$  element in a doublet, and breaks  $S_3^Q$  to  $S_2^Q \simeq \mathbf{Z}_2 = \{e, (12)\}$ . This  $\mathbf{Z}_2$  flips the sign of second generation and Higgs fields, while leaving first generation field unchanged. Therefore  $Q_2$  can acquire a Yukawa coupling while  $Q_1$  cannot.  $h_c$  and  $h_s$  belong to breaking parameters  $(\mathbf{2}, \mathbf{2}, \mathbf{1}_S)$  and  $(\mathbf{2}, \mathbf{1}_A, \mathbf{2})$ , respectively, and break the diagonal  $S_3^{U,D}$  to  $\mathbf{Z}_2$  as well, which still keep all first generation fields massless. After

including the smaller breaking parameters, the symmetry  $(S_3)^3$  is completely broken. In this way, the hierarchical pattern of the Yukawa matrices can be understood as a sequential breaking of the flavor symmetry.

Now we turn to the squark mass matrices. Since the constraints from the flavor-changing neutral currents are at best of order a few times  $10^{-3}$ , we work out the non-degeneracy in squark masses down to this order. It is straightforward to work out how the breaking parameters enter the scalar matrices. For  $m_Q^2$  matrix, the leading correction comes from  $(\mathbf{2}, \mathbf{1}_S, \mathbf{1}_S)$  with  $\mathbf{v}_1$  and  $\mathbf{v}_2$  components of  $O(h_t \lambda^2 A)$  and  $O(h_t \lambda^3 A(\rho + i\eta))$ , respectively. Therefore,

$$m_Q^2 \sim \left( \begin{array}{cc|c} M_1^2 + m^2 h_t \lambda^2 A & m^2 h_t \lambda^3 A(\rho + i\eta) & -m'^2 h_t \lambda^2 A \\ m^2 h_t \lambda^3 A(\rho + i\eta) & M_1^2 - m^2 h_t \lambda^2 A & m'^2 h_t \lambda^3 A(\rho + i\eta) \\ \hline -m'^2 h_t \lambda^2 A & m'^2 h_t \lambda^3 A(\rho - i\eta) & M_3^2 \end{array} \right), \quad (4)$$

where a possible correction to  $(m_Q^2)_{33}$  was absorbed into  $M_3^2$ . Here and hereafter,  $m^2$  and  $m'^2$  are arbitrary numbers comparable to  $M_1^2$  and  $M_3^2$ , and they are in general different for  $Q, U, D$ . For the  $m_U^2$  matrix, the only correction comes from the square of  $(\mathbf{2}, \mathbf{2}, \mathbf{1}_S)$  breaking parameter of  $O(h_c^2)$ . The resulting form is

$$m_U^2 \sim \left( \begin{array}{cc|c} M_1^2 + m^2 h_c^2 & m^2 \sqrt{h_c^3 h_u} & -m'^2 h_c^2 \\ m^2 \sqrt{h_c^3 h_u} & M_1^2 - m^2 h_c^2 & m'^2 \sqrt{h_c^3 h_u} \\ \hline -m'^2 h_c^2 & m'^2 \sqrt{h_c^3 h_u} & M_3^2 \end{array} \right). \quad (5)$$

The  $m_D^2$  matrix receives corrections from two sources at the leading order. One is the square of the  $(\mathbf{2}, \mathbf{1}_A, \mathbf{2})$  breaking parameter of  $O(h_s^2)$ , and the other is a product of three breaking parameters  $(\mathbf{1}_S, \mathbf{1}_A, \mathbf{1}_A)$ ,  $(\mathbf{2}, \mathbf{1}_A, \mathbf{2})$ , and  $(\mathbf{2}, \mathbf{1}_S, \mathbf{1}_S)$  of  $O(h_s h_b h_t A \lambda^2)$ . They are of the same order of magnitude and have the same group theoretical structure  $(\mathbf{1}_S, \mathbf{1}_S, \mathbf{2})$ . We keep only the first for simplicity and obtain

$$m_D^2 \sim \left( \begin{array}{cc|c} M_1^2 m^2 h_s^2 & m^2 h_s^2 \lambda & -m'^2 h_s^2 \\ m^2 h_s^2 \lambda & M_1^2 - m^2 h_s^2 & m'^2 h_s^2 \lambda \\ \hline -m'^2 h_s^2 & m'^2 h_s^2 \lambda & M_3^2 \end{array} \right). \quad (6)$$

The authors of [3] listed the constraints on the off-diagonal mass matrix elements for  $m_{\tilde{q}} \sim 1$  TeV in the basis where the Yukawa matrices are diagonal.

We adopt their notation and list the constraints in Tables 2, 3, 4. It is clear that our mass matrices satisfy all constraints rather easily. We have not discussed the left-right mixing mass matrix so far, but they are tightly constrained by the  $S_3^3$  symmetry as well. The breaking parameters enter the mixing mass matrix in the same manner as in the Yukawa matrices. It is easy to work them out and see that the constraints are easily satisfied.

A natural question is how much stronger the constraints become when we introduce further breaking parameters and introduce mixing in the right-handed fields as well. The off-diagonal elements of  $m_D^2$  and  $m_Y^2$  can be much larger than the above estimates. However, they are at most of the same order as those in  $m_Q^2$  if we assume a similar order of mixing angles in the right-handed fields. On the other hand, constraints become even weaker if we attribute all CKM angles to the down sector, since the breaking parameters are then proportional to  $h_b$  rather than  $h_t$ . A potentially dangerous breaking is that in  $(\mathbf{1}_S, \mathbf{1}_S, \mathbf{1}_A)$  or  $(\mathbf{1}_A, \mathbf{1}_S, \mathbf{1}_S)$ , which do not contribute to the Yukawa matrices. However they are presumably as small as  $h_u$  or  $h_d$  because they break the  $Z_2$  symmetry which keeps the first generation fields massless.

In summary, we proposed a geometric theory of flavor based on the discrete group  $(S_3)^3$ . The group acts symmetrically on three objects, yet gives fundamentally different characteristics to each generation. The three generations belong to a reducible representation  $\mathbf{2} + \mathbf{1}_A$ ; although they are not unified, they require each other for anomaly cancellations. Only the top quark is heavy in the symmetry limit, and first- and second-generation squarks are degenerate. Hierarchical Yukawa matrices can be understood as a consequence of sequential symmetry breaking. Flavor-changing processes are highly suppressed, allowing squarks at Tevatron energies.

### Acknowledgements

This work was supported in part by the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, Division of High Energy Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098 and in part by the National Science Foundation under grant PHY-90-21139.

## References

- [1] S. Dimopoulos and H. Georgi, *Nucl. Phys.* **B193**, 150 (1981).
- [2] This degeneracy is not a necessity, as flavor-changing effects in the down sector can be greatly suppressed by assuming that the origin of the CKM matrix resides in the up sector [3]. However, it is somewhat contrived to choose the flavor symmetries of the theory so that the flavor violations arise only in the sector of the theory where the experimental constraints are the weakest.
- [3] Y. Nir and N. Seiberg, *Phys. Lett.* **B309**, 337 (1993).
- [4] L.J. Hall, V.A. Kostelecky, and S. Raby, *Nucl. Phys.* **B267**, 415 (1986).
- [5] L.J. Hall, J. Lykken, and S. Weinberg, *Phys. Rev.* **D27**, 2359 (1983).
- [6] See, *e.g.*, C.D. Froggatt and H.B. Nielsen, *Nucl. Phys.* **B147**, 277 (1979); L. Ibañez and G.G. Ross, *Phys. Lett.* **B332**, 100 (1994).
- [7] M. Dine, A. Kagan, and R. Leigh, *Phys. Rev.* **D48**, 4269 (1993).
- [8] L.M. Krauss and F. Wilczek, *Phys. Rev. Lett.* **62**, 1221 (1989).
- [9] S. Giddings and A. Strominger, *Nucl. Phys.* **B307**, 854 (1988); S. Coleman, *Nucl. Phys.* **B310**, 643 (1988); G. Gilbert, *Nucl. Phys.* **B328**, 159 (1989).
- [10] Y. Kawamura, H. Murayama and M. Yamaguchi, *Phys. Rev.* **D51**, 1337 (1995); H. Murayama, LBL-36962, hep-ph/9503392.
- [11] H. Fritzsch, *Nucl. Phys.* **B 155**, 189 (1979).
- [12] H. Fritzsch and J. Plankl, *Phys. Lett.* **237 B**, 451 (1990).
- [13] P.H. Frampton and T.W. Kephart, *Phys. Rev.* **D51**, 1 (1995); IFP-702-UNC, hep-ph/9409330.
- [14] D.B. Kaplan and M. Schmaltz, *Phys. Rev.* **D49**, 3741 (1994).
- [15] T. Banks and M. Dine, *Phys. Rev.* **D45**, 1424 (1992); L.E. Ibañez, *Nucl. Phys.* **B398**, 301 (1993).

- [16] C.D. Carone and H. Murayama, LBL-37099, hep-ph/9504393, to appear in *Phys. Rev. D*.

$\otimes$	$\mathbf{1}_S$	$\mathbf{1}_A$	$\mathbf{2}$
$\mathbf{1}_S$	$\mathbf{1}_S$	$\mathbf{1}_A$	$\mathbf{2}$
$\mathbf{1}_A$	$\mathbf{1}_A$	$\mathbf{1}_S$	$\mathbf{2}$
$\mathbf{2}$	$\mathbf{2}$	$\mathbf{2}$	$\mathbf{1}_A \oplus \mathbf{1}_S \oplus \mathbf{2}$

Table 1: Decomposition of tensor product of two representations into irreducible representations.

	$Q$	$U$	$D$	$H_u$	$H_d$
$S_3^Q$	$(\mathbf{1}_A, \mathbf{2})$	-	-	$\mathbf{1}_A$	$\mathbf{1}_A$
$S_3^U$	-	$(\mathbf{1}_A, \mathbf{2})$	-	$\mathbf{1}_A$	$\mathbf{1}_A$
$S_3^D$	-	-	$(\mathbf{1}_A, \mathbf{2})$	-	-

Table 2: Quantum number assignments of the fields under  $(S_3)^3$  symmetry.  $Q$  refers to left-handed quark doublets,  $U$  ( $D$ ) to right-handed up(down)-type quarks.

	$(\delta_{LL}^d)_{12}$	$(\delta_{RR}^d)_{12}$	$(\delta_{LR}^d)_{12}$	$\langle \delta_{12}^d \rangle$
upper bound [3]	0.05	0.05	0.008	0.006
this model	$h_t A \lambda^3$	$h_s^2 \lambda^2$	$h_s^2 \lambda$	$\sqrt{(\delta_{LL}^d)_{12} (\delta_{RR}^d)_{12}}$

Table 3: The constraints and the consequence of  $(S_3)^3$  symmetry on the mass splittings in  $\tilde{d}$ - $\tilde{s}$ .

	$(\delta_{LL}^d)_{13}$	$(\delta_{RR}^d)_{13}$	$(\delta_{LR}^d)_{13}$	$\langle \delta_{13}^d \rangle$
upper bound [3]	0.1	0.1	0.06	0.04
this model	$h_t A \lambda^2$	$h_s^2$	$h_b h_t A \lambda^3$	$\sqrt{(\delta_{LL}^d)_{13} (\delta_{RR}^d)_{13}}$

Table 4: The constraints and the consequence of  $(S_3)^3$  symmetry on the mass splittings in  $\tilde{d}$ - $\tilde{b}$ .

	$(\delta_{LL}^u)_{12}$	$(\delta_{RR}^u)_{12}$	$(\delta_{LR}^u)_{12}$	$(\delta_{LR}^u)_{12}$
upper bound [3]	0.1	0.1	0.06	0.04
this model	$h_t A \lambda^3$	$\sqrt{h_u h_c^3}$	$\sqrt{h_u h_c^3}$	$\sqrt{(\delta_{LL}^u)_{12} (\delta_{RR}^u)_{12}}$

Table 5: The constraints and the consequence of  $(S_3)^3$  symmetry on the mass splittings in  $\tilde{u}$ - $\tilde{c}$ . We assumed that the rotation angle between  $u$  and  $c$  is  $O(\sqrt{h_u/h_c})$ .




Figure 1:  $S_3$  acts as a rotation of the triangle spanned by three orthonormal vectors  $e_{1,2,3}$ . The vector  $v_3$  corresponds to the  $1_A$  representation, and two vectors  $v_{1,2}$ , in the plane of the triangle, to the  $2$  representation.

LAWRENCE BERKELEY NATIONAL LABORATORY  
UNIVERSITY OF CALIFORNIA  
TECHNICAL & ELECTRONIC INFORMATION DEPARTMENT  
BERKELEY, CALIFORNIA 94720