Submitted to Inorganic Chemistry

LBL-3769
Preprint c.1

KINETICS OF THE REACTION OF AMMONIUM ION WITH HYDROBORATE ION IN LIQUID AMMONIA

Thomas S. Briggs and William L. Jolly

March 1975

Prepared for the U. S. Atomic Energy Commission
under Contract W-7405-ENG-48

For Reference

Not to be taken from this room

LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY

LBL-3769
c.1

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

0 0 0 0 4 3 0 0 1 3 0

Kinetics of the Reaction of Ammonium Ion with Hydroborate Ion in Liquid Ammonia

by

Thomas S. Briggs and William L. Jolly

Chemistry Department, University of California, and
Inorganic Materials Research Division,
Lawrence Berkeley Laboratory,
Berkeley, California 94720

The acid-catalyzed hydrolysis of hydroborate has been thoroughly studied,[1] but little has been reported on the analogous ammonolysis in liquid ammonia. Armstrong[2] studied the reaction

$$(NH_4)_2SO_4 + 2NaBH_4 \longrightarrow Na_2SO_4 + 2H_2 + 2BNH_6 \qquad (1)$$

in ammonia from 40 to 70°. The BNH_6 product was not completely identified. We have investigated the kinetics of this reaction from 25 to 45° using NH_4Br instead of $(NH_4)_2SO_4$. In addition to obtaining rate constants and a heat of activation, we have determined the effect of ionic strength from $\mu = 0.01$ M to $\mu = 1.92$ M and have identified the product as BH_3NH_3.

Experimental Section

Reagent grade ammonium and sodium bromide (dried at 85°) and analytical reagent sodium hydroborate (Ventron) were used. Ammonia was distilled from a sodium solution.

The reactions were carried out in a magnetically-stirred, glass-lined 128-ml Parr general-purpose bomb. This was connected by high pressure fittings to one side of a mercury-containing U-tube manometer. The other side of the manometer was connected to a reference bomb containing pure liquid ammonia. The bombs were held in a thermostatted water bath, and the entire apparatus was enclosed in an air bath thermostatted at a temperature slightly above that of the water bath. The reaction rates were measured by reading the manometer with the aid of a cathetometer as a function of time.

The ammonia was first distilled into a separate bomb and weighed. The amounts of $NaBH_4$, NH_4Br, and NaBr required to make a solution of predetermined ionic strength were then weighed out into the reaction bomb, and the ammonia was finally distilled into the reaction bomb. Either the $NaBH_4$ or the NH_4Br was always in at least a 9-fold excess. After each run, the evolved hydrogen was Toepler-pumped into a gas buret and measured.

The pressures measured with the manometer corresponded to the partial pressures of hydrogen gas in the reaction bomb side. Rate constants were determined from plots of $\log[(P_\infty - P_t)/P_\infty]$ vs time.

The BNH_6 product was isolated by removal of ammonia on a vacuum line and extraction of the residue with ethyl ether. Evaporation of the ether left a white, crystalline solid which melted at 110-113°.

This melting temperature corresponds to NH_3BH_3.[3] Anal. Calcd for BNH_6: B, 35.03; N, 45.48. Found: B, 35.24; N, 46.16.

Results and Conclusions

From $\log[P_\infty - P)/P_\infty]$-vs-time plots such as that shown in Fig. 1, the reaction of NH_4^+ with BH_4^- was found to be first order in both NH_4^+ and BH_4^-. The second order rate constants calculated from the slopes of such plots for various temperatures are given in Table I. A log k-vs-1/T

Table I

Rate Constants in Liquid Ammonia at $\mu = 0.9$

Temperature, °C	k (M^{-1} sec^{-1})
25	5.88×10^{-6}
30	1.79×10^{-5}
35	5.04×10^{-5}
40	1.43×10^{-4}[a]
42	2.17×10^{-4}
45	3.40×10^{-4}[b]

[a]Average of 1.49×10^{-4} and 1.37×10^{-4}, from separate experiments.

[b]This value obtained in two experiments.

plot of the data, shown in Fig. 2, gave a least-squares heat of activation of 38.5 ± 0.6 kcal/mol. Our data may be compared with those of Gardiner and Collat,[4] who studied the aqueous reaction of

NH_4^+ with BH_4^-. They obtained a rate constant of 2.4×10^{-4} $\underline{M}^{-1}$ sec^{-1} at $\mu = 1$ and 25° and a heat of activation of 23 ± 2 kcal/mol at $\mu = 0.5$. The higher rate constant and lower activation energy of the aqueous reaction may be due to the participation, in the aqueous reaction, of water molecules hydrogen bonded to the ammonium ion. A Grotthuss-type proton transfer such as the following may be involved:



Because hydrogen bonding is relatively unimportant in liquid ammonia, the analogous mechanism in liquid ammonia is unlikely.

The Bronsted-Bjerrum theory of ionic reaction leads to the expression

$$\log_{10}(k/k_0) = \frac{2\alpha Z_A Z_B \sqrt{\mu}}{1 + \beta a \sqrt{\mu}}$$

which was applied to rate constants measured at various ionic strengths at 30° (Table II). Using the dielectric constant[5] of 16.5 for ammonia

Table II

Rate Constants in Liquid Ammonia at 30°

Ionic Strength	BH_4^-, conc., $\underline{M}$	NH_4^+, conc., $\underline{M}$	k ($\underline{M}^{-1}$ sec^{-1})
0.0098	0.00888	0.00094	2.07×10^{-3}
0.049	0.0446	0.0045	6.23×10^{-4}
0.20	0.1954	0.0046	1.94×10^{-4}
0.90	0.818	0.0818	1.79×10^{-5}
1.92[a]	0.818	0.0818	4.95×10^{-6}

[a]NaBr was added to increase the ionic strength.

at 30°, we calculated[6] $\alpha = 5.16$ and $\beta = 0.711$. A value for a of 4.7 Å was used in constructing the Bronsted-Bjerrum plot of the data, shown in Figure 3. The experimental points at relatively low ionic strengths have a slope close to that predicted by equation 2, whereas the points at higher ionic strengths are low, probably because of the pairing of reactant ions with inert ions. Dewald and Browall[7] correlated rate data for the reaction of the electron with the tri-n-butyl-n-propylammonium ion with the Bronsted-Bjerrum equation using a value for a of 6 Å.

Acknowledgement

This work was supported by the Energy Research and Development Administration.

-References-

(1) R. E. Davis, and C. G. Swain, J. Amer. Chem. Soc., 82, 5949 (1960); R. E. Davis, C. L. Kibby and C. G. Swain, J. Amer. Chem. Soc., 82, 5950 (1960); W. H. Stockmayer, R. R. Miller and R. J. Zeto, J. Phys. Chem., 65, 1076 (1961); R. E. Mesmer and W. L. Jolly, Inorg. Chem., 1, 608 (1962); J. A. Gardiner and J. W. Collat, J. Amer. Chem. Soc., 86, 3165 (1964); J. A. Gardiner and J. W. Collat, J. Amer. Chem. Soc., 87, 1692 (1965); F. T. Wang and W. L. Jolly, Inorg. Chem., 11, 1933 (1972).

(2) D. L. Armstrong, Aerojet Engineering Corp. Annual Report No. 367, Defense Documentation Center File No. ATI50631 (11 March 1949).

(3) E. Mayer, Inorg. Chem., 11, 866 (1972).

(4) J. A. Gardiner and J. W. Collat, J. Amer. Chem. Soc., 87, 1692 (1965).

(5) J. Jander, "Anorganische und allgemeine Chemie in flüssigem Ammoniak," John Wiley, New York, 1966, p. 23.

(6) W. J. Moore, "Physical Chemistry," Prentice-Hall, New Jersey, 1963, 3rd Ed., p. 358.

(7) R. R. Dewald and K. W. Browall, Chem. Comm., 1511 (1968).

-Figure Captions-

Fig. 1 Plot of $\log[(P_\infty - P)/P_\infty]$ vs time for reactions at 45° and $\mu = 0.9$: O from 0.08 M BH_4^- with 0.82 M NH_4^+; Δ from 0.08 M NH_4^+ with 0.82 M BH_4^-.

Fig. 2 Temperature dependence of k at $\mu = 0.9$.

Fig. 3 Ionic strength dependence of k at 30°. The line is drawn with the slope given by eq. 2.



XBL 753-665

Fig. 1



XBL 753-664

Fig. 2



XBL 753-663

Fig. 3

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720