



# Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

## Physics Division

Submitted to Physical Review D

### **R-Parity Violation in Flavor Changing Neutral Current Processes and Top Quark Decays**

K. Agashe and M. Graesser

October 1995



REFERENCE COPY  
Does Not  
Circulate  
Bldg. 50 Library.  
Copy 1  
LBL-37823

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

October 27, 1995

LBL-37823

UCB-PTH-95/33

***R-Parity Violation in Flavor Changing Neutral Current  
Processes and Top Quark Decays \****

K. Agashe<sup>†</sup> and M. Graesser<sup>† §</sup>

*Theoretical Physics Group*

*Lawrence Berkeley National Laboratory*

*University of California*

*Berkeley, California 94720*

---

\*This work was supported in part by the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, Division of High Energy Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098 and in part by the National Science Foundation under grant PHY-90-21139.

<sup>†</sup>email: [agashe@theor1.lbl.gov](mailto:agashe@theor1.lbl.gov)

<sup>†</sup>email: [graesser@theor1.lbl.gov](mailto:graesser@theor1.lbl.gov)

<sup>§</sup>Supported by a Natural Sciences and Engineering Research Council of Canada Fellowship.

## Abstract

We show that supersymmetric  $R$ -parity breaking ( $R_p$ ) interactions always result in Flavor Changing Neutral Current (FCNC) processes. Within a single coupling scheme, these processes can be avoided in either the charge  $+2/3$  or the charge  $-1/3$  quark sector, but not both. These processes are used to place constraints on  $R_p$  couplings. The constraints on the first and the second generations are better than those existing in the literature. The  $R_p$  interactions may result in new top quark decays. Some of these violate electron-muon universality or produce a surplus of  $b$  quark events in  $t\bar{t}$  decays. Results from the CDF experiment are used to bound these  $R_p$  couplings.

# 1 Introduction

The Minimal Supersymmetric Standard Model (MSSM) [1] with the gauge group  $G = SU(3)_c \times SU(2)_L \times U(1)_Y$  contains the Standard Model particles and their superpartners, and an additional Higgs doublet. In order to produce the observed spectrum of particle masses, the superpotential is given by

$$\lambda_{ij}^L L_i E_j^c H + \lambda_{jk}^D H Q_j D_k^c + \lambda_{ij}^U U_i^c Q_j H' + \mu H H' \quad (1)$$

where  $L = \begin{pmatrix} N \\ E \end{pmatrix}$  and  $Q = \begin{pmatrix} U \\ D \end{pmatrix}$  denote the chiral superfields containing the lepton and quark  $SU(2)_L$  doublets and  $E^c$ ,  $U^c$  and  $D^c$  are the  $SU(2)_L$  singlets, all in the weak basis.  $H$  and  $H'$  are the Higgs doublets with hypercharges  $-1$  and  $+1$  respectively. The  $SU(2)_L$  and  $SU(3)_c$  indices are suppressed, and  $i, j$  and  $k$  are generation indices. However, requiring the Lagrangian to be gauge invariant does not uniquely determine the form of the superpotential. In addition, the following renormalizable terms

$$\lambda_{ijk} L_i L_j E_k^c + \bar{\lambda}_{ijk} L_i Q_j D_k^c + \lambda''_{ijk} U_i^c D_j^c D_k^c \quad (2)$$

are allowed\*. Unlike the interactions of the MSSM, these terms violate lepton number and baryon number. They may be forbidden by imposing a discrete symmetry,  $R$ -parity, which is  $(-1)^{3B+L+2S}$  on a component field with baryon

---

\*A term  $\mu_i L_i H'$  is also allowed. This may be rotated away through a redefinition of the  $L$  and  $H$  fields [2].

number  $B$ , lepton number  $L$  and spin  $S$ . Whether this symmetry is realized in nature must be determined by experiment. If both lepton and baryon number violating interactions are present, then limits on the proton lifetime place stringent constraints on the products of most of these couplings. So, it is usually assumed that if  $R$ -parity is violated, then either lepton or baryon number violating interactions, but not both, are present. It is interesting that despite the large limits on the proton lifetime, some products of the  $R$ -parity violating couplings remain bounded only by the requirement that the theory remain perturbative [3]. If either  $L_i Q_j D_k^c$  or  $U_i^c D_j^c D_k^c$  terms are present, flavor changing neutral current (FCNC) processes are induced. It has been assumed that if only one  $R$ -parity violating ( $\mathcal{R}_p$ ) coupling with a particular flavor structure is non-zero, then these flavor changing processes are avoided. In this *single coupling scheme* [4] then, efforts at constraining  $R$ -parity violation have concentrated on flavor conserving processes [5, 6, 7, 8, 9, 10]. It is surprising that, even though individual lepton or baryon number is violated in this scheme, the constraints are rather weak.

In Section 2, we demonstrate that the *single coupling scheme* cannot be realized in the quark mass basis. Despite the general values the couplings may have in the weak basis, after electroweak symmetry breaking there is at least one large  $\mathcal{R}_p$  coupling and many other  $\mathcal{R}_p$  couplings with different flavor structure. Therefore, in the mass basis the  $R$ -parity breaking couplings *cannot* be diagonal

in generation space. Thus, flavor changing neutral current processes are always present in either the charge  $2/3$  or the charge  $-1/3$  quark sectors. We use these processes to place constraints on  $R$ -parity breaking. We find constraints on the first and the second generations that are much stronger than existing limits.

The recent discovery of the top quark [11, 12] with the large mass of  $176 \text{ GeV}$  opens the possibility for the tree level decays  $t \rightarrow \tilde{l}_i^+ + d_k$  and  $t \rightarrow \tilde{d}_j + \bar{d}_k$  if  $R$ -parity is broken. If the  $R_p$  couplings are large enough, then these decay channels may be competitive with the Standard Model decay  $t \rightarrow b + W$ . As no inconsistencies between the measured branching fractions and production cross-section of the top quark and those predicted by the Standard Model (SM) have been reported, limits on the branching fractions for the  $R_p$  decay channels may be obtained. Since the existing lower bound on the mass of the lightest slepton is  $\sim 45 \text{ GeV}$  [13], while the strong interactions of the squarks make it likely that the squarks are heavier than the sleptons, the decay  $t \rightarrow \tilde{l}_i^+ + d_k$  is more probable. In our analysis, we therefore assume that only the slepton decay channel is present. In Section 3 we analyse the  $R_p$  top decay channels to place constraints on the  $t \rightarrow \tilde{l}_i^+ + d_k$  coupling. For this reason, in this paper we assume that only the  $\mathcal{L}$  terms  $L_i Q_j D_k^c$  are present. The conclusions of Section 2, however, are valid even if the  $L_i L_j E_k^c$  terms are also present. Constraints on products of couplings when both  $\mathcal{L}$  interactions are present may be found in reference [14]. In Section 4 we summarize our results and compare them with

limits existing in the literature.

## 2 Flavor Changing Neutral Current Processes

Flavor changing neutral current processes are more clearly seen by examining the structure of the interactions in the quark mass basis. In this basis, the  $\bar{\lambda}_{ijk}$  interactions are

$$\lambda'_{ijk}(N_i^m(V_{KM})_{jl}D_l^m - E_i^m U_j^m)D_k^{cm} \quad (3)$$

where

$$\lambda'_{ijk} = \bar{\lambda}_{imn}U_{Ljm}D_{Rnk}^* \quad (4)$$

The superfields in Equation (3) have their fermionic components in the mass basis so that the Cabibbo-Kobayashi-Masakawa (CKM) matrix [15]  $V_{KM}$  appears explicitly. The rotation matrices  $U_L$  and  $D_R$  appearing in the previous equation are defined by

$$u_{Li} = U_{Lij}u_{Lj}^m \quad (5)$$

$$d_{Ri} = D_{Rij}d_{Rj}^m \quad (6)$$

where  $q_i$  ( $q_i^m$ ) are quark fields in the weak (mass) basis. Henceforth, all the fields will be in the mass basis and we drop the superscript  $m$ .

Unitarity of the rotation matrices implies that the couplings  $\lambda'_{ijk}$  and  $\bar{\lambda}_{ijk}$

satisfy

$$\sum_{jk} |\lambda'_{ijk}|^2 = \sum_{mn} |\bar{\lambda}_{imn}|^2 \quad (7)$$

So any constraint on the  $R_p$  couplings in the quark mass basis also places a bound on the  $R_p$  couplings in the weak basis.

In terms of component fields, the interactions are

$$\lambda'_{ijk} [(V_{KM})_{jl} (\tilde{\nu}_L^i \bar{d}_R^k d_L^l + \bar{d}_L^l \bar{d}_R^k \nu_L^i + (\bar{d}_R^k)^* \overline{(\nu_L^i)^c} d_L^l) - \tilde{e}_L^i \bar{d}_R^k u_L^j - \tilde{u}_L^j \bar{d}_R^k e_L^i - (\bar{d}_R^k)^* \overline{(e_L^i)^c} u_L^j] \quad (8)$$

where  $e$  denotes the electron and  $\tilde{e}$  it's scalar partner and similarly for the other particles.

The contributions of the  $R$ -parity violating interactions to low energy processes involving no sparticles in the final state arise from using the  $R_p$  interactions an even number of times. If two  $\lambda'$  's or  $\lambda''$  's with different flavor structure are non-zero, flavor changing low energy processes can occur. These processes are considered in references [2] and [16], respectively. Therefore, it is usually assumed that either only one  $\lambda'$  with a particular flavor structure is non-zero, or that the  $R$ -parity breaking couplings are diagonal in generation space. However, Equation (8) indicates that this does not imply that there is only one set of interactions with a particular flavor structure, or even that they are diagonal in flavor space. In fact, in this case of one  $\lambda'_{ijk} \neq 0$ , the CKM matrix generates couplings involving each of the three down-type quarks. Thus, flavor violation

occurs in the down quark sector, though suppressed by the small values of the off-diagonal CKM elements. Below, we use these processes to obtain constraints on  $R$ -parity breaking, assuming only one  $\lambda'_{ijk} \neq 0$ .

It would be more natural to assume that there is only one large  $R_p$  coupling in the *weak* basis, i.e., only one  $\bar{\lambda}_{ijk} \neq 0$ . As we have indicated, this generates many couplings with different flavor structure in the *mass* basis, e.g., many  $\lambda'_{imn}$ 's. It is possible that

$$\lambda'_{imn} \simeq \bar{\lambda}_{ijk} V_{KMjm} \delta_{kn} \quad (9)$$

This will be the case if, for example, the rotation to the mass basis occurs only for the charge  $+2/3$  quark sector. Then, in addition to the Feynman diagrams that contribute to the flavor changing neutral current processes when only one  $\lambda'_{ijk}$  is present, there are new contributions involving the  $\lambda'_{imn}$  ( $m \neq j, n = k$ ) vertices. However, these new contributions interfere constructively with the operators that are present in the effective Lagrangian that is generated when there is only one non-zero  $\lambda'_{ijk}$ . So if these more natural assumptions are made, any constraint found for  $\bar{\lambda}_{ijk}$  is slightly better than the constraint that is obtained when only one  $\lambda'_{ijk}$  is present.

It would seem that the flavor changing neutral current processes may be rotated away by making a different physical assumption concerning which  $R_p$  coupling is non-zero. For example, while leaving the quark fields in the mass

basis, Equation (3) gives

$$W_{R_p} = \lambda'_{ijk}(N_i(V_{KM})_{jl}D_l - E_i U_j)D_k^c \quad (10)$$

$$= (\lambda'_{ijk}V_{KMjl})(N_i D_l - E_i(V_{KMlp}^{-1})U_p)D_k^c \quad (11)$$

$$= \tilde{\lambda}_{ijk}(N_i D_j - E_i(V_{KMjp}^{-1})U_p)D_k^c \quad (12)$$

where

$$\tilde{\lambda}_{ijk} \equiv \lambda'_{imk}(V_{KM})_{mj} \quad (13)$$

With the assumption that the  $\lambda'_{ijk}$  coefficients have values such that only one  $\tilde{\lambda}_{ijk}$  is non-zero, there is only one interaction of the form  $N_L D_L D^c$ . There is then no longer any flavor violation in the down quark sector. In particular, there are no  $R_p$  contributions to the processes discussed below. But now there are couplings involving each of the three up type quarks. So these interactions contribute to FCNC in the up sector; for example,  $D^0-\bar{D}^0$  mixing. We use  $D^0-\bar{D}^0$  mixing to place constraints on  $R$ -parity violation assuming only one  $\tilde{\lambda}_{ijk} \neq 0$ . Thus, there is no basis in which FCNC can be avoided in both sectors.

## 2.1 $K^0-\bar{K}^0$ Mixing

With one  $\lambda'_{ijk} \neq 0$ , the interactions of Equation (8) involve down and strange quarks. So, there are contributions to  $K^0-\bar{K}^0$  mixing through the box diagrams shown in Figure (1).




Figure 1:  $\mathcal{R}_p$  contributions to  $K^0-\bar{K}^0$  mixing with one  $\lambda'_{ijk} \neq 0$ . Arrows indicate flow of propagating left handed fields.

Evaluating these diagrams at zero external momentum and neglecting the down quark masses, the following effective Hamiltonian is generated

$$\mathcal{H}_{\mathcal{R}_p}^{\Delta S=2} = \frac{1}{128\pi^2} |\lambda'_{ijk}|^4 \left( \frac{1}{m_{\tilde{\nu}_i}^2} + \frac{1}{m_{\tilde{d}_{Rk}}^2} \right) ((V_{KM})_{j2}(V_{KM})_{j1}^*)^2 (\bar{d}_L \gamma^\mu s_L)^2 \quad (14)$$

where  $m_{\tilde{\nu}_i}$  is the sneutrino mass and  $m_{\tilde{d}_{Rk}}$  is the right-handed down squark mass. As this operator is suppressed by the CKM angles, it is largest when  $\lambda'_{ijk}$  is non-zero for  $j = 1$  or  $j = 2$ .

The SM effective Hamiltonian is [17]

$$\mathcal{H}_{SM}^{\Delta S=2} = \frac{G_F^2}{4\pi^2} m_c^2 ((V_{KM})_{12}(V_{KM})_{11}^*)^2 (\bar{d}_L \gamma^\mu s_L)^2 \quad (15)$$

where the CKM suppressed top quark contribution, the up quark mass, and QCD radiative corrections have been ignored. As the uncertainty in hadronic matrix elements of the Standard Model effective Hamiltonian are at most 40%, a conservative constraint on the  $\mathcal{R}_p$  coupling is obtained by demanding that  $\mathcal{L}_{\mathcal{R}_p}^{\Delta S=2} \leq 0.5 \mathcal{L}_{SM}^{\Delta S=2}$ . This gives the constraint

$$|\lambda'_{ijk}| \leq 0.08 \left( \frac{1}{z_i^2} + \frac{1}{w_k^2} \right)^{-\frac{1}{4}} \quad (16)$$

where  $z_i = m_{\bar{\nu}_i}/(100 \text{ GeV})$  and  $w_k = m_{\bar{d}_{Rk}}/(100 \text{ GeV})$ . This constraint applies for  $j = 1$  or  $j = 2$  and for any  $i$  or  $k$ . The constraint for  $j = 3$  is not interesting as the CKM angles suppress the  $\mathcal{R}_p$  operator relative to the Standard Model operator.

## 2.2 $B^0-\bar{B}^0$ Mixing

The  $\mathcal{R}_p$  interactions also contribute to both  $B^0-\bar{B}^0$  mixing and  $B_s^0-\bar{B}_s^0$  mixing through box diagrams similar to those given in the previous section. As  $B_s^0-\bar{B}_s^0$  mixing is expected to be nearly maximal, it is not possible at present to place a constraint on any non-Standard Model effects that would *add* more mixing. However,  $B^0-\bar{B}^0$  mixing has been observed [18] with a moderate  $x_d \sim 0.7$  [13]. As lattice QCD calculations predict  $B_K \sim 0.6$  [19] and  $B_B \sim 1.2$  [20], it is reasonable to expect that any  $\mathcal{R}_p$  contributions to  $B^0-\bar{B}^0$  mixing should not exceed 50% of the amount expected from the Standard Model alone.

The effective Hamiltonian generated by these  $\mathcal{R}_p$  processes is

$$\mathcal{H}_{\mathcal{R}_p} = \frac{1}{128\pi^2} |\lambda'_{ijk}|^4 \left( \frac{1}{m_{\bar{\nu}_i}^2} + \frac{1}{m_{\bar{d}_{Rk}}^2} \right) ((V_{KM})_{j3}(V_{KM})_{j1}^*)^2 (\bar{d}_L \gamma^\mu b_L)^2 \quad (17)$$

This is largest when  $\lambda'_{i3k}$  is non-zero.

The dominant contribution to  $B^0-\bar{B}^0$  mixing in the Standard Model is [21]

$$\mathcal{H}_{SM}^{\Delta S=2} = \frac{G_F^2 m_t^2}{4\pi^2} ((V_{KM})_{33}(V_{KM})_{31}^*)^2 G(x_t) (\bar{d}_L \gamma^\mu b_L)^2 \quad (18)$$




Figure 2:  $\tilde{H}_p$  contribution to  $K^+ \rightarrow \pi^+ \nu \bar{\nu}$  with one  $\lambda'_{ijk} \neq 0$ .

where  $x_t = m_t^2/m_W^2$ , and

$$G(x) = \frac{4 - 11x + x^2}{4(x-1)^2} - \frac{3x^2 \ln x}{2(1-x)^3} \quad (19)$$

For a top mass of 176 GeV,  $G(x_t) = 0.54$ .

This gives the constraint

$$|\lambda'_{i3k}| \leq 0.77 \left( \frac{1}{z_i^2} + \frac{1}{w_k^2} \right)^{-\frac{1}{4}} \quad (20)$$

with  $z_i$  and  $w_k$  as previously defined.

In addition to inducing  $B^0-\bar{B}^0$  mixing, these interactions also contribute to the  $b \rightarrow s + \gamma$  amplitude. However, with reasonable values for squark and sneutrino masses, the constraint is significantly weaker than that found from the top quark analysis.

### 2.3 $K^+ \rightarrow \pi^+ \nu \bar{\nu}$

The tree level Feynman diagram in Figure (2) generates an effective Hamiltonian which contributes to the branching ratio for  $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ . Using a Fierz

rearrangement, a straightforward evaluation of this diagram gives

$$\mathcal{H}_{\mathcal{R}_p} = \frac{1}{2} \frac{|\lambda'_{ijk}|^2}{m_{\tilde{d}_{Rk}}^2} (V_{KMj1} V_{KMj2}^*) (\bar{s}_L \gamma^\mu d_L) (\bar{\nu}_{Li} \gamma_\mu \nu_{Li}) \quad (21)$$

There is also a Standard Model contribution to this decay [21]. This is an order of magnitude lower than the existing experimental limit. To obtain a bound on the  $\mathcal{R}_p$  coupling, we shall assume that the  $\mathcal{R}_p$  effects dominate the decay rate.

As the matrix element for this semi-leptonic decay factors into a leptonic and a hadronic element, the isospin relation

$$\langle \pi^+(\mathbf{p}) | \bar{s} \gamma_\mu d | K^+(\mathbf{k}) \rangle = \sqrt{2} \langle \pi^0(\mathbf{p}) | \bar{s} \gamma_\mu u | K^+(\mathbf{k}) \rangle \quad (22)$$

can be used to relate  $\Gamma[K^+ \rightarrow \pi^+ \nu \bar{\nu}]$  to  $\Gamma[K^+ \rightarrow \pi^0 \nu e^+]$ . The effective Hamiltonian for the neutral pion decay channel arises from the spectator decay of the strange quark. It is

$$\mathcal{H}_{eff} = \frac{4G_F}{\sqrt{2}} V_{KM12}^* (\bar{s}_L \gamma^\mu u_L) (\bar{\nu}_{Li} \gamma_\mu e_{Li}) \quad (23)$$

So in the limit where the lepton masses can be neglected,

$$\frac{\Gamma[K^+ \rightarrow \pi^+ \nu_i \bar{\nu}_i]}{\Gamma[K^+ \rightarrow \pi^0 \nu e^+]} = \left( \frac{|\lambda'_{ijk}|^2}{4G_F m_{\tilde{d}_{Rk}}^2} \right)^2 \left( \frac{|V_{KMj1} V_{KMj2}^*|}{|V_{KM12}^*|} \right)^2 \quad (24)$$

This ratio is valid for  $i = 1, 2$  or  $3$ , since in the massless neutrino and electron approximation, the integrals over phase space in the numerator and denominator cancel. So using  $BR[K^+ \rightarrow \pi^+ \nu \bar{\nu}] \leq 5.2 \times 10^{-9}$  [22] (90%CL) and  $BR[K^+ \rightarrow$

$\pi^0\nu e^+$ ] = 0.0482 [13], the constraint is

$$|\lambda'_{ijk}| \leq 0.012 \left( \frac{m_{\tilde{d}_{Rk}}}{100 \text{ GeV}} \right) (90\%CL) \quad (25)$$

for  $j = 1$  or  $j = 2$ . Using  $|V_{KM13}| \geq 0.004$  [13] and  $|V_{KM23}| \geq 0.03$  [13], a conservative upper bound for  $\lambda'_{i3k}$  is

$$|\lambda'_{i3k}| \leq 0.52 \left( \frac{m_{\tilde{d}_{Rk}}}{100 \text{ GeV}} \right) (90\%CL) \quad (26)$$

## 2.4 $D^0-\bar{D}^0$ Mixing

If there is only one  $\tilde{\lambda}_{ijk}$  in the *mass* basis, then from Equation (12) it is clear that flavor changing neutral current processes will occur in the charge +2/3 quark sector. Rare processes such as  $D^0-\bar{D}^0$  mixing,  $D^0 \rightarrow \mu^+\mu^-$  and  $D^+ \rightarrow \pi^+l^+l^-$ , for example, may be used to place tight constraints on  $\tilde{\lambda}_{ijk}$ . For illustrative purposes, in this section we will consider  $D^0-\bar{D}^0$  mixing.

The interactions in Equation (12) generate box diagrams identical to those discussed in the previous sections if both the internal sneutrino (neutrino) propagators are replaced with slepton (lepton) propagators and the external quarks lines are suitably corrected. Using the same approximations that were made earlier, the  $\mathcal{H}_p$  effects generate the following effective Hamiltonian

$$\mathcal{H}_{\mathcal{H}_p} = \frac{1}{128\pi^2} |\tilde{\lambda}_{ijk}|^4 \left( \frac{1}{m_{\tilde{l}_i}^2} + \frac{1}{m_{\tilde{d}_{Rk}}^2} \right) ((V_{KM})_{2j}(V_{KM})_{1j}^*)^2 (\bar{c}_L \gamma^\mu u_L)^2 \quad (27)$$

$$\equiv G(\tilde{\lambda}_{ijk}, m_{\tilde{l}_i}, m_{\tilde{d}_{Rk}}) (\bar{c}_L \gamma^\mu u_L)^2 \quad (28)$$

In the vacuum saturation approximation, the  $R_p$  effects contribute an amount

$$(\Delta m)_{th} \equiv m_{D_L} - m_{D_S} = \frac{2}{3} f_D^2 m_D \text{Re} G(\tilde{\lambda}_{ijk}, m_{\tilde{l}_i}, m_{\tilde{d}_{Rk}}) \quad (29)$$

to the  $D_L - D_S$  mass difference. With  $f_D = 200 \text{ MeV}$  [23],  $m_D = 1864 \text{ MeV}$  [13], and  $|(\Delta m)_{exp}| \leq 1.32 \times 10^{-10} \text{ MeV}$  [13](90%CL), the constraint on  $\tilde{\lambda}_{ijk}$  for  $j = 1$  or  $j = 2$  is

$$|\tilde{\lambda}_{ijk}| \leq 0.16 \left( \left( \frac{100 \text{ GeV}}{m_{\tilde{l}_i}} \right)^2 + \left( \frac{100 \text{ GeV}}{m_{\tilde{d}_{Rk}}} \right)^2 \right)^{-\frac{1}{4}} \quad (90\%CL) \quad (30)$$

### 3 Top Quark Decay

In the Standard Model, the dominant decay mode for the top quark is

$$t \rightarrow b + W \quad (31)$$

with a real  $W$  gauge boson produced. This has a partial decay width

$$\Gamma[t \rightarrow W + b] = \frac{G_F m_t^3}{8\pi\sqrt{2}} |V_{tb}|^2 (1 - x_W^2)(1 - 2x_W^4 + x_W^2) \quad (32)$$

where  $x_W = m_W/m_t$ . The  $b$  quark mass has been neglected.

The  $R$ -parity violating interactions (see Equation (8) with  $j = 3$ )  $\lambda'_{i3k} \tilde{e}_L^i \tilde{d}_{Rk}^k t_L$  contribute to the decay  $t_L \rightarrow \tilde{l}_i^+ + d_{Rk}$  at tree level [24], if kinematically allowed. This is possible only if there exist sleptons lighter than the top quark. The partial width for this process is

$$\Gamma[t \rightarrow \tilde{l}_i^+ + d_k] = \frac{|\lambda'_{i3k}|^2 m_t (1 - y_i^2)^2}{32\pi} \quad (33)$$

with  $y_i \equiv m_{\tilde{l}_i}/m_t$  [24]. The mass of the down type quark has been neglected. If this is the only non-zero  $R$ -parity coupling, the two top quark decay channels are  $t \rightarrow b + W$  and  $t \rightarrow d_{Rk} + \tilde{l}_i^+$ , with branching fractions  $1 - x$  and  $x$ , respectively.

We assume that the Lightest Supersymmetric Particle (LSP), denoted by  $\tilde{\chi}^0$ , is neutral and that the real slepton decays with 100% branching fraction to the  $\tilde{\chi}^0$  and a lepton. The presence of a non-zero  $R$ -parity breaking coupling implies that the  $\tilde{\chi}^0$  is no longer stable [1]. The two dominant decays are [24]  $\tilde{\chi}^0 \rightarrow \nu_i + b + \bar{d}_k$  and  $\tilde{\chi}^0 \rightarrow \bar{\nu}_i + \bar{b} + d_k$ . The LSP decays inside the detector if [7]

$$|\lambda'_{i3k}| \geq 6 \times 10^{-5} \sqrt{\gamma} \left( \left( \frac{100 \text{ GeV}}{m_{\bar{d}_{Rk}}} \right)^2 + \left( \frac{100 \text{ GeV}}{m_{\bar{b}_R}} \right)^2 \right) \left( \frac{100 \text{ GeV}}{m_{\tilde{\chi}^0}} \right)^{5/2} \quad (34)$$

where  $\gamma$  is the Lorentz boost factor of  $\tilde{\chi}^0$ . For this decay chain to be kinematically allowed, we require that  $m_{\tilde{\chi}^0} \geq m_b$  for  $k = 1$  or  $k = 2$ , and  $m_{\tilde{\chi}^0} \geq 2m_b$  for  $k = 3$ . Using the previous equation, the maximum lower bound on  $\lambda'_{i3k}$  such that the LSP decays inside the detector is  $0.0003 \times \sqrt{\gamma}$  for  $k = 3$ , and  $0.002 \times \sqrt{\gamma}$  for  $k = 1$  or  $k = 2$ ; all for 300 GeV squark masses. We shall assume that  $\lambda'_{i3k}$  is larger than this value so that the LSP decays within the detector.

If a top quark decays through this  $R$ -parity violating process, the final state will contain one lepton, at least one  $b$  quark and missing transverse energy. The two novel features of this decay channel are that it spoils lepton universality and, when  $k = 3$ , produces a surplus of  $b$  quark events. Both of these signatures can be used to test the strength of  $R$ -parity violation.

The CDF collaboration reconstructs  $t\bar{t}$  quark events from observing: (1) dilepton (electron or muon) events coming from the leptonic decays of both the  $W$ 's; or (2) one lepton event arising from leptonic decay of one  $W$  and jets from the hadronic decay of the remaining  $W$  boson. CDF also requires a  $b$ -tag in the lepton+jets channel. If the lightest slepton has a mass between 50 and 100  $GeV$ , then the kinematics of the decay  $\tilde{l}_i \rightarrow \tilde{\chi}^0 + l_i$  will be similar to that of the leptonic decay of the  $W$  boson. A slepton of mass less than 45  $GeV$  is ruled out by the LEP limit on the  $Z$  decay width [13]. If the slepton mass is close to the top mass, then the  $b$  quark produced in the top decay via this channel will have less energy than the  $b$  quark from the top decay via the SM channel. Also, the lepton from the slepton decay will have more energy than the lepton from the  $W$  decay. These will affect the lepton and the  $b$  quark detection efficiencies. Although these decay channels will be present for any slepton lighter than the top quark, for the purpose of obtaining a constraint, we shall assume that there is a slepton with a mass in the range given above. The presence of the  $R$ -parity violating coupling will then contribute signals to all of these channels.

We assume that the  $i = 1$  coupling is non-zero. However, all that is required is that the slepton in the generation with the non-zero coupling have a mass in the range quoted above, i.e., if  $\lambda'_{13k} \neq 0$  then we require  $50 GeV < m_{\tilde{e}} < 100 GeV$ , and if  $\lambda'_{23k} \neq 0$  then we require  $50 GeV < m_{\tilde{\mu}} < 100 GeV$ . Assuming also that the CDF data is consistent with lepton universality, the constraints we

obtain for  $\lambda'_{13k}$  and  $\lambda'_{23k}$  are identical.

In the  $k = 1, 2$  cases, two  $b$  quarks are always produced in a  $t\bar{t}$  event. In the  $k = 3$  case, the LSP decays into  $\bar{b}b\nu_i$  or  $\bar{b}b\bar{\nu}_i$ . Thus, four or six  $b$  quarks may be produced if one or both of the top quarks decay through the  $R$ -parity breaking channel; this possibility must be treated separately.

### 3.1 $\lambda'_{i3k}, k \neq 3$

The branching fraction for the di-electron event is

$$BR[t\bar{t} \rightarrow ee + X] = x^2 + L^2(1-x)^2 + 2Lx(1-x) \quad (35)$$

with  $L =$  leptonic branching fraction of  $W$ , approximately  $1/9$ . The first term arises from both top quarks decaying via the  $R$ -parity violating interaction; the second is the Standard Model contribution; and the third is the contribution from one top quark decaying through the  $R$ -parity breaking channel and the other top quark decaying through the Standard Model channel. The other branching fractions are

$$BR[t\bar{t} \rightarrow \mu\mu + X] = L^2(1-x)^2 \quad (36)$$

$$BR[t\bar{t} \rightarrow \mu e + X] = 2(1-x)^2L^2 + 2x(1-x)L \quad (37)$$

$$BR[t\bar{t} \rightarrow \mu + \text{jets}] = 2(1-x)^2L(1-3L) \quad (38)$$

$$BR[t\bar{t} \rightarrow e + \text{jets}] = 2(1-x)^2L(1-3L) + 2x(1-x)(1-3L) \quad (39)$$

The factor of  $1 - 3L$  is the hadronic branching fraction of the  $W$  boson. We have also assumed that the branching fraction for  $\tilde{l} \rightarrow l + \tilde{\chi}^0$  is close to one. We are ignoring leptonic events produced from the Standard Model decay of the  $W$  boson into  $\tau\nu_\tau$ .

Two independent constraints on the  $\tilde{R}_p$  interactions may be obtained from the top quark data. CDF has observed the  $t\bar{t}$  cross section to be  $\sigma(t\bar{t})_{exp} = 6.8^{+3.6}_{-2.4}$  pb [12]. The QCD calculation [25] gives the value  $\sigma(t\bar{t})_{th} = 4.79^{+0.67}_{-0.41}$  pb for  $m_t = 176 \text{ GeV}$ .

The first method is to compare the ratio of theoretically predicted values for the numbers of events found in two channels with the experimentally observed ratio. For example,  $\sigma(t\bar{t})_{th} \times BR[t\bar{t} \rightarrow \mu + \text{jets}] \times \int L dt \times (\text{detection efficiencies})$  is the number of  $\mu + \text{jets}$  events that should have been observed where  $\int L dt$  is the integrated luminosity. This theoretical prediction contains uncertainties in both the value for the  $t\bar{t}$  production cross section and in the lepton and the  $b$  quark detection efficiencies. In comparing the ratio

$$(\sigma(t\bar{t})_{th} \times BR[t\bar{t} \rightarrow e + \text{jets}]) / (\sigma(t\bar{t})_{th} \times BR[t\bar{t} \rightarrow \mu + \text{jets}]) \quad (40)$$

the uncertainties in the  $t\bar{t}$  cross section cancel. The  $b$ -detection efficiencies also cancel. If the electron and the muon detection efficiencies in the lepton + jets channel are equal, these uncertainties will also cancel. The only remaining errors are statistical. The CDF collaboration reported observing 37  $b$ -tagged events in

the lepton  $+ \geq 3$  jets channel. In this set there were 50  $b$ -tags, with a background of 22  $b$ -tags. A conservative estimate for the background in the 37 events is 22. This leaves 15  $t\bar{t}$  events in the lepton +jets channel. Since no inconsistencies with electron-muon universality have been reported, a central value of 7  $\mu$  +jets and 7  $e$  +jets events will be assumed. This leads to

$$\frac{BR[t\bar{t} \rightarrow e + \text{jets}]_{th}}{BR[t\bar{t} \rightarrow \mu + \text{jets}]_{th}} = \frac{\#(e + \text{jets events})}{\#(\mu + \text{jets events})} = 1_{-b}^{+a} \quad (41)$$

Inserting the theoretical predictions for the branching ratios leads to the constraint  $x < La/(1 + La)$ , where  $a$  is the uncertainty in the previous ratio. In this case,  $a = b = 1/\sqrt{7}$ . This gives  $x < 0.077$  at 95% $CL$  which leads to

$$|\lambda'_{13k}| \leq 0.41 \text{ (95\% } CL) \quad (42)$$

for  $k = 1$  or  $k = 2$  and a slepton of mass 100  $GeV$ .

A similar analysis may be performed for the dilepton channels. In principle these channels should lead to a good constraint since a non-zero  $\lambda'_{13k}$  coupling will lead to an excess of electrons observed in the di-electron channel over the number of muons observed in the di-muon channel. However at present only a small number of dilepton events have been observed and an interesting constraint cannot be obtained.

In the other method we will compare the number of events produced in a given channel with the theoretical expectation. The number of produced events is  $\sigma[t\bar{t}]_{th} \times BR[t \rightarrow l + \text{jets}]_{th} \times \int L dt$ . Here  $\sigma[t\bar{t}]_{th}$  is the production cross section

calculated in perturbative QCD for the assumed top quark mass of 176 GeV. We will use the fact that the number of experimentally observed events in any given channel is consistent with, within experimental errors, the number expected in the standard model. The actual number of events detected depends upon the detection efficiency. We will use the number of observed events in any channel to determine the statistical accuracy with which the rate in that channel is measured, and then constrain the strength of the  $R_p$  terms by requiring that the rate is not changed by more than the error.

This leads to the constraint

$$\frac{BR[t\bar{t} \rightarrow l + \text{jets}, x]_{th}}{BR[t\bar{t} \rightarrow l + \text{jets}, x = 0]_{th}} = \frac{\sigma[t\bar{t}]_{exp}}{\sigma[t\bar{t}]_{th}} \quad (43)$$

within theoretical and experimental errors. Using the theoretical and experimental values for the production cross sections [12, 25] leads to

$$\epsilon^2 \leq \frac{BR[t\bar{t} \rightarrow l + \text{jets}, x]_{th}}{BR[t\bar{t} \rightarrow l + \text{jets}, x = 0]_{th}} \leq 1 + d \quad (44)$$

with  $\epsilon = 0.9$  and  $d = 1.37$ . The constraint on  $x$  is then

$$x \leq \min \left( 1 - \epsilon, \frac{1 - 2L - \sqrt{(1 - 2L)^2 - 4Ld(1 - L)}}{2(1 - L)} \right) \quad (45)$$

The first entry is the constraint from the  $\mu + \text{jets}$  channel and the second entry is from the  $e + \text{jets}$  channel. For these values of  $\epsilon$  and  $d$ , the constraint is  $x \leq 0.1$ . For a 100 GeV slepton this translates into the constraint

$$|\lambda'_{13k}| \leq 0.46 \quad (46)$$

for  $k = 1$  or  $k = 2$ .

### 3.2 $\lambda'_{i33}$

For this coupling the analysis of the previous section must be modified in the lepton + jets channel since the  $b$ -detection efficiencies no longer cancel. This is because in the  $R$ -parity breaking decay channel three  $b$  quarks are produced. To correct for this, introduce the function  $P(k, n)$  that gives the probability that, given that  $n$   $b$  quarks are produced,  $k$  of them are detected. Then the number of observed single  $b$  quark events expected in the  $e$ +jets channel is

$$\begin{aligned} \#(e + \text{jets events}) &= \left(2(1-x)^2L(1-3L)P(1,2) + 2x(1-x)(1-3L)P(1,4)\right) \\ &\quad \times \mathcal{N} \end{aligned} \tag{47}$$

where

$$\mathcal{N} \equiv \int Ldt \times \sigma(t\bar{t})_{th} \tag{48}$$

With  $P(1,2) \leq P(1,n)$  for  $n \geq 2$ , then

$$\#(e + \text{jets events}) \geq \left(2(1-x)^2L(1-3L) + 2x(1-x)(1-3L)\right) P(1,2) \times \mathcal{N} \tag{49}$$

These approximations will give a conservative limit for  $\lambda'_{133}$ . The analysis of the previous section may now be carried out with the following restrictions:

(i) In comparing the ratio of the numbers of events detected in two channels

with the theoretical prediction, the inequality in Equation (49) indicates that only upper limit in Equation (41) may be used;

(ii) In comparing the number of events detected in a channel with the theoretically predicted value for that channel, only the upper bound in Equation (44) may be used in the  $e$ +jets channel, and either limit may be used in the  $\mu$ + jets channel. With these caveats, a conservative limit on the branching fraction for  $t \rightarrow b + \tilde{l}_i^+$  is then

$$x \leq \min \left( La/(1+La), 1 - \epsilon, \frac{1 - 2L - \sqrt{(1 - 2L)^2 - 4Ld(1 - L)}}{2(1 - L)} \right) \quad (50)$$

For the errors quoted in the previous section, the result is

$$|\lambda'_{133}| \leq 0.41 \text{ (95\% CL)} \quad (51)$$

As the  $R$ -parity breaking decay channels produce three  $b$  quarks, then for moderate values of  $\lambda'_{133}$  or  $\lambda'_{233}$ , semi-leptonic events containing four and six  $b$  quarks should be observable at the Tevatron. The non-observance of these events should provide the strongest test for the  $R$ -parity breaking couplings  $\lambda'_{133}$  or  $\lambda'_{233}$ . If limits on the branching fractions for the  $t\bar{t}$  pair to decay into these excess  $b$  quark channels are known, then the  $R$ -parity branching fraction  $x$  is constrained.

Namely,

$$1. \quad BR[t\bar{t} \rightarrow X + \geq 3b's] \leq B_1 \Rightarrow x \leq \left( 1 - \sqrt{1 - B_1} \right) \quad (52)$$

$$2. \quad BR[t\bar{t} \rightarrow X + \geq 3b's + 2e] \leq B_2 \Rightarrow x \leq \frac{\sqrt{L^2 + B_2(1 - 2L)} - L}{1 - 2L} \quad (53)$$

$$3. \quad BR[t\bar{t} \rightarrow X + \geq 6b's + 2e] \leq B_3 \Rightarrow x \leq \sqrt{B_3} \quad (54)$$

$$4. \quad BR[t\bar{t} \rightarrow X + \geq 3b's + e] \leq B_4 \Rightarrow x \leq \frac{1}{2} \left( 1 - \sqrt{1 - \frac{2B_4}{1 - 3L}} \right) \quad (55)$$

$$(56)$$

This constrains  $|\lambda'_{133}|$ . To constrain  $|\lambda'_{233}|$ , interchange  $e$  with  $\mu$  in the previous equations.

The constraints on  $|\lambda'_{133}|$  and  $|\lambda'_{233}|$  found in this section are comparable to those obtained from examining  $\mathcal{R}_p$  contributions either to  $Z \rightarrow b\bar{b}$  and  $Z \rightarrow l^+l^-$  decays [9] or to forward-backward asymmetry measurements in  $e^+e^-$  collisions [5]. We have engaged in this exercise to illustrate how comparable  $\mathcal{R}_p$  constraints may be obtained from analysing top quark decays even though the experimental and theoretical errors are still large. These processes will provide much better tests of  $R$ -parity violation once more top quark decays are seen.

## 4 Summary

In this paper we have argued that  $R$ -parity breaking interactions always lead to flavor changing neutral current processes. It is possible that there is a single  $\mathcal{R}_p$  coupling in the charge  $+2/3$  quark sector. But requiring consistency with electroweak symmetry breaking demands that  $\mathcal{R}_p$  couplings involving all the charge  $-1/3$  quarks exist. That is, a single coupling scheme may only be possible

in either the charge  $2/3$  or the charge  $-1/3$  quark sector, but not both. As a result, flavor changing neutral current processes always exist in one of these sectors. We have used  $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ ,  $K^0 - \bar{K}^0$  mixing,  $B^0 - \bar{B}^0$  mixing and  $D^0 - \bar{D}^0$  mixing to constrain the  $R_p$  couplings. The constraints we obtain for the first two generations are more stringent than those presently existing in the literature.

The  $R$ -parity breaking interactions lead to the top quark decay  $t \rightarrow \tilde{l}_i + d_k$ , if the slepton is lighter than the top quark. Some of the new top quark decays spoil electron-muon universality or result in  $t\bar{t}$  events with more than 2  $b$  quarks. At present, the CDF collaboration has not reported any inconsistencies with lepton universality or reported any events with more than 2  $b$  quarks. These decays also lower the branching fractions for Standard Model top quark decays. These observations are used to constrain some  $R_p$  couplings.

A list of the known constraints on the  $\lambda'_{ijk}$  couplings is presented in Table (1). Although several of these couplings are constrained by different low energy processes, we have only listed the smallest known upper limit.

The tightest constraint is on  $|\lambda'_{ijk}|$  for  $j = 1, 2$  and any  $i$  and  $k$ . This comes from the rare decay  $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ . With the exception of  $\lambda'_{133}$ , the constraints on the third quark generation couplings are only of order  $e/\sin\theta_w$ . Once more top quark decays are observed the signatures discussed in this paper will more

$ \lambda'_{1jk} $		$ \lambda'_{2jk} $		$ \lambda'_{3jk} $	
111	0.012 <sup>a</sup>	211	0.012 <sup>a</sup>	311	0.012 <sup>a</sup>
112	0.012 <sup>a</sup>	212	0.012 <sup>a</sup>	312	0.012 <sup>a</sup>
113	0.012 <sup>a</sup>	213	0.012 <sup>a</sup>	313	0.012 <sup>a</sup>
121	0.012 <sup>a</sup>	221	0.012 <sup>a</sup>	321	0.012 <sup>a</sup>
122	0.012 <sup>a</sup>	222	0.012 <sup>a</sup>	322	0.012 <sup>a</sup>
123	0.012 <sup>a</sup>	223	0.012 <sup>a</sup>	323	0.012 <sup>a</sup>
131	0.26 <sup>c</sup>	231	0.22 <sup>d</sup>	331	0.26 <sup>e</sup>
132	0.4 <sup>b</sup>	232	0.4 <sup>b</sup>	332	0.26 <sup>e</sup>
133	0.001 <sup>f</sup>	233	0.4 <sup>b</sup>	333	0.26 <sup>e</sup>

Table 1: Constraints on  $|\lambda'_{ijk}|$  from:(a)  $K^+ \rightarrow \pi^+ \nu \bar{\nu}$  (90%*CL*); (b) top quark decay (95%*CL*); (c) atomic parity violation and  $eD$  asymmetry (90%*CL*) [5]; (d)  $\nu_\mu$  deep-inelastic scattering (95%*CL*) [5]; (e) partial  $Z^0$  decay width (95%*CL*) [9]; (f)  $\nu_e$  mass (90%*CL*) [6]. All limits are for 100 *GeV* sparticle masses.

tightly constrain these couplings.

## 5 Acknowledgements

The authors would like to acknowledge both I. Hinchliffe and M. Suzuki for many useful discussions. This work was supported in part by the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, Division of High Energy Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098 and in part by the National Science Foundation under grant PHY-90-21139. M.G. would also like to thank the Natural Sciences and Engineering Research Council of Canada (NSERC) for their support.

## References

- [1] For reviews of supersymmetry and supersymmetry phenomenology, see:  
P. Fayet and S. Ferrara, *Phys. Rep.* **5**, 249 (1977); H.P. Nilles, *Phys. Rep.* **110**, 1 (1984); M.F. Sohnius, *Phys. Rep.* **128**, 2 (1985); I. Hinchliffe, *Ann. Rev. Nucl. Part. Sci.* **36**, 505 (1986).
- [2] L.J. Hall and M. Suzuki, *Nucl. Phys.* **B231**, 419 (1984).
- [3] C.E. Carlson, P. Roy and M. Sher, *Phys. Lett.* **B357**, 99 (1995).
- [4] S. Dimopoulos and L.J. Hall, *Phys. Lett.* **B207**, 210 (1987).

- [5] V. Barger, G.F. Giudice and T. Han, Phys. Rev. **D40**, 2987 (1989).
- [6] R. Godbole, P. Roy and X. Tata, Nucl. Phys. **B401**, 67 (1993).
- [7] S. Dawson, Nucl. Phys. **B261**, 297 (1985).
- [8] G. Bhattacharyya and D. Choudhury, Mod. Phys. Lett. **A10**, 1699 (1995).
- [9] G. Bhattacharyya, J. Ellis and K. Sridhar, Mod. Phys. Lett. **A10**, 1583 (1995).
- [10] R. Mohapatra, Phys. Rev. **D34**, 3457 (1986).
- [11] S. Abachi *et al.*, Phys. Rev. Lett. **74**, 2632 (1995).
- [12] F. Abe *et al.*, Phys. Rev. Lett. **74**, 2626 (1995).
- [13] *Review of Particle Properties*, Phys. Rev. **D50**, 1177 (1994).
- [14] E. Ma and P. Roy, Phys. Rev. **D41**, 988 (1990); E. Ma and D. Ng, Phys. Rev. **D41**, 1005 (1990).
- [15] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [16] R. Barbieri and A. Masiero, Nucl. Phys. **B267**, 679 (1986).
- [17] M.K. Gaillard and B.W. Lee, Phys. Rev. **D10**, 897 (1974).

- [18] C. Albajar *et al.*, Phys. Lett. **B186**, 247 (1987); H. Albrecht *et al.*, Phys. Lett. **B197**, 452 (1987).
- [19] C. Bernard, Nucl. Phys. (**Proc. Suppl.**) **B34**, 47 (1994); S. Sharpe, Lectures given at the Theoretical Advanced Study Institute in Particle Physics (TASI 94), Boulder, CO, 29 May - 24 Jun 1994.
- [20] G. Martinelli, in Proceedings of the 6th Rencontres De Bois, Bois, France 20 - 25 Jun 1994.
- [21] T. Inami and C.S. Lim, Prog. Theor. Phys. **65**, 292 (1981).
- [22] M.S. Atiya *et al.*, Phys. Rev. Lett. **70**, 2521 (1993); (erratum) M.S. Atiya *et al.*, Phys. Rev. Lett. **71**, 305 (1993).
- [23] J. Shigemitsu, in *Proceedings of the XXVII International Conference on High Energy Physics* (Glasgow, Scotland, UK, July 1994), edited by P.J. Bussey and I.G. Knowles (Institute of Physics Publishing, Bristol and Philadelphia, 1995).
- [24] H. Dreiner and R.J.N. Phillips, Nucl. Phys. **B367**, 591 (1991).
- [25] E. Laenen, J. Smith and W.L. van Neerven, Phys. Lett. **B321**, 254 (1994).

LAWRENCE BERKELEY NATIONAL LABORATORY  
UNIVERSITY OF CALIFORNIA  
TECHNICAL & ELECTRONIC INFORMATION DEPARTMENT  
BERKELEY, CALIFORNIA 94720