NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

How fast does the multiplicity of particle production in p-p collisions really increase with primary energy?

Erwin M. Friedlander

Lawrence Berkeley Laboratory, Berkeley, California 94720

December 1978

ABSTRACT

Multiplicity distributions of particles created in p-p collisions with E_{lab} = 50-200 GeV are shown to contain at least one Poisson component which becomes dominant at multiplicities $k \geqslant 1.5 \langle k \rangle$. The mean values of both this component and of the (relatively low multiplicity residue) increase like $E_{lab}^{1/2}$; the relative weight of the Poisson component decreases approximately like $E_{lab}^{-0.3}$. If these trends continue beyond the available accelerator energies, one might expect the mean multiplicity to approach an $E_{lab}^{1/2}$ law beyond 10 TeV.

How fast does the multiplicity of particle production in p-p collisions really increase with primary energy?

It has been well-known for a long time that the dependence of the mean multiplicity, m, of particles produced in high energy p-p collisions on the primary laboratory energy E_o is much weaker than $\sim E_o^{1/2}$, the fastest rise allowed kinematically or implied e.g by Heisenberg's original theory.[1] At the same time the rise is faster than $\sim \ln E_o$ as predicted by scaling models [2] and even somewhat faster than the $E_o^{1/4}$ dependence predicted by thermodynamical-hydrodynamical considerations [3], [4] implying a Lorentz contracted production volume.

This paper is concerned with the problem of predicting the evolution of m with E_o at super-high energies (say, beyond 10 TeV), not just from an extrapolation or fits to values of m observed in the energy range covered by present accelerators, but by using regularities observed in the structure of multiplicity distributions (abbreviated hereafter as MD), too. It turns out that—provided these regularities continue to hold at very high E_o—a very fast increase, $\sim E_o^{1/2}$ is asymptotically expected.

In order to emphasize the Poisson-like shape of MD's† it is preferable to use instead of the probability W(k) for observing k secondaries the quantity

$$Y(k) \equiv \ln[k!\ W(k)] \tag{1}$$

Obviously, if W(k) is a Poisson distribution (PD) of mean, say, a, Y is linear in k:

† Hereafter we will be concerned only with negative multiplicities $k = (n_{ch}-2)/2$ where n_{ch} is the total multiplicity of charged secondaries; k refers to created particles only.

$$Y = -a + k \ln a \tag{2}$$



Fig. 1

Fig. 1 shows the MD at E_o = 69 GeV [5] on a Y vs. k plot. The PD shape is obvious. However, with increasing energy the shape becomes more complicated.

Assume† that W(k) is a superposition of two MD's, say, W_1 and W_2 with means m_1 and $m_2 > m_1$.

$$W(k) = (1 - \alpha)\, W_1\,(k) + \alpha\, W_2\,(k) \tag{3}$$

where one component, say W_2, is Poisson and all we know about W_1 is that it practically dies out beyond some value k_c of k. Then, beyond k_c,

$$Y \approx - m_2 + \ln \alpha + \ln m_2 \quad ; \tag{4}$$

i.e., we get again a straight line with slope $\ln m_2$ but with an intercept depending on both m_2 <u>and</u> the relative weight α of the Poisson component.

† As has been done in Ref. [6] for mathematical expediency, but without presenting compelling evidence for Poisson components!



Fig. 2

This is well seen in fig. 2 which shows the MD at 300 GeV, [7] the statistically best measured sample to date (10^4 events).

The fact that this pattern is consistently repeated over the whole range of accelerator energies is shown in fig. 3, which displays the scaling variable Z, defined as

$$Z = \frac{m + y}{m \ln m} \quad , \tag{5}$$

as a function of

$$X \equiv k/m \quad , \tag{6}$$

for all available data from 50 to 2100 GeV. On such a plot PD would be a straight line of slope 1 passing through the origin.

As can be seen, beyond $x \sim 1.5$ the points (drawn from both HBC exposures between 50 and 405 GeV and ISR results between 500 and 2100 GeV equivalent laboratory energy) do indeed cluster along a straight line confirming up to the highest available energy the presence of a PD component (W_2). As to the residue (W_1), it can be shown that, in spite of the apparently high statistics gathered to this day, the accuracy is insufficient to define its shape. A PD is not excluded, although a more complicated structure



Fig. 3

may be indicated by the HBC-data.† However, the accuracy is just sufficient to estimate the mean values m_2 and m_1 (the Poisson component and the residue).

These values are plotted against E_o (dilog plot) in fig. 4, together with the overall mean m of k.

As can be seen, the energy dependence of both m_1 and m_2 can be well parameterized as $\sim E_o^{\delta}$ with δ close to 1/2 (the fitted values are $\delta = 0.54 \pm 0.03$ for m_2 and—understandably with lower accuracy—$\delta = 0.57 \pm 0.13$ for m_1).

It thus appears that p-p collisions can be regarded as a mixture of (at least) two types of events, each of which produces particles with a multiplicity law $\sim E_o^{1/2}$.

This can be reconciled with the relatively slow variation of the <u>mean of the mixture</u>, namely m (indeed, for m, $\delta = 0.35 \pm 0.01$, if it is parameterized in the same way as m_1 and m_2) only if α (the relative weight of W_2) <u>decreases</u> with E_o. Within the statistical

† The largest uncertainty resides in $W_1(0)$ because of the systematical effects connected with estimation of the elastic contribution.



Fig. 4

and systematic uncertainties of the data, α appears to decrease like $E_o^{-(0.3 \pm 0.1)}$. If this trend continues at very high energies, W_1 should become dominant beyond, say, 50 TeV (at ~ 400 GeV W_1 and W_2 have comparable weights) and there one might expect m to increase like $E_o^{1/2}$. This may be essential in understanding the development of extensive air showers in the early stages of their evolution.

This report was prepared under the auspices of the Nuclear Physics Division of the Department of Energy.

References

1. W. Heisenberg, Kosmische Strahlung, Springer, Berlin 1953, p. 148 ff.
2. R. P. Feynman, Phys. Rev. Lett. 23, 1415 (1969).
3. E. Fermi, Progr. Theor. Phys. 5, 270 (1950).
4. L. Landau, Izv. A. N. USSR, s. fiz. 17, 1 (1953).
5. V. Ammosov et al, Phys. Rev. Lett. 42B, 519 (1972).
6. J. Lach and E. Malamud, Phys. Lett. 44B, 474 (1973).
7. A. Firestone et al, Phys. Rev. D10, 2080 (1974).