



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

ENERGY & ENVIRONMENT DIVISION

Submitted to the Federation of European Biochemical
Societies, Vol. 91, No. 1, July 1978

pH GRADIENTS ACROSS THYLAKOID MEMBRANES MEASURED
WITH A SPIN-LABELED AMINE

A. T. Quintanilha and R. J. Mehlhorn

July 1978

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782*



DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

pH GRADIENTS ACROSS THYLAKOID MEMBRANES MEASURED WITH A SPIN-LABELED AMINE

A. T. QUINTANILHA and R. J. MEHLHORN


Membrane Bioenergetics Group, Lawrence Berkeley Laboratory, University of California, Berkeley, CA 94720, USA

Received 4 May 1978


1. Introduction

In their uncharged form amines have been shown to be permeable across membranes and to establish concentration gradients proportional to H^+ gradients [1,2]. In their protonated form they may bind to negative moieties on the surface of proteins or lipid layers. Amines have therefore been used to study pH gradients across membranes [2-9] as well as the surface charge of membranes [10-13]. In particular the quenching of 9-aminoacridine fluorescence has been used to measure the pH gradient across thylakoid membranes [4,8], while that changes in the fluorescence of acridines are claimed [10,12,13] to be closely associated with their strong interaction with the membrane surface and that transport across the thylakoid membrane may not be required.


When using amine probes to measure pH gradients across membranes it becomes important to distinguish between the free and membrane bound populations of the probe; we have therefore used the spin-labeled amine 4-amino-2,2,6,6-tetramethyl-piperidine-*N*-oxyl (Tempamine) with $pK = 9.5$, which shows distinct bound and free EPR signals. An impermeable analogue, 4-trimethyl ammonium-2,2,6,6-tetramethyl-piperidine-*N*-oxyl bromide (CAT_1) and the impermeable spin broadening agent $K_3Fe(CN)_6$ were used to demonstrate that it is the uncharged form of Tempamine that is permeable to the thylakoid membrane. The permeable uncharged spin label 2,2,6,6-tetramethyl-4-piperidone (^{15}N -Tempone) was used to show that at low concentrations of Tempamine ($150 \mu M$) virtually no swelling of the thylakoids occurred during illumination.



TEMPAMINE



CAT_1



^{15}N TEMPONE

Our studies have allowed us to determine the internal aqueous concentration of this amine in the thylakoid and to accurately measure the pH gradient across its membrane.

2. Materials and methods

Thylakoid membranes were prepared from spinach leaves in 0.4 M sucrose, 10 mM NaCl buffered with 10 mM tricine, at pH 8, and resuspended in this medium at 6 mg chl/ml [14]. For the present studies thylakoid suspensions (0.6 mg chl/ml) in 90 mM NaCl or KCl, tricine > 1 mM (pH 8), in the presence of spin label and 0.1 mM methylviologen were placed in quartz EPR tubes of 1 mm internal diameter. Where indicated, samples were illuminated by a 400 W tungsten halogen lamp (GE 3476) at heat filtered saturating light intensities ($\sim 200 W/m^2$) inside the microwave cavity and the spectra recorded in an E-109 E spectrometer at microwave power of 10 mW, time constant 0.128 s and scan time 9 min. $K_3Fe(CN)_6$, the spin broadening agent, was usually used at conc. ≤ 40 mM. Tempamine was from Aldrich Chemical Co., CAT_1 was synthesized [15] and ^{15}N -Tempone was a gift from A. D. Keith.

tion and the impermeability of CAT_1 across the thylakoid membrane. We have also been able to demonstrate that at low concentrations of Tempamine ($< 20 \mu M$) and fairly high concentrations of chlorophyll (0.6 mg chl/ml) virtually no binding of Tempamine to the thylakoid membrane is detectable either in the dark or in the light. Furthermore, this technique allows unambiguous determinations of inside and outside aqueous concentrations of Tempamine in a thylakoid suspension and therefore accurate determinations of the pH gradient across these membranes during illumination. Our results give ΔpH values of the order of 3.2 ± 0.2 in good agreement with the data in [22].

Acknowledgements

We acknowledge useful discussions with L. Packer. Research supported by the Department of Energy.

References

- [1] Packer, L. and Crofts, A. R. (1967) in: *Current Topics in Bioenergetics* (Sanadi, D. R. ed) pp. 23-64.
- [2] Deamer, D. W., Prince, R. C. and Crofts, A. R. (1972) *Biochim. Biophys. Acta* 274, 323-335.
- [3] Rottenberg, H., Grunwald, T. and Avron, M. (1972) *Eur. J. Biochem.* 25, 54-63.
- [4] Shuldiner, S., Rottenberg, H. and Avron, H. (1972) *Eur. J. Biochem.* 25, 64-70.
- [5] Rottenberg, H. and Grunwald, T. (1972) *Eur. J. Biochem.* 25, 71-74.
- [6] Portis, A. R. and McCarty, R. E. (1973) *Arch. Biochem. Biophys.* 156, 621-625.
- [7] Pick, U., Rottenberg, H. and Avron, M. (1975) in: *Proc. 3rd Int. Cong. Photosynthesis* (Avron, M. ed) pp. 967-974, Elsevier, Amsterdam.
- [8] De Benedetti, E. and Garlaschi, F. M. (1977) *J. Bioenerg. Biomembr.* 9, 195-201.
- [9] Rottenberg, H. and Lee, C. P. (1975) *Biochemistry* 14, 2675-2680.
- [10] Fiolet, J. W. T., Van der Erf-ter Haar, L., Kraayenhof, R. and Van Dam, K. (1975) *Biochim. Biophys. Acta* 387, 320-334.
- [11] Barber, J. and Searle, G. F. W. (1977) in *Abst. 4th Int. Cong. Photosynthesis*, p. 18.
- [12] Searle, G. F. W., Barber, J. and Mills, J. D. (1977) *Biochim. Biophys. Acta* 461, 413-425.
- [13] Kraayenhof, R. and Arents, J. C. (1977) in: *Electrical Phenomena at the Biological Membrane Level* (Roux, E. ed) pp. 493-504, Elsevier, Amsterdam.
- [14] Avron, M. (1961) *Anal. Biochem.* 2, 535-543.
- [15] Mehlhorn, R. J. and Packer, L. (1976) *Biochim. Biophys. Acta* 423, 382-397.
- [16] Mehlhorn, R. J. and Keith, A. D. (1972) in: *Membrane Molecular Biology* (Fox, C. F. and Keith, A. D. eds) pp. 192-227, Sinauer Ass. Inc.
- [17] Weaver, E. C. and Chen, H. P. (1966) *Science* 153, 301-303.
- [18] Torres-Pereira, J., Mehlhorn, R. J., Keith, A. D. and Packer, L. (1974) *Arch. Biochem. Biophys.* 160, 90-99.
- [19] Keith, A. D., Snipes, A. D., Mehlhorn, R. J. and Gunter, T. (1977) *Biophys. J.* 19, 205-218.
- [20] Morse, P. D. ii (1977) *Biochem. Biophys. Res. Commun.* 77, 1486-1491.
- [21] Chow, W. S. and Hope, A. B. (1976) *Aust. J. Plant Physiol.* 3, 141-152.
- [22] Rottenberg, H. (1975) *Bioenerg.* 7, 61-74.

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720