

Submitted to Physical Review Letters

LBL-990
Preprint

RECEIVED
LAWRENCE
BERNARD LABORATORY

DOCUMENTS SECTION

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

**GAUGE THEORY OF STRONG, WEAK, AND
ELECTROMAGNETIC INTERACTIONS**

I. Bars, M. B. Halpern, and M. Yoshimura

July 3, 1972

AEC Contract No. W-7405-eng-48



34

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

GAUGE THEORY OF STRONG, WEAK, AND ELECTROMAGNETIC INTERACTIONS*

I. Bars,† M. B. Halpern, and M. Yoshimura†§

Department of Physics and Lawrence Berkeley Laboratory
University of California, Berkeley, California 94720

July 3, 1972

ABSTRACT

Starting from the recent $U(3) \otimes U(3)$ gauge model of strong interactions, we discuss the very natural synthesis with Weinberg's model of leptons. Due to the interplay of the two models (a) neutral $\Delta S = 1$ currents are eliminated without enlarging the number of quarks, and (b) a natural $(3, \bar{3}) \oplus (\bar{3}, 3)$ symmetry breaking emerges for the hadrons. Incorporation of alternative lepton theories is mentioned.

Although there are now a number of renormalizable gauge models of leptons,¹ only one class of such theories has been constructed for hadrons.² We have asked the question: Can we effect a marriage between the hadron and lepton theories, such that, for simplicity, we have just $[(3, \bar{3}) \oplus (\bar{3}, 3) \text{ broken}] U(3) \otimes U(3)$ (3 quarks) for the hadrons, and only observed leptons? The answer, modulo anomalies, is a very natural yes. Our removal of anomalies, for reasons mentioned below, involves fermionic doubling; this will be discussed after presentation of the model, along with inclusion of other lepton models.

* Supported in part by the U.S. Atomic Energy Commission.

† Participating guest, Lawrence Berkeley Laboratory.

§ Address after September 1972: Department of Physics, University of Pennsylvania, Philadelphia, Pa. 19104.

Our approach to the synthesis is orderly: Starting with the $U(3) \otimes U(3)$ gauge model of hadrons, and Weinberg's $SU(2) \otimes U(1)$ model, we study embedding the latter in progressively larger "primed"² groups. Details of this study (and intermediate models), will appear in a larger paper. Here we sketch the emerging picture: The structure of the hadron theory requires the "primed" group at least as large as $SU(3) \otimes SU(3)$,² and Weinberg's model can be embedded therein. Unfortunately, models of this type have trouble with strangeness-changing processes. When we embed the leptons in $U(4) \otimes U(4)$ however, everything falls together beautifully, and it is this model we now present.

Groups and Representations. The hadronic group, entirely local, is $U(3)_L \otimes U(3)_R$. We represent its generators F_{OL}, F_{OR} by $\frac{1}{2} \lambda^\alpha$ ($\alpha = 0, \dots, 8$, left or right), being the usual 3×3 $SU(3)$ matrices. The local leptonic group is $SU(2)_L \otimes U(1)$, embedded in a "primed" $U(4)_L \otimes U(4)_R$. We call the latter's generators $F'_{\beta L}, F'_{\beta R}$ ($\beta = 0, \dots, 15$), but only four are realized locally. These are \tilde{F}'_{kL} ($k = 1, 2, 3$) and $\tilde{Y} = \tilde{F}'_{3R} + \frac{1}{3} (\tilde{F}'_{OR} + \tilde{F}'_{OL})$, with representations

$$\tilde{t}_\beta = R \begin{bmatrix} \frac{1}{2} \tau_\beta & 0 \\ 0 & \frac{1}{2} \hat{\tau}_\beta \end{bmatrix} R^{-1}, \quad R = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \theta & \sin \theta & 0 \\ 0 & -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad (1)$$

where τ_β ($\beta = 0, 1, 2, 3$, left or right) are the usual Pauli matrices, $\hat{\tau}_\beta \equiv \tau_2 \tau_\beta \tau_2$, and the twiddle operation R is the Cabibbo rotation. The charge operator is

$$Q = (F_{3L} + F_{3R}) + \frac{1}{\sqrt{3}} (F_{8L} + F_{8R}) + (F'_{3L} + F'_{3R}) + \frac{1}{3} (F'_{OL} + F'_{OR}). \quad (2)$$

Neutral operators do not rotate under R : $\tilde{t}_3 = t_3$, $\tilde{Y} = Y$, etc.

Local transformations. We represent the general local operator transformation in a unified super matrix notation

$$\mathcal{U} = \exp i(\alpha_L \cdot F_L + \alpha_R \cdot F_R + \beta \cdot F'_L + \gamma Y) \rightarrow S = \begin{pmatrix} S_L & 0 & 0 & 0 \\ 0 & S_R & 0 & 0 \\ 0 & 0 & \tilde{S}'_R & 0 \\ 0 & 0 & 0 & \tilde{S}'_L \end{pmatrix} \quad (3)$$

where, e.g., $\tilde{S}'_L = \exp i(\tilde{t} \cdot \beta + \frac{1}{3} t_0 \gamma)$, $\tilde{S}'_R = \exp i(t_3 + \frac{1}{3} t_0 \gamma)$, etc.

Fields. Let $V_\alpha^\mu, A_\alpha^\mu, W_k^\mu, B^\mu$ be (respectively) the strong vector and axial vector fields and the weak gauge bosons. Defining

$$V_{L,R}^\mu \equiv \sum_0^8 (V_\alpha^\mu \mp A_\alpha^\mu) \frac{\lambda_\alpha}{2}, \quad \tilde{W} \equiv \sum_1^3 \tilde{t}_k W_k, \quad \text{we specify the transformation}$$

properties of all gauge bosons at once in terms of a diagonal supermatrix V^μ (like S), with entries³ $V^\mu: [fV_L^\mu, fV_R^\mu, g'(t_3 + \frac{1}{3} t_0)B^\mu, g\tilde{W}^\mu + \frac{1}{3} g' t_0 B^\mu]$. Then $\mathcal{U} V_\mu \mathcal{U}^{-1} = S(V_\mu - i\partial_\mu)S^{-1}$. Similarly, for Weinberg's leptons, we introduce ($v_L = v_e, v_R = v_\mu^c$, D=doublet, S=singlet)

$$\Psi_D = \begin{bmatrix} v_L & \mu_R^c & & 0 \\ e_L^- & -\nu_e & & 0 \\ \hline 0 & ? & & ? \end{bmatrix}, \quad \Psi_S = \begin{bmatrix} 0 & \mu_L^c & & 0 \\ e_R & 0 & & 0 \\ \hline 0 & ? & & ? \end{bmatrix}, \quad \ell = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & \tilde{\Psi}_S & \frac{1}{2} \tilde{\Psi}_D \\ 0 & 0 & \frac{1}{2} \tilde{\Psi}_D & 0 \end{bmatrix}$$

(?) will be replaced later by heavy leptons to eliminate anomalies and Ψ_D^c is the charge conjugate of Ψ_D ; then $\mathcal{U} \ell \mathcal{U}^{-1} = S \ell S^{-1}$.

There is a great deal of freedom in the quark assignment. We will take here the simplest case, three fractionally charged quarks: (T=transpose) $q^T = (q_L, q_R, 0, 0)$, $\mathcal{U} q \mathcal{U}^{-1} = S q$. The supermatrix notation is most symmetric for the scalar mesons,

$$M = \begin{bmatrix} 0 & \Sigma & 0 & M_L \\ \Sigma^+ & 0 & M_R & 0 \\ 0 & M_R^+ & 0 & \phi^+ \\ M_L^+ & 0 & \phi & 0 \end{bmatrix}, \quad \mathcal{U} M \mathcal{U}^{-1} = S M S^{-1}. \quad (5)$$

Here, $\Sigma = \sigma + i\pi$ is the usual $(3, \bar{3}) \oplus (\bar{3}, 3)$ multiplet of scalars and pseudoscalars; $M_{L,R}$ are three-by-four complex matrices, being the scalars of Ref. (2), now with an extra fourth column. These scalars are the only connections of the weak and electromagnetic interactions with the strong, and will give mass to strong vector mesons;² ϕ is the Weinberg scalar field $\phi = \phi_0 t_0 + i\phi \cdot t$, which we see (in this notation) is to the leptonic system what Σ is to the hadronic.

Covariant derivatives and Lagrangian. The covariant derivatives can be read from the covariant momentum operator

$$\mathcal{P}^\mu \equiv P^\mu + f(V_L^\mu \cdot F_L + V_R^\mu \cdot F_R) + gW^\mu \cdot \tilde{F}'_L + g'B^\mu Y \quad (6)$$

by commuting this operator with each field. We find, $\Delta_\mu q \equiv \partial_\mu q - iV_\mu q$, $\Delta_\mu \ell \equiv \partial_\mu \ell - i[V_\mu, \ell]$, $\Delta_\mu M \equiv \partial_\mu M - i[V_\mu, M]$ and the usual $F_{\mu\nu}$'s for each gauge meson. Our locally invariant Lagrangian is

$$\begin{aligned} \mathcal{L} = & -\frac{1}{4} \text{Tr}(F_L^{\mu\nu} F_{\mu\nu}^L + F_R^{\mu\nu} F_{\mu\nu}^R) - \frac{1}{4} F_{\mu\nu}^B F_B^{\mu\nu} - \frac{1}{4} \text{Tr}(F_{\mu\nu}^W F_W^{\mu\nu}) \\ & - i\bar{q} \Delta^\mu \gamma_\mu q - i \text{Tr}(\bar{\ell} \Delta^\mu \gamma_\mu \ell) - \frac{1}{4} \text{Tr}(\Delta^\mu M)^\dagger \Delta_\mu M + \alpha(\bar{q}_L \Sigma q_R + \text{h.c.}) \\ & + \text{Tr}(\tilde{\Psi}_D \tilde{\phi} \tilde{\Psi}_S \tilde{G} + \text{h.c.}) + V(M_L) + V(M_R) + V(\Sigma) + V(\phi) \\ & + \text{Tr}(G_2 \tilde{\phi}^+ M_L^+ M_L \tilde{\phi} + G_2 M_R^+ M_R \tilde{\phi}^+) + \text{Tr}(G_1 \tilde{\phi}^+ M_L^+ \Sigma M_R + \text{h.c.}) \end{aligned} \quad (7)$$

where the $V(\dots)$'s are the usual quartic, and quadratic terms,³ and the G "insertions",² which are 4×4 diagonal matrices with entries

$$G: \frac{2}{\lambda}(m_e, m_\mu, ?, ?), \quad G_1: \frac{\sqrt{2}}{2\lambda} \begin{pmatrix} \frac{f m^2}{\pi \pi} & \frac{f m^2}{\pi \pi} & \frac{2f m^2 - f m^2}{2 \pi \pi} \\ \kappa_1 & \kappa_1 & \kappa_2 \end{pmatrix},$$

$G_2: (a, a, b, c)$, do not spoil the unified gauge invariance. The interpretation of the parameters in these insertions will be clarified in the following paragraph.

Spontaneous breakdown and symmetries. A detailed study of the complicated scalar system will be presented in a larger paper. Here we sketch

the general ideas. First, we use 21 degrees of gauge freedom (all but Q) to eliminate the 3×3 submatrices of $M_L - M_L^+$ and $M_R - M_R^+$, and all the components of ϕ except ϕ_0 . Next, in order to give masses to all the gauge fields except the photon, we assign vacuum expectation values $\langle \phi \rangle \equiv \lambda t_0$, $\langle M_L \rangle = \langle M_R \rangle \equiv \kappa$. These then generate a linear term in Σ (last term in \mathcal{L}). Thus Σ itself acquires a vacuum expectation value $\langle \Sigma \rangle = v$, which is the usual $(3, \bar{3}) \oplus (\bar{3}, 3)$ hadronic symmetry breaking in the spirit of Gell-Mann, Oakes, and Renner.⁴ It further turns out that the system allows the following arbitrary vacuum expectation values:

$$\kappa = \begin{bmatrix} \kappa_1 & 0 & 0 & 0 \\ 0 & \kappa_1 & 0 & 0 \\ 0 & 0 & \kappa_2 & 0 \end{bmatrix} \quad v = \frac{1}{\sqrt{2}} \begin{bmatrix} f_\pi & 0 & 0 \\ 0 & f_\pi & 0 \\ 0 & 0 & 2f_K - f_\pi \end{bmatrix} \quad (8)$$

and no Goldstone bosons. Except for d , the interpretation of the parameters in G_1 and v is standard,⁵ while G_2 , d , and $V(\dots)$ can be adjusted to give arbitrarily large masses to ϕ_0 and the remaining scalars in M_L and M_R ; hence with Ref. (2), we continue to regard these as unobservable entities. Actually, the model is perhaps more satisfactory with $\kappa_1 = \kappa_2$, leaving $\omega - \phi$ splitting until higher order in strong interactions. For this case we preserve the Weinberg sum rules⁶ and so we specialize to $\kappa_1 = \kappa_2$ below.

Photon system and diagonalization. Our spontaneous breakdown is such that the only unbroken gauge symmetry is generated by Q. Rewriting the covariant momentum (6), we isolate the (massless, universal) photon as the coefficient of Q:

$$A^\mu = \cos \eta (\sin \phi W_3^\mu + \cos \phi B^\mu) + \sin \eta \left(\frac{\sqrt{3}}{2} V_3^\mu + \frac{1}{2} V_8^\mu \right) \quad (9)$$

$$e = g \sin \phi \cos \eta; \quad \tan \phi = \frac{g'}{g}, \quad \tan \eta = \frac{2g \sin \phi}{\sqrt{3} f}.$$

With $f^2/4\pi \sim 2$ and g, g' small, we obtain approximately Weinberg's $e \sim gg'/(g^2 + g'^2)^{1/2}$. This diagonalization induces electromagnetic mixing of bare ρ_0, ϕ , and ω , such that the physical particles have order e^2/f couplings directly to the leptonic electromagnetic currents. This simulates vector-dominated electromagnetic form factors in lowest order, and gives a hadronic correction to the muonic $\frac{g-2}{2}$ which agrees with previous estimates.⁷ To keep the usual universality of weak interactions, we do not diagonalize the W^\pm -strong vector meson mixings involved in the term

$$\mathcal{L}' = gf \text{Tr} \{ V_L (M_L + \kappa) \tilde{W} (M_L^+ + \kappa) \}. \quad (10)$$

Thus, charged currents at low energy proceed via vector dominance in lowest order.

$\Delta S = 1$ neutral currents. Because our Cabibbo rotation rotates only W^\pm , we find no neutral $\Delta S = 1$ currents. In this model then, although we need four "things" to eliminate such currents, they are the columns of the unobservable $M_{L,R}$, and not extra quarks.

Fermions and anomalies. As thus far presented, the model has anomalies.⁸ Further, there appears to be no way, in the presence of both strong and weak vector mesons, to cancel hadronic against leptonic anomalies. Thus, we mention a flexible doubling scheme that for hadrons is in the spirit of dual models. We introduce $q', \psi_{S,D}^9$ that couple to gauge bosons just as $q, \psi_{S,D}$, but with the opposite sign of γ_5 . In the leptonic system anomalies are cancelled without complication. To avoid

suppressing $\pi_0 \rightarrow 2\gamma$, however, we also need a new Σ' , which transforms like Σ , but by choice, couples only to q' . It is then easy to arrange, with other terms in \mathcal{L} like $\text{Tr}[\bar{\psi}_L^+ M_L^+ \Sigma' M_R G_L^+]$ and $(\alpha' \bar{q}'_L \Sigma' q'_R + \text{h.c.})$, that the masses of q' , Σ' are high with negligible effect on V, A masses. Then, $\pi^0 \rightarrow 2\gamma$ proceeds only through q . To get an extra factor of three¹⁰ in amplitude, there are a number of choices--the simplest being the introduction of two more "pairs" of cancelling quarks (like q, q') with large mass.¹¹

Other lepton models. Among the other lepton models in the literature, one stands out as fitting our hadrons as well as Weinberg's. This is the "second" model of Prentki and Zumino¹ which may confront neutral current measurements more successfully than Weinberg. The Prentki-Zumino leptons fit into our $\psi_{D,S}$ using the lower right-hand corners as well. All other details are essentially the same as above. Not all leptonic theories fit our hadron theory however. For example, the model of Georgi and Glashow,¹ if it fits at all, appears unnatural. In the first place we need a $U(5) \otimes U(5)$ hadronic group (five quarks before anomaly cancellation), and worse, their scalar field transforms such that, without further scalars, we cannot find a $(3, \bar{3}) \oplus (\bar{3}, 3)$ symmetry breaking term like $\text{Tr}[\bar{\psi}_L^+ M_L^+ \Sigma M_R]$.

Conclusions. To the best of our knowledge, our unified model is consistent with known low-energy data, including vector meson dominance at low energies for electromagnetic and weak form factors, and accepted theoretical ideas about broken hadron symmetries etc.--in the presence of explicit hadron dynamics.

The question of deep inelastic scaling for our model (in perturbation theory) remains to be investigated. Although it turns out that

the current algebra generally resembles algebra of fields, we do not expect worse scaling properties than other renormalizable (longitudinally-damped) theories.¹² Possibly relevant to this question is the interesting fact that this unified model can be taken formally scale invariant before spontaneous breakdown: By omitting quadratic mass terms and adding terms of the form $\text{Tr}(M_L M_L^+ \Sigma \Sigma^+)$ etc., all masses for the scalars are generated spontaneously. In this case, scale invariance and internal symmetries are broken together.

We find it very encouraging that a unified renormalizable gauge theory of strong, weak, and electromagnetic interactions exists in which all three forces derive from a single, stringent principle: gauge invariance.

We acknowledge helpful conversations with K. Bardakci, M. Suzuki, and D. Levy.

FOOTNOTES AND REFERENCES

1. S. Weinberg, Phys. Rev. Letters 19, 1264 (1967) and 27, 1688 (1971).
M. Georgi and S. L. Glashow, Phys. Rev. Letters 28, 1494 (1972).
B. W. Lee, to be published. J. Prentki and B. Zumino, to be published.
2. K. Bardakci and M. B. Halpern, to be published. The "primed" or "leptonic" groups here are the "global" groups of this reference.
3. For different ninth meson couplings, let $f_{V_{L,R}} \rightarrow f \sum_1^8 (V \mp A) \cdot \frac{\lambda}{2} + f'(V_0 \mp A_0) \cdot \frac{\lambda_0}{2}$. The ninth axial vector meson seems needed to avoid a Goldstone boson in $(M - M^+)_{L,R}$, thus we cannot use the term $\det \Sigma + \det \Sigma^+$. As an effective Lagrangian then our model has a $\pi - \eta$ degeneracy.
4. M. Gell-Mann, R. J. Oakes, and B. Renner, Phys. Rev. 175, 2195 (1968). S. L. Glashow and S. Weinberg, Phys. Rev. Letters 20, 224 (1968). We emphasize that our ability to construct this important last term in \mathcal{L} depends on the ϕ, Σ symmetry observed above. A similar mechanism was suggested by Weinberg (Ref. 1).
5. S. Gasiorowicz and D. A. Geffen, Rev. Mod. Phys. 41, 531 (1969). Actually, there is freedom to induce $\langle \sigma_3 \rangle$ as well, if desired, for a lowest order model of $\eta \rightarrow 3\pi$.
6. S. Weinberg, Phys. Rev. Letters 18, 507 (1967) and T. D. Lee, S. Weinberg, and B. Zumino, Phys. Rev. Letters 18, 1209 (1967).
7. M. Gourdin and E. de Rafael, Nucl. Phys. B10, 667 (1969).
8. D. Gross and R. Jackiw; also C. Bouchiat, J. Iliopoulos, Ph. Meyer, to be published.

9. ψ'_S must also include (heavy) neutrino entries on the diagonal, which do not couple to gauge bosons. With minor rearrangement, the new heavy leptons fit in Eq. (4) in place of (?).
10. S. Adler, Lectures on Elementary Particles and Quantum Field Theory, edited by S. Deser et al. (MIT Press, 1970).
11. An alternate mechanism uses two "pairs" of integrally charged quarks with Y providing a "charm". Only one triplet couples to Σ by choice.
12. S. Adler and Wu-Ki Tung, Phys. Rev. Letters 22, 978 (1969).

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720