



ERNEST ORLANDO LAWRENCE BERKELEY NATIONAL LABORATORY

Glueball Mass Spectrum from Supergravity

Csaba Csáki, Hirosi Ooguri, Yaron Oz,
and John Terning

Physics Division

June 1998

Submitted to
Nuclear Physics B



Lawrence Berkeley National Laboratory

REFERENCE COPY
Does Not
Circulate

Bldg. 50 Library - Ref.

Copy 1

LBNL-41875

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Glueball Mass Spectrum from Supergravity

Csaba Csáki,^{1,2} Hirosi Ooguri,^{1,2,3} Yaron Oz,^{1,2,3} and John Terning^{1,2}

¹Department of Physics
University of California, Berkeley
Berkeley, California 94720

²Theoretical Physics Group
Physics Division
Ernest Orlando Lawrence Berkeley National Laboratory
University of California
Berkeley, California 94720

³Institute for Theoretical Physics
University of California
Santa Barbara, California 93106

June 1998

Glueball Mass Spectrum From Supergravity

Csaba Csáki^{1,2*}, Hiroshi Ooguri^{1,2,3}, Yaron Oz^{1,2,3} and John Terning^{1,2}

¹ Department of Physics, University of California at Berkeley,
Berkeley, CA 94720

² Theoretical Physics Group, Mail Stop 50A-5101,
Lawrence Berkeley National Laboratory,
Berkeley, CA 94720

³ Institute for Theoretical Physics, University of California,
Santa Barbara, CA 93106

csaki@thwk5.lbl.gov, ooguri@thsrv.lbl.gov, yaronoz@thsrv.lbl.gov,
terning@alvin.lbl.gov

Abstract

We calculate the spectrum of glueball masses in non-supersymmetric Yang-Mills theory in three and four dimensions, based on a conjectured duality between supergravity and large N gauge theories. The glueball masses are obtained by solving supergravity wave equations in a black hole geometry. We find that the mass ratios are in good numerical agreement with the available lattice data. We also compute the leading $(g_{YM}^2 N)^{-1}$ corrections to the glueball masses, by taking into account stringy corrections to the supergravity action and to the black hole metric. We find that the corrections to the masses are negative and of order $(g_{YM}^2 N)^{-3/2}$. Thus for a fixed ultraviolet cutoff the masses decrease as we decrease the 't Hooft coupling, in accordance with our expectation about the continuum limit of the gauge theories.

*Research Fellow, Miller Institute for Basic Research in Science.

1 Introduction

Recently Maldacena formulated a conjecture [1] stating that the large N limit of the maximally supersymmetric conformal theories in 3, 4 and 6 dimensions are dual to superstring/ M theory on $\text{AdS}_4 \times \text{S}^7$, $\text{AdS}_5 \times \text{S}^5$ and $\text{AdS}_7 \times \text{S}^4$ respectively, where AdS_d is a d -dimensional anti-de Sitter space. More recently Witten proposed [2] that one can extend this duality to non-supersymmetric theories such as pure QCD. In this case the AdS space is replaced by the Schwarzschild geometry describing a black hole in the AdS space. When the curvature of the spacetime is small compared to the string scale and the Planck scale, superstring/ M theory is well-approximated by supergravity. It was found that the supergravity description gives results that are in qualitative agreement with expectations for QCD at strong coupling. This includes the area law behavior of Wilson loops, the relation between confinement and monopole condensation, the existence of a mass gap for glueball states, the behavior of Wilson loops for higher representations, and the construction of heavy quark baryonic states [2–7].

In this paper, we use the supergravity description of large N gauge theories to compute the scalar glueball mass spectrum explicitly for pure QCD_3 and QCD_4 . The glueball masses in QCD can be obtained by computing correlation functions of gauge invariant local operators or the Wilson loops, and looking for particle poles. According to the refinement of Maldacena’s conjecture given in [8, 9], correlation functions of a certain class of local operators (chiral primary operators and their superconformal descendants) are related at large N and large $g_{YM}^2 N$ to tree level amplitudes of supergravity. The correspondence between the chiral operators and the supergravity states has been worked out in [8–14]. For example, the operator $\text{tr} F^2$ in four dimensions corresponds to the dilaton field of supergravity in ten dimensions. Therefore the scalar glueball* $J^{PC} = 0^{++}$ in QCD which couples to $\text{tr} F^2$ is related to the dilaton propagating in the black hole geometry. In particular, its mass is computable by solving the dilaton wave equation [2]. In [7], it was shown that the correlation function of Wilson loops is also expressed in terms of supergraviton exchange if the distance between the loops becomes larger than their sizes, leading again to the supergravity wave equation.

In this paper we will solve the wave equations numerically to obtain the glueball masses. Since this description preserves all the symmetries of QCD, we can identify the spin and the other quantum numbers of the glueballs. The mass ratios turn out to be in excellent agreement with the available lattice data in the continuum limit. This

*In the following we will use the notation J^{PC} for the glueballs, where J is the glueball spin, and P , C refer to the parity and charge conjugation quantum numbers respectively.

is surprising since a priori the supergravity computations are to be compared with the strong ultraviolet coupling limit of the gauge theory $g_{YM}^2 N \gg 1$.

As we will see, the supergravity computation at $g_{YM}^2 N \gg 1$ gives the glueball masses in units of the fixed ultraviolet cutoff Λ_{UV} . For finite 't Hooft coupling $\lambda = g_{YM}^2 N$, the glueball mass M would be a function of the form,

$$M^2 = f(\lambda)\Lambda_{UV}^2. \quad (1.1)$$

To take the continuum limit $\Lambda_{UV} \rightarrow \infty$, we have to simultaneously take $\lambda \rightarrow 0$ so that the right-hand side of this equation becomes of the order of the QCD mass scale Λ_{QCD} . This in particular requires that $f(\lambda)$ decreases as we decrease the 't Hooft coupling λ .

We compute the leading λ^{-1} corrections to the supergravity computation and show that this is indeed the case. On the superstring side, the λ^{-1} corrections are due to the finite string tension. The leading order string correction to the low-energy supergravity action was computed in [15, 16]. This modifies both the background black hole metric and the supergravity wave equation in that background. Recently the stringy correction to the black hole metric was obtained in [17] by solving the modified supergravity equation. We use both this metric and the string corrected wave equation to compute the leading λ^{-1} corrections to the 0^{++} glueball masses in QCD_3 . We find:

(1) The corrections to the masses are negative and of order $\lambda^{-3/2}$:

$$f(\lambda) = c_0 + c_1\lambda^{-3/2} + \dots, \quad c_1 < 0, \quad (1.2)$$

for the ground state and the first 5 excited levels of the 0^{++} glueball. Thus, for a fixed ultraviolet cutoff, the masses decrease as we decrease the 't Hooft coupling, in accordance with the expectation about the continuum limit of QCD.

(2) The corrections to the ratios of the glueball masses are relatively small compared to the correction to each glueball mass, suggesting that the corrections are somewhat universal for all the glueball masses. This may indicate that the good agreement between the supergravity computation and the lattice gauge theory results is not a coincidence but is due to small λ^{-1} corrections to the mass ratios.

This paper is organized as follows.

In section 2, we solve the supergravity wave equations in the AdS_5 black hole geometry to obtain glueball masses in QCD_3 and compare the results with lattice computations.

In section 3, we solve the supergravity wave equations in the AdS_7 black hole geometry to obtain glueball masses in QCD_4 and compare the results with lattice computations.

In section 4, we use the string theory corrections to the low-energy supergravity action and to the AdS₅ black hole geometry to estimate corrections to the glueball masses in QCD₃.

We close the paper with a summary and discussions.

2 Glueballs in Three Dimensions

The $\mathcal{N} = 4$ superconformal $SU(N)$ gauge theory in four dimensions is realized as a low energy effective theory of N coinciding parallel D3 branes. One can construct a three-dimensional non-supersymmetric theory [2] by compactifying this theory on $\mathbf{R}^3 \times \mathbf{S}^1$ with anti-periodic boundary conditions on the fermions around the compactifying circle \mathbf{S}^1 . Supersymmetry is broken explicitly by the boundary conditions. As the radius R of the circle becomes small, the fermions decouple from the system since there are no zero frequency Matsubara modes. The scalar fields in the 4D theory will acquire masses at one-loop, since supersymmetry is broken, and these masses become infinite as $R \rightarrow 0$. Therefore in the infrared we are left with only the gauge field degrees of freedom and the theory should be effectively the same as pure QCD₃.

According to Maldacena [1], the $\mathcal{N} = 4$ theory in Euclidean \mathbf{R}^4 is dual to type IIB superstring theory on AdS₅ \times \mathbf{S}^5 with the metric

$$\frac{ds^2}{l_s^2 \sqrt{4\pi g_s N}} = \rho^{-2} d\rho^2 + \rho^2 \sum_{i=1}^4 dx_i^2 + d\Omega_5^2 \quad (2.1)$$

where l_s is the string length related to the superstring tension, g_s is the string coupling constant and $d\Omega_5$ is the line element on \mathbf{S}^5 . The $x^{1,2,3,4}$ directions in AdS₅ correspond to \mathbf{R}^4 where the gauge theory lives. The gauge coupling constant g_4 of the 4D theory is related to the string coupling constant g_s as $g_4^2 = g_s$. In the 't Hooft limit ($N \rightarrow \infty$ with $g_4^2 N = g_s N$ fixed), the string coupling constant vanishes $g_s \rightarrow 0$. Therefore we can study the 4D theory using the first quantized string theory in the AdS space (2.1). Moreover if $g_s N \gg 1$, the curvature of the AdS space is small and the string theory is approximated by classical supergravity.

Upon compactification on \mathbf{S}^1 and imposing the supersymmetry breaking boundary conditions, (2.1) is replaced by the Euclidean black hole geometry [2]

$$\frac{ds^2}{l_s^2 \sqrt{4\pi g_s N}} = \left(\rho^2 - \frac{b^4}{\rho^2} \right)^{-1} d\rho^2 + \left(\rho^2 - \frac{b^4}{\rho^2} \right) d\tau^2 + \rho^2 \sum_{i=1}^3 dx_i^2 + d\Omega_5^2 \quad (2.2)$$

where τ parameterizes the compactifying circle and the $x^{1,2,3}$ direction corresponding to

the \mathbf{R}^3 where QCD_3 lives. The horizon of this geometry is located at $\rho = b$ with

$$b = \frac{1}{2R}. \quad (2.3)$$

Once again, the supergravity approximation is applicable for $N \rightarrow \infty$ and $g_s N \gg 1$.

According to [8, 9, 12–14], there is a one-to-one correspondence between supergravity wave solutions on $\text{AdS}_5 \times \mathbf{S}^5$ and chiral primary fields (and their descendants) in the $\mathcal{N} = 4$ superconformal theory in four dimensions. The mass m of a p -form C on the AdS space is related to the dimension Δ of a $(4 - p)$ form operator in the $\mathcal{N} = 4$ theory by

$$m^2 = (\Delta - p)(\Delta + p - 4). \quad (2.4)$$

The supergravity fields on $\text{AdS}_5 \times \mathbf{S}^5$ can be classified by decomposing them into spherical harmonics (the Kaluza-Klein modes) on \mathbf{S}^5 . They fall into irreducible representations of the $SO(6)$ isometry group of \mathbf{S}^5 , which is also the R -symmetry group of the 4D superconformal theory. The spectrum of Kaluza-Klein harmonics of type IIB supergravity on $\text{AdS}_5 \times \mathbf{S}^5$ was derived in [18, 19]. Among them, there are four Kaluza-Klein modes that are $SO(6)$ singlets, coming from the s -wave components on \mathbf{S}^5 of bosonic fields. They are:

- (1) The graviton $g_{\mu\nu}$ polarized along the \mathbf{R}^4 in (2.2). It couples to the dimension 4 stress-energy tensor $T_{\mu\nu}$ of the $\mathcal{N} = 4$ theory.
- (2) The dilaton and the R-R scalar, which combine into a complex massless scalar field. Its real and imaginary parts couple to the dimension 4 scalar operators $\mathcal{O}_4 = \text{tr } F^2$ and $\tilde{\mathcal{O}}_4 = \text{tr } F \wedge F$ of the $\mathcal{N} = 4$ theory respectively.
- (3) The NS-NS and R-R two-forms, which combine into a complex-valued antisymmetric field $A_{\mu\nu}$, polarized along the \mathbf{R}^4 . Its $(\text{AdS mass})^2 = 16$ and using (2.4) we see that it couples to a dimension 6 two-form operator of the $\mathcal{N} = 4$ theory. This operator has been identified as $\mathcal{O}_6 = d^{abc} F_{\mu\alpha}^a F^{b\alpha\beta} F_{\beta\nu}^c$ [22, 23].
- (4) The s -wave component of the metric g_α^α and the R-R 4-form $A_{\alpha\beta\gamma\delta}$ polarized along \mathbf{S}^5 . They combine into a massive scalar with $(\text{AdS mass})^2 = 32$ and couple to a dimension 8 scalar operator constructed from the gauge field strength $F_{\mu\nu}$ of the $\mathcal{N} = 4$ theory [24, 23]. Only these $SO(6)$ singlet fields are related to glueballs of QCD_3 since $SO(6)$ non-singlets are supposed to decouple in the limit $R \rightarrow 0$.

Let us discuss now how to identify the quantum numbers of the glueballs. The spin and the parity of a glueball in three dimensions can be easily found from the transformation properties of the corresponding supergravity field. The charge conjugation, C , for

gluons is defined by $A_\mu^a T_{ij}^a \rightarrow -A_\mu^a T_{ji}^a$ where the T^a 's are the hermitian generators of the gauge group [25]. In the string theory, charge conjugation corresponds to the worldsheet parity transformation changing the orientation of the open string attached to D-branes. Therefore, for example, the NS-NS two-form in supergravity is odd under the charge conjugation. This is consistent with the fact that it couples to \mathcal{O}_6 , which indeed has $C = -1$.

From the point of view of QCD₃, the radius R of the compactifying circle provides the ultraviolet cutoff scale. To obtain large N QCD₃ in the continuum, one has to take $g_4^2 N \rightarrow 0$ as $R \rightarrow 0$ so that $g_3^2 N = g_4^2 N/R$ remains at the intrinsic energy scale of QCD₃. Here g_3 is the dimensionful gauge coupling of QCD₃. This is the opposite of the limit that is required for the supergravity description to be valid. As we mentioned, the supergravity description is applicable for $g_4^2 N \gg 1$. Therefore, with the currently available techniques, the Maldacena-Witten conjecture can only be used to study large N QCD with a fixed ultraviolet cutoff R^{-1} in the strong ultraviolet coupling regime. The results we find are, however, surprisingly close to those of the lattice computation, leading us to suspect that $(g_4^2 N)^{-1}$ corrections to the mass ratios are small. In section 4, we will estimate the leading $(g_4^2 N)^{-1}$ correction to our computation.

Consider first the 0^{++} glueball masses. These can be derived from the 2-point function of the operator $\text{tr } F_{\mu\nu} F^{\mu\nu}$. In the supergravity description we have to solve the classical equation of motion of the massless dilaton,

$$\partial_\mu [\sqrt{g} \partial_\nu \Phi g^{\mu\nu}] = 0, \quad (2.5)$$

on the AdS₅ black hole background (2.2). In order to find the lowest mass modes we assume following [2] that Φ is independent of τ and has the form $\Phi = f(\rho)e^{ikx}$. Using the metric of (2.2) one obtains the following differential equation for f :

$$\rho^{-1} \frac{d}{d\rho} \left((\rho^4 - b^4) \rho \frac{df}{d\rho} \right) - k^2 f = 0. \quad (2.6)$$

Since the glueball mass M^2 is equal to $-k^2$, the task is to solve this equation as an eigenvalue problem for k^2 . In the following we set $b = 1$, so the masses are computed in units of b . If one changes variables to $x = \rho^2$, the equation takes the form,

$$\frac{d^2 f}{dx^2} + \left(\frac{1}{x} + \frac{1}{x-1} + \frac{1}{x+1} \right) \frac{df}{dx} - \frac{k^2}{4x(x^2-1)} f = 0, \quad (2.7)$$

namely it is an ordinary differential equation with four regular singularities at $x = 0, \pm 1$ and ∞ .

Unlike the equation with three regular singularities (known as the hypergeometric equation), analytic solutions are not known for this type of equation. Fortunately there is an analytical method to compute its eigenvalues k^2 . It is the exact WKB analysis recently developed by mathematicians at RIMS, Kyoto University [20]. To use their approach, we note that the differential equation (2.7) can be written as the Schrödinger-type equation

$$\left(-\frac{d^2}{dx^2} + Q(x)\right)g(x) = 0, \quad (2.8)$$

where $g(x) = \sqrt{x(x^2 - 1)}f(x)$ and

$$Q(x) = \frac{3x^4 - 6x^2 - 1}{4x^2(x^2 - 1)^2} + \frac{k^2}{4x(x^2 - 1)}. \quad (2.9)$$

To apply the WKB analysis, one can perturb the equation as

$$\left(-\frac{d^2}{dx^2} + Q(x) + (\eta^2 - 1)R(x)\right)g(x) = 0, \quad (2.10)$$

by introducing a large parameter η . With a suitable choice of $R(x)$, the secular equation, which determines the values of k^2 so that the equation admits a solution regular at both $x = 1$ and ∞ , becomes explicitly solvable as a asymptotic power series expansion in η^{-1} . Assuming the expansion is Borel summable at $\eta = 1$, the eigenvalues are approximated by the following expression [21]

$$k^2 = -6n(n + 1), \quad (n = 1, 2, 3, \dots). \quad (2.11)$$

We should note that the differential equation in question is degenerate from the point of view of the exact WKB analysis and a mathematical proof of the Borel summability in this case has not been given. It is possible that the formula (2.11) receives small corrections.

Since the analytical expression (2.11) for k^2 is still preliminary and we would like to find masses for the other glueball states, we also solved the differential equation (2.6) numerically. For large ρ , the black hole metric (2.2) asymptotically approaches the AdS metric, and the behavior of the solution for a p -form for large ρ takes the form ρ^λ , where λ is determined from the mass m of the supergravity field:

$$m^2 = \lambda(\lambda + 4 - 2p). \quad (2.12)$$

Indeed both (2.6) and (2.12) give the asymptotic forms $f \sim 1, \rho^{-4}$, and only the later is a normalizable solution [2]. Changing variables to $f = \psi/\rho^4$ we have:

$$(\rho^2 - \rho^6) \psi'' + (3\rho^5 - 7\rho) \psi' + (16 + k^2\rho^2) \psi = 0. \quad (2.13)$$

For large ρ this equation can be solved by series solution with negative even powers:

$$\psi = \sum_{n=0}^{\infty} a_{2n} \rho^{-2n} . \quad (2.14)$$

Since the normalization is arbitrary we can set $a_0 = 1$. The first few coefficients are given by:

$$\begin{aligned} a_2 &= \frac{k^2}{12} \\ a_4 &= \frac{1}{2} + \frac{k^4}{384} \\ a_6 &= \frac{7k^2}{120} + \frac{k^6}{23040} . \end{aligned} \quad (2.15)$$

For $n \geq 5$ the coefficients are given by the recursive relation:

$$(n^2 + 4n)a_n = k^2 a_{n-2} + n^2 a_{n-4} . \quad (2.16)$$

Since the black hole geometry is regular at the horizon $\rho = 1$, k^2 has to be adjusted so that f is also regular at $\rho = 1$ [2]*. This can be done numerically in a simple fashion using a “shooting” technique as follows. For a given value of k^2 the equation is numerically integrated from some sufficiently large value of ρ ($\rho \gg k^2$) by matching $f(\rho)$ with the asymptotic solution set by (2.14) and (2.15). The glueball mass M is related to the eigenvalues of k^2 by $M^2 = -k^2$ in units of b^2 . The results of the numerical work are listed in Table 1. They agree with the formula (2.11). The 4% discrepancy of the two results are either due to some systematic error in the numerical analysis or due to corrections to the analytical formula (2.11).

state	numerical method	exact WKB method	ratio
0^{++}	11.59	12	1.03
0^{+++}	34.53	36	1.04
0^{++++}	68.98	72	1.04
0^{+****}	114.9	120	1.04
0^{+*****}	172.3	180	1.04
0^{+*****}	241.2	252	1.04

Table 1: $(\text{Mass})^2$ of 0^{++} glueball in QCD_3 obtained by solving the supergravity wave equation in the black hole geometry (in units of b^2) using the two different methods.

*We thank A. Jevicki and J. P. Nunes for communication on the boundary condition.

Since both methods give the same results within a 4% error, we are ready to compare them with the lattice gauge theory computations [27]. Since the lattice results are in units of string tension, we normalize the supergravity results so that the lightest 0^{++} state agrees with the lattice result. The results are listed in Table 2. One should also expect a systematic error in addition to the statistical error denoted in Table 2 for the lattice computations.

state	lattice, $N = 3$	lattice, $N \rightarrow \infty$	supergravity
0^{++}	4.329 ± 0.041	4.065 ± 0.055	4.07 (input)
0^{+++}	6.52 ± 0.09	6.18 ± 0.13	7.02
0^{++++}	8.23 ± 0.17	7.99 ± 0.22	9.92
0^{+****}	-	-	12.80
0^{+*****}	-	-	15.67
0^{+*****}	-	-	18.54

Table 2: 0^{++} glueball masses in QCD_3 coupled to $\text{tr } F_{\mu\nu}F^{\mu\nu}$. The lattice results are in units of the square root of the string tension. The denoted error in the lattice results is only the statistical one.

Next we consider the two-form of the supergravity theory. As noted previously, it couples to the operator \mathcal{O}_6 . This operator contains 1^{+-} and 1^{--} components, which correspond to the fields $A_{\tau i}$ and A_{ij} , where $i, j = 1, 2, 3$ correspond to the three coordinates x_i of \mathbf{R}^3 . The remaining components $A_{\rho\tau}$ and $A_{\rho i}$ can be set to zero by an appropriate gauge transformation. In the QCD_3 limit $R \rightarrow 0$, the 1^{--} component A_{ij} is reduced to a 0^{--} operator in 3D, and thus has a non-zero overlap with the 0^{--} glueball[†]. On the other hand, the 1^{+-} components $A_{\tau i}$ couple to an operator which is supposed to decouple in the $R \rightarrow 0$ limit. Therefore they do not correspond to glueball states in QCD_3 .

The s -wave component of the two-form field satisfies the equation [18, 26]

$$\frac{3}{\sqrt{g}}\partial_\mu \left[\sqrt{g} \partial_{[\mu'} A_{\mu'_1 \mu'_2]} g^{\mu'\mu} g^{\mu'_1 \mu'_1} g^{\mu'_2 \mu'_2} \right] - 16g^{\mu'_1 \mu'_1} g^{\mu'_2 \mu'_2} A_{\mu'_1 \mu'_2} = 0, \quad (2.17)$$

where $[]$ denotes antisymmetrization with strength one. As before we look for solutions which are independent of τ and are of the form $A_{ij} = h_{ij}(\rho)e^{ikx}$. The $\rho\tau$ and the ρi components of this equation simply result in a constraint which sets the transverse component of A_{ij} to zero. For the remaining pseudoscalar component from A_{ij} the equation reduces

[†]The parity $P = -1$ is due to the fact that the 2-form is dual to a pseudoscalar. The charge conjugation $C = -1$ is inferred from the string worldsheet parity.

to

$$\rho(\rho^4 - 1)h'' + (3 + \rho^4)h' - (k^2\rho + 16\rho^3)h = 0, \quad (2.18)$$

in units where $b = 1$. At large ρ the solution has the form

$$h = \rho^{-4} \sum_{n=0}^{\infty} a_{2n} \rho^{-2n}. \quad (2.19)$$

Since the normalization is arbitrary we can again set $a_0 = 1$. The first few coefficients are given by:

$$\begin{aligned} a_2 &= \frac{k^2}{20} \\ a_4 &= \frac{640 + k^4}{960} \\ a_6 &= \frac{3520k^2 + k^6}{80640}. \end{aligned} \quad (2.20)$$

We have solved the differential equation and obtained the eigenvalues k^2 by the same numerical method described above. The results are shown in Table 3. The supergravity results are displayed in the same normalization as the one used in Table 2.

state	lattice, $N = 3$	lattice, $N \rightarrow \infty$	supergravity
0^{--}	6.48 ± 0.09	5.91 ± 0.25	6.10
0^{--*}	8.15 ± 0.16	7.63 ± 0.37	9.34
0^{--**}	9.81 ± 0.26	8.96 ± 0.65	12.37
0^{--***}	-	-	15.33
0^{--****}	-	-	18.26
$0^{--*****}$	-	-	21.16

Table 3: 0^{--} glueball masses in QCD_3 coupled to \mathcal{O}_6 . The lattice results are in units of square root of the string tension. The normalization of the supergravity results is the same as in Table 2.

Since the supergravity method and the lattice gauge theory compute the glueball masses in different units, one cannot compare the absolute values of the lowest glueball mass obtained using these methods. However it makes sense to compare the lowest glueball masses of different quantum numbers. Using Tables 2 and 3, we find that the supergravity results are in good agreement with the lattice gauge theory computation [27]:

$$\begin{aligned} \left(\frac{M_{0^{--}}}{M_{0^{++}}}\right)_{\text{supergravity}} &= 1.50 \\ \left(\frac{M_{0^{--}}}{M_{0^{++}}}\right)_{\text{lattice}} &= 1.45 \pm 0.08. \end{aligned} \quad (2.21)$$

There are still two more $SO(6)$ singlet supergravity fields. One of them is the s -wave component of the metric g_α^α and the R-R 4-form $A_{\alpha\beta\gamma\delta}$ polarized along \mathbf{S}^5 . From (2.4) we see that it should couple to a dimension 8 scalar operator \mathcal{O}_8 . In [23, 24], this operator is identified as a symmetrized form of $[F^4 - \frac{1}{4}(F^2)^2]$. By using the prescription of Tseytlin [28] to symmetrize the group indices, one finds that the operator is even under the charge conjugation. This is also seen from the fact that g_α^α is clearly even both spacetime and worldsheet parity transformations. Therefore g_α^α has the quantum numbers of the 0^{++} glueball. The classical equation of motion of g_α^α is that of a massive scalar with $(\text{AdS mass})^2 = 32$ (in units of b^2) [18] on the AdS_5 black hole background (2.2). The mass spectrum that we get is given in Table 4. In the $g_s N \rightarrow \infty$ limit the operators \mathcal{O}_8 and $\text{tr}F^2$ are not mixed since they couple to different states in the supergravity theory. However, we expect that for finite $g_s N$ these operators will mix, thus the full 0^{++} spectrum is expected to be given by the interleaving of Tables 2 and 4. For example the 0^{+++} presumably corresponds to the first state in Table 4.

<u>g_α^α and $A_{\alpha\beta\gamma\delta}$</u>
8.85
12.06
15.00
17.98

Table 4: 0^{++} glueball masses in QCD_3 coupled to \mathcal{O}_8 , the normalization is the same as in Table 2.

The remaining $SO(6)$ singlet is the graviton $g_{\mu\nu}$. It couples to the energy-momentum tensor $T_{\mu\nu}$ and therefore corresponds to the 2^{++} glueball. It would be interesting to compute its mass and compare with the lattice result.

3 Glueballs in Four Dimensions

To construct QCD_4 , one starts with the superconformal theory in six dimensions realized on N parallel coinciding M5-branes. The compactification of this theory on a circle of radius R_1 gives a five-dimensional theory whose low-energy effective theory is the maximally supersymmetric $SU(N)$ gauge theory with gauge coupling constant $g_5^2 = R_1$. To obtain QCD_4 , one compactifies this theory further on another \mathbf{S}^1 of radius R_2 . The gauge coupling constant g_4 in 4D is given by $g_4^2 = g_5^2/R_2 = R_1/R_2$. To break supersymmetry, one imposes the anti-periodic boundary condition on the fermions around the second \mathbf{S}^1 .

According to Maldacena [1], the large N limit of the six-dimensional theory is M theory on $\text{AdS}_7 \times \mathbf{S}^4$. Upon compactification on $\mathbf{S}^1 \times \mathbf{S}^1$ and imposing the anti-periodic boundary conditions around the second \mathbf{S}^1 , we find M theory to be on the black hole geometry [2]. To take the large N limit while keeping $g_4^2 N$ finite, we have to take $R_1 \ll R_2$. In this limit, M theory reduces to type IIA string theory and the M5 brane wrapping on \mathbf{S}^1 of radius R_1 becomes a D4 brane. The large N limit of QCD_4 then becomes string theory on the black hole geometry given by

$$\frac{ds^2}{l_s^2 g_5^2 N / 4\pi} = \frac{d\rho^2}{4\rho^{3/2} \left(1 - \frac{b^6}{\rho^3}\right)} + \rho^{3/2} \left(1 - \frac{b^6}{\rho^3}\right) d\tau^2 + \rho^{3/2} \sum_{i=1}^4 dx_i^2 + \rho^{1/2} d\Omega_4^2, \quad (3.1)$$

with a dilaton $e^\phi \sim \rho^{3/4}$ [29]. The location of the horizon $\rho = b^2$ is related to the radius R_2 of the compactifying circle as

$$b = \frac{1}{3R_2} \quad (3.2)$$

As in the case of three dimensions, we will compute the spectrum of glueball masses by solving the classical equations of motions of Kaluza-Klein modes of the supergravity theory. We will consider only singlets of the $SO(5)$ isometry group of \mathbf{S}^4 , which corresponds to the R -symmetry group of the six-dimensional theory.

Consider first the 0^{++} glueball. The non-extremal D4 brane solution has a non constant dilaton background. As shown in [30] the dilaton is a linear combination of two scalars. One of them is massless and couples to the relevant glueball operator. The equation of motion for the scalar is given by (2.5) in the background of the metric (3.1). Again assuming that the solution is independent of τ and of the form $\Phi = f(\lambda)e^{ikx}$ (with $\lambda^2 = \rho$), one obtains the differential equation in the units where $b = 1$ as

$$(\lambda^7 - \lambda)f''(\lambda) + (10\lambda^6 - 4)f'(\lambda) - \lambda^3 k^2 f(\lambda) = 0. \quad (3.3)$$

The asymptotic solutions to this equation are $f \sim 1, \lambda^{-9}$, with the latter corresponding to normalizable solutions. In order to solve the equation and find the allowed values of k^2 we introduce the function $g(\lambda)$ as $f(\lambda) = \lambda^{-9}g(\lambda)$. This way $g(\lambda)$ has to be asymptotically constant for $\lambda \rightarrow \infty$, and one can again look for a solution in terms of a negative power series in λ . The differential equation for $g(\lambda)$ is

$$(\lambda^8 - \lambda^2)g'' + (14\lambda - 8\lambda^7)g' - (\lambda^4 k^2 + 54)g = 0. \quad (3.4)$$

The first few coefficients in the power series solution $g = \sum_{n=0}^{\infty} a_{2n} \lambda^{-2n}$ are given by (for $a_0 = 1$)

$$a_2 = \frac{k^2}{22}$$

$$\begin{aligned}
a_4 &= \frac{k^4}{1144} \\
a_6 &= \frac{61776 + k^6}{102960} .
\end{aligned} \tag{3.5}$$

The regularity of f at $\lambda = 1$, after numerically solving the equation (3.4) as described in the previous section, results in the allowed values of k^2 . The first six masses (normalized so that the lightest 0^{++} state agrees with the lattice calculation) together with the available lattice results [32, 33] are given in Table 5.

state	lattice, $N = 3$	supergravity
0^{++}	1.61 ± 0.15	1.61 (input)
0^{++*}	2.8	2.38
0^{++**}	-	3.11
0^{++***}	-	3.82
0^{++****}	-	4.52
$0^{++*****}$	-	5.21

Table 5: Masses of the first few 0^{++} glueballs in QCD_4 , in GeV, from supergravity compared to the available lattice results. Note that the authors of ref. [33] do not quote errors for the 0^{++*} since it is not yet clear whether it is a genuine excited state or merely a two glueball bound state.

In order to calculate the masses of the 0^{-+} glueball in four dimensions we will consider the 3-form $A_{\alpha\beta\gamma}$ of the eleven dimensional supergravity. In this case, it is more useful to use the eleven-dimensional metric

$$ds^2 = \frac{d\lambda^2}{\left(\frac{\lambda^2}{b^2} - \frac{b^4}{\lambda^4}\right)} + \left(\frac{\lambda^2}{b^2} - \frac{b^4}{\lambda^4}\right) d\tau^2 + \lambda^2 \sum_{i=1}^5 dx_i^2 + d\Omega_4^2, \tag{3.6}$$

which reduces to (3.1) upon compactifying x_5 on \mathbf{S}^1 and by going to the string frame [2] by multiplying the metric by λ , setting $\lambda^2 = \rho$, and rescaling the other coordinates. The s -wave component of the 3-form in the harmonic expansion on \mathbf{S}^4 is a singlet of the $SO(5)$ isometry group [31]. Its mass squared* is 36 in units of b^2 and using (2.4) we see that it couples to a dimension 9 operator of the six-dimensional theory. The 3-form obeys the following equation of motion:

$$\frac{4}{\sqrt{g}} \partial_\mu \left[\sqrt{g} \partial_{[\mu'} A_{\mu'_1 \mu'_2 \mu'_3]} g^{\mu' \mu} g^{\mu'_1 \mu_1} g^{\mu'_2 \mu_2} g^{\mu'_3 \mu_3} \right] - 36 g^{\mu'_1 \mu_1} g^{\mu'_2 \mu_2} g^{\mu'_3 \mu_3} A_{\mu'_1 \mu'_2 \mu'_3} = 0 . \tag{3.7}$$

*The value of the mass term is fixed by matching the supergravity computation [31] to (2.4) [34].

Choosing a gauge where $A_{\rho\tau i}$ and $A_{\rho ij}$ vanish, where $i, j = 1, \dots, 5$, and assuming that the remaining components are independent of the coordinate τ and the only dependence on x_i is through e^{ikx} , one finds that there are two independent modes after compactification to 4D:

(1) A three-index tensor A_{ijk} . This is dual to a massive scalar and can be identified with the 0^{-+} glueball of the 4D theory.

(2) A massive vector $A_{\tau ij}$. This couples to an operator which is supposed to decouple in the limit $R_2 \rightarrow 0$. Therefore it does not correspond to a glueball state in QCD_4 .

The scalar component of A_{ijk} satisfies the differential equation

$$(\lambda^7 - \lambda)f''(\lambda) + (\lambda^6 + 5)f'(\lambda) - \lambda^3(k^2 + 36\lambda^2)f(\lambda) = 0, \quad (3.8)$$

in the same units as in the equation (3.3) for the dilaton. The normalizable asymptotic solution behaves like $1/\lambda^6$, thus we introduce the function $g(\lambda)$ by $f(\lambda) = \lambda^{-6}g(\lambda)$. This satisfies

$$(\lambda^7 - \lambda)g''(\lambda) - (11\lambda^6 - 17)g'(\lambda) - (72 + k^2\lambda^4)g(\lambda) = 0. \quad (3.9)$$

The power series expansion $g(\lambda) = \sum_{n=0}^{\infty} a_{2n}\lambda^{-2n}$ with $a_0 = 1$ has the first few coefficients

$$\begin{aligned} a_2 &= \frac{k^2}{28} \\ a_4 &= \frac{k^4}{1792} \\ a_6 &= \frac{129024 + k^6}{193536} \end{aligned} \quad (3.10)$$

The single-valuedness of the solution requires $g' = 6g$ at $\lambda = 1$. With this boundary condition, the numerical solution of (3.9) yields the allowed values of k^2 , and the corresponding masses in the units defined above are displayed in Table 6.

state	lattice, $N = 3$	supergravity
0^{-+}	2.19 ± 0.32	1.83
0^{-+*}	-	2.67
0^{-+**}	-	3.42
0^{-+***}	-	4.14
0^{-+****}	-	4.85
$0^{-+*****}$	-	5.55

Table 6: Masses of 0^{-+} glueball in QCD_4 . The lattice result is in GeV.

Unlike the 3D case, there exists little lattice data on the masses of the excited glueball states. We can however compare the ratio of masses of the lowest glueball states 0^{-+} and 0^{++}

$$\begin{aligned} \left(\frac{M_{0^{-+}}}{M_{0^{++}}}\right)_{\text{supergravity}} &= 1.14 \\ \left(\frac{M_{0^{-+}}}{M_{0^{++}}}\right)_{\text{lattice}} &= 1.36 \pm 0.32, \end{aligned} \quad (3.11)$$

and the results are in agreement within the one σ error.

4 Leading String Theory Corrections

As we mentioned earlier, the supergravity computation is valid in the strong ultraviolet coupling limit $g_s N \gg 1$. In order to compare with the lattice computations in the continuum limit, we have to take $g_s N \rightarrow 0$ as we take the ultraviolet cutoff $R^{-1} \rightarrow \infty$ so that the scale set by the Yang-Mills coupling constant remains at the intrinsic energy scale of QCD. The fact that the glueball masses computed in the supergravity limit are in good agreement with the lattice results leads us to suspect that, for this particular computation, α' corrections are small. In this section, we test this idea.

For $g_s N \ll 1$, the curvature of the black hole geometry becomes larger than the string scale. Therefore stringy corrections (to be precise, the worldsheet sigma-model corrections) are expected to become important. The leading stringy corrections to the low-energy supergravity action were obtained in [15, 16]. Recently Gubser, Klebanov and Tseytlin [17] used the modified action to obtain the leading order string corrections to the black hole metric. We use their result to calculate the leading corrections to the glueball mass spectrum. We will perform this computation only for the 0^{++} glueballs in QCD_3 . We expect, however, that the conclusions will be similar for the other glueball states.

According to [17], the leading (in units of the curvature) $\alpha' = (4\pi g_s N)^{-1/2}$ correction to the AdS_5 black hole metric (2.2) is

$$\frac{ds^2}{l_s^2 \sqrt{4\pi g_4^2 N}} = (1 + \delta_1) \frac{d\rho^2}{\left(\rho^2 - \frac{b^4}{\rho^2}\right)} + (1 + \delta_2) \left(\rho^2 - \frac{b^4}{\rho^2}\right) d\tau^2 + \rho^2 \sum_{i=1}^3 dx_i^2, \quad (4.1)$$

where the correction terms $\delta_{1,2}$ are given by the formulae

$$\begin{aligned} \delta_2 &= -15\gamma \left(5 \frac{b^4}{\rho^4} + 5 \frac{b^8}{\rho^8} - 3 \frac{b^{12}}{\rho^{12}}\right) \\ \delta_1 &= +15\gamma \left(5 \frac{b^4}{\rho^4} + 5 \frac{b^8}{\rho^8} - 19 \frac{b^{12}}{\rho^{12}}\right), \end{aligned} \quad (4.2)$$

and γ is given by $\gamma = \frac{1}{8}\zeta(3)\alpha'^3$. With these corrections of the metric, the dilaton is no longer constant, instead it is given by

$$\Phi_0 = -\frac{45}{8}\gamma \left(\frac{b^4}{\rho^4} + \frac{b^8}{2\rho^8} + \frac{b^{12}}{3\rho^{12}} \right). \quad (4.3)$$

There is also a correction to the ten-dimensional dilaton action [15, 16], given by

$$I_{dilatons} = -\frac{1}{16\pi G_{10}} \int d^{10}x \sqrt{g} \left[-\frac{1}{2} g^{\mu\nu} \partial_\mu \Phi \partial_\nu \Phi + \gamma e^{-\frac{3}{2}\Phi} W \right], \quad (4.4)$$

where W is given in terms of the Weyl tensor, and in our background $W = 180/\rho^{16}$ in units where $b = 1$. To the leading order in γ , the dilaton perturbation does not mix with the metric perturbation, so we can study the dilaton equation derived from the action (4.4) in the fixed metric background (4.1). In subleading order in γ , the term $\gamma e^{-\frac{3}{2}\Phi} W$ would generate a mixing of the dilaton and the graviton.

We now have all pieces needed in order to obtain the first order correction to the dilaton equation. We write $\Phi = \Phi_0 + f(\rho)e^{ikx}$, with Φ_0 given by (4.3), and expand $f(\rho)$ and k^2 in γ as

$$f(\rho) = f_0(\rho) + \gamma h(\rho), \quad k^2 = k_0^2 + \gamma \delta k^2. \quad (4.5)$$

Here $f_0(\rho)$ obeys the lowest order equation (2.6) and is a numerically given function, and k_0^2 are the eigenvalues numerically obtained from the solution of (2.6). The first order term of the differential equation obtained from the action (4.4) using the background (4.1) and (4.3) is given by

$$\begin{aligned} \rho^{-1} \frac{d}{d\rho} \left((\rho^4 - 1) \rho \frac{dh}{d\rho} \right) - k_0^2 h &= (75 - 240\rho^{-8} + 165\rho^{-12}) \frac{d^2 f_0}{d\rho^2} \\ &+ (75 + 1680\rho^{-8} - 1815\rho^{-12}) \rho^{-1} \frac{df_0}{d\rho} \\ &+ (\delta k^2 - 120k_0^2 \rho^{-12} - 405\rho^{-14}) f_0(\rho). \end{aligned} \quad (4.6)$$

With $f_0(\rho)$ and k_0^2 given, one may regard this as an inhomogeneous version of the equation (2.6). We solve this equation to determine $h(\rho)$ and δk^2 .

At large ρ the solution for the first order correction has the form

$$h = \rho^{-4} \sum_{n=0}^{\infty} b_{2n} \rho^{-2n}. \quad (4.7)$$

Since (4.6) is inhomogeneous for $h(\rho)$, one can always add to a solution $h(\rho)$ a constant multiple of the solution $f_0(\rho)$ to the corresponding homogeneous equation (2.6) to obtain

another solution. We use this freedom to set $b_0 = 0$. The first few coefficients are then given by:

$$\begin{aligned}
b_2 &= \frac{\delta k^2}{12} \\
b_4 &= \frac{14400 + 2\delta k^2 k_0^2}{384} \\
b_6 &= \frac{1344\delta k^2 + 100800k_0^2 + 3\delta k^2 k_0^4}{23040} .
\end{aligned} \tag{4.8}$$

We can now determine δk^2 by the same ‘‘shooting’’ method described above for each eigenvalue of k_0^2 and its corresponding eigenfunction $f_0(\rho)$. It turns out that, for each eigenvalue k_0^2 , there is a unique solution with h being regular at $\rho = 1$. The first few solutions are shown in Table 7.

state	$(-k_0^2)$	$(-\delta k^2)$	$\delta k^2/k_0^2$
0^{++}	11.59	89.75	7.74
0^{+++}	34.53	365.7	10.59
0^{++++}	68.98	809.8	11.74
0^{+++++}	114.9	1397	12.16
0^{+++++}	172.3	2122	12.32
0^{+++++}	241.2	2991	12.40

Table 7: Leading string correction to the 0^{++} glueball masses in QCD_3 . The first column gives the zeroth order supergravity result for the mass squared, the second column gives the coefficient of the leading string correction and the third column gives their ratio.

Recalling that the squared mass of each glueball states is given by

$$M^2 = - \left(k_0^2 + \frac{1}{8} \delta k^2 \zeta(3) \alpha'^3 + \dots \right) b^2, \tag{4.9}$$

we see that the leading stringy corrections to the 0^{++} glueball masses are

$$\begin{aligned}
M_{0^{++}}^2 &= 11.59 \times (1 + 0.97\zeta(3)\alpha'^3 + \dots)b^2 \\
M_{0^{+++}}^2 &= 34.53 \times (1 + 1.32\zeta(3)\alpha'^3 + \dots)b^2 \\
M_{0^{++++}}^2 &= 68.98 \times (1 + 1.47\zeta(3)\alpha'^3 + \dots)b^2 \\
M_{0^{+++++}}^2 &= 114.9 \times (1 + 1.52\zeta(3)\alpha'^3 + \dots)b^2 \\
M_{0^{+++++}}^2 &= 172.3 \times (1 + 1.54\zeta(3)\alpha'^3 + \dots)b^2 \\
M_{0^{+++++}}^2 &= 241.2 \times (1 + 1.55\zeta(3)\alpha'^3 + \dots)b^2 .
\end{aligned} \tag{4.10}$$

It is important to note that the relation between the radius R of the compactifying circle $\mathbf{R}^4 \rightarrow \mathbf{R}^3 \times \mathbf{S}^1$ and the location $\rho = b$ of the horizon also receives an α' -correction. Instead of (2.3), we now have [17]

$$b = \left(1 - \frac{15}{8}\zeta(3)\alpha'^3 + \dots\right) \frac{1}{2R}. \quad (4.11)$$

Therefore, in units of the ultraviolet cutoff $\Lambda_{UV} = (2R)^{-1}$, the glueball masses are expressed as

$$\begin{aligned} M_{0^{2++}}^2 &= 11.59 \times (1 - 2.78\zeta(3)\alpha'^3 + \dots)\Lambda_{UV}^2 \\ M_{0^{2+3+}}^2 &= 34.53 \times (1 - 2.43\zeta(3)\alpha'^3 + \dots)\Lambda_{UV}^2 \\ M_{0^{2+4+}}^2 &= 68.98 \times (1 - 2.28\zeta(3)\alpha'^3 + \dots)\Lambda_{UV}^2 \\ M_{0^{2+5+}}^2 &= 114.9 \times (1 - 2.23\zeta(3)\alpha'^3 + \dots)\Lambda_{UV}^2 \\ M_{0^{2+6+}}^2 &= 172.3 \times (1 - 2.21\zeta(3)\alpha'^3 + \dots)\Lambda_{UV}^2 \\ M_{0^{2+7+}}^2 &= 241.2 \times (1 - 2.20\zeta(3)\alpha'^3 + \dots)\Lambda_{UV}^2. \end{aligned} \quad (4.12)$$

Thus the glueball masses are indeed modified by the $\alpha' = (4\pi g_s N)^{-1/2}$ correction. The corrections are negative for all the 6 levels we computed and are of the order α'^3 . Therefore the glueball masses decrease as we decrease the 't Hooft coupling $\lambda = g_s N$. As we discussed in the introduction of this paper, this is in accordance with our expectation about the continuum limit of QCD.

At the same time, the α' corrections to the ratios of the masses appear to be smaller than the corrections to each glueball mass, suggesting that the corrections are somewhat universal. This may indicate that the good agreement between the supergravity computation and the lattice gauge theory results is not a coincidence but is due to small λ^{-1} corrections to the mass ratios. Obviously, with the given data, we cannot tell whether the stringy corrections for the mass ratios remain small in the continuum limit $\lambda \rightarrow 0$. It would be very interesting to see whether this trend continues in the subleading corrections in α' .

5 Summary and Discussion

In this paper, we computed the glueball masses of large N QCD in three and four dimensions by solving supergravity wave equations in the AdS black hole geometry. The supergravity approximation is valid for large N and large $\lambda = g_{YM}^2 N$ and therefore the results are to be compared with a fixed ultraviolet cutoff in the strong ultraviolet coupling regime.

We computed the ratios of the masses of the excited glueball states with the mass of the lowest state, as well as the ratio of masses of two different lowest glueball states. These ratios are in surprisingly good agreement with the available lattice data. We also computed the leading $\lambda^{-3/2}$ corrections to the glueball masses taking into account stringy corrections to the black hole geometry. We found that the corrections to the masses are in accordance with our expectation about the continuum limit of QCD. The corrections to the ratios of the masses appear to be smaller than the corrections to each glueball mass, suggesting that the corrections are somewhat universal.

The above computations can be generalized to higher spin glueballs. As noted previously, the graviton couples to the energy-momentum tensor and solving its equation of motion will give the masses of the 2^{++} glueball. In general the higher spin glueballs will correspond to operators that couple to massive string excitations. The dimensions of these operators are $\Delta \sim \lambda^{1/4}$ for large λ [8]. It would be interesting to see how to extrapolate this result to the continuum $\lambda \rightarrow 0$.

Another interesting issue is the existence of $SO(6)$ non-singlet states in supergravity. For large λ , their masses are of the same order as the $SO(6)$ singlet states we studied in this paper. In the continuum limit, $\Lambda_{UV} \rightarrow \infty$ and $\lambda \rightarrow 0$, those states should decouple. Presumably λ^{-1} corrections make them heavy.

Maldacena's conjecture reduces the problem of solving large N QCD in three and four dimensions to that of controlling the α' corrections to the two-dimensional sigma-model with the Ramond-Ramond background. In this paper, we have extracted information about glueballs in strongly coupled QCD using the α' -expansion of the sigma-model. It would certainly be interesting to understand properties of such a sigma-model better.

Acknowledgments

We would like to thank M. Chanowitz, D. Gross, A. Hashimoto, C. Morningstar, S. Sharpe and Y. Takei for discussions and communications. We also thank Harlan Robins and Jonathan Tannenhauser, who participated in an early stage of this project, for discussions. H.O. and Y.O. thank the Institute for Theoretical Physics at Santa Barbara for its hospitality. This work was supported in part by the NSF grant PHY-95-14797 and in part the DOE grant DE-AC03-76SF00098. In addition, H.O. and Y.O. are supported by the NSF grant PHY-94-07194 through the ITP. C.C. is a research fellow of the Miller Institute for Basic Research in Science.

References

- [1] J. M. Maldacena, “The Large N Limit of Superconformal Field Theories and Supergravity,” hep-th/9711200.
- [2] E. Witten, “Anti-de Sitter Space, Thermal Phase Transition, And Confinement in Gauge Theories,” hep-th/9803131.
- [3] A. Brandhuber, N. Itzhaki, J. Sonnenschein, S. Yankielowicz, “Wilson Loops in the Large N Limit at Finite Temperature,” hep-th/9803137.
- [4] S.-J. Rey, S. Theisen and J. Yee, “Wilson-Polyakov Loop at Finite Temperature in Large N Gauge Theory and Anti-de Sitter Supergravity,” hep-th/9803135.
- [5] M. Li, “’t Hooft Vortices on D-branes,” hep-th/9803252; “’t Hooft vortices and phases of large N gauge theory,” hep-th/9804175.
- [6] E. Witten, “Baryons and Branes in Anti-de Sitter Space,” hep-th/9805112.
- [7] D. J. Gross and H. Ooguri, “Aspects of Large N Gauge Theory Dynamics as Seen by String Theory,” hep-th/9805129.
- [8] S.S. Gubser, I.R. Klebanov, A.M. Polyakov, “Gauge Theory Correlators from Non-Critical String Theory,” hep-th/9802109.
- [9] E. Witten, “Anti-de Sitter Space and Holography,” hep-th/9802150.
- [10] I. R. Klebanov, “World Volume Approach to Absorption by Non-Dilatonic Branes,” hep-th/9702076; Nucl. Phys. **B496** (1997) 231.
- [11] S. S. Gubser, I. R. Klebanov and A. A. Tseytlin, “String Theory and Classical Absorption by Threebranes,” hep-th/9703040; Nucl. Phys. **B499** (1997) 217.
- [12] G. Horowitz and H. Ooguri, “Spectrum of Large N Gauge Theory from Supergravity,” hep-th/9802116, Phys. Rev. Lett. **80** (1998) 4116.
- [13] S. Ferrara, C. Fronsdal and A. Zaffaroni, “On $\mathcal{N} = 8$ Supergravity on AdS_5 and $\mathcal{N} = 4$ Superconformal Yang-Mills theory,” hep-th/9802203.
- [14] Y. Oz and J. Terning, “Orbifolds of $\text{AdS}_5 \times S^5$ and 4d Conformal Field Theories,” hep-th/9803167.
- [15] M. T. Grisaru, A. E. M. van de Ven and D. Zanon, “Four-Loop Beta Function for the $\mathcal{N} = 1$ and $\mathcal{N} = 2$ Supersymmetric Nonlinear Sigma-Model in Two Dimensions,” Phys. Lett. **B173** (1986) 423; M. T. Grisaru and D. Zanon, “Sigma-Model Superstring Corrections to the Einstein-Hilbert Action,” Phys. Lett. **B177** (1986) 347.
- [16] D. J. Gross and E. Witten, “Superstring Modifications of Einstein Equation,” Nucl. Phys. **B277** (1986) 1.

- [17] S. S. Gubser, I. R. Klebanov and A. A. Tseytlin, “Coupling Constant Dependence in the Thermodynamics of $\mathcal{N} = 4$ Supersymmetric Yang-Mills Theory,” hep-th/9805156.
- [18] H. J. Kim, L. J. Romans, P. Van Nieuwenhuizen, “Mass Spectrum of Chiral Ten-Dimensional $\mathcal{N} = 2$ Supergravity on S^5 ,” Phys. Rev. **D32** (1985) 389.
- [19] M. Gunaydin, N. Marcus, “The Spectrum of The S^5 Compactification of Chiral $\mathcal{N} = 2, D = 10$ Supergravity and The Unitary Supermultiplets of $U(2, 2/4)$,” Class.Quant.Grav. **2** (1985) L11.
- [20] T. Aoki, T. Kawai and Y. Takei, “Algebraic Analysis of Singular Perturbations – On Exact WKB Analysis,” Sugaku Exposition **8-2** (1995) 217, and references therein.
- [21] Y. Takei and T. Koike, private communication.
- [22] S. Das and S. Trivedi, “Three Brane Action and The Correspondence Between $\mathcal{N} = 4$ Yang Mills Theory and Anti-de Sitter Space,” hep-th/9804149.
- [23] S. Ferrara, M. A. Lledo, A. Zaffaroni, “Born-Infeld Corrections to D3 brane Action in $AdS_5 \times S_5$ and $\mathcal{N} = 4, d = 4$ Primary Superfields,” hep-th/9805082.
- [24] S. S. Gubser, A. Hashimoto, I. R. Klebanov, M. Krasnitz, “Scalar Absorption and the Breaking of the World Volume Conformal Invariance,” hep-th/9803023.
- [25] J. Mandula, G. Zweig, and J. Govaerts, “Covariant Lattice Glueball Fields,” Nucl. Phys. **B228** (1983) 109.
- [26] S. Mathur and A. Matusis, “Absorption of Partial Waves by Three-Branes,” hep-th/9805064.
- [27] M.J. Teper, “ $SU(N)$ gauge Theories in $(2 + 1)$ Dimensions”, hep-lat/9804008.
- [28] A.A. Tseytlin, “On Non-Abelian Generalization of Born-Infeld Action in String Theory,” Nucl. Phys. **B501** (1997) 41, hep-th/9701125.
- [29] N. Itzhaki, J. Maldacena, J. Sonnenschein and S. Yankielowicz, “Supergravity and The Large N Limit of Theories With Sixteen Supercharges,” hep-th/9802042.
- [30] G. Horowitz and A. Strominger, “Black Strings and p -Branes,” Nucl. Phys. **B360** (1991) 197.
- [31] P. Van Nieuwenhuizen, “The Complete Mass Spectrum of $d = 11$ Supergravity Compactified on S_4 and a General Mass Formula for Arbitrary Cosets M_4 ”, Class. Quantum Grav. **2** (1985) 1.
- [32] M.J. Teper, ”Physics from lattice: glueballs in QCD; topology; $SU(N)$ for all N ,” hep-lat/9711011.
- [33] C. Morningstar and M. Peardon, “Efficient Glueball Simulations on Anisotropic Lattices,” Phys. Rev. **D56** (1997) 4043, hep-lat/9704011.

- [34] O. Aharony, Y. Oz and Z. Yin, “*M* Theory on $\text{AdS}_p \times S^{11-p}$ and Superconformal Field Theories,” hep-th/9803051.

ERNEST ORLANDO LAWRENCE BERKELEY NATIONAL LABORATORY
ONE CYCLOTRON ROAD | BERKELEY, CALIFORNIA 94720

2
4