RECEIVED
LAWRENCE
RADIATION LABORATORY

LIBRARY AND
DOCUMENTS SECTION

UCRL-10154
c.2

University of California

Ernest O. Lawrence Radiation Laboratory

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

Berkeley, California

UCRL-10154
c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

submitted to Physical Review

UCRL-10154
Limited distribution

UNIVERSITY OF CALIFORNIA
Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

PRODUCTION OF Σ^+K^+ BY (π^+, p) INTERACTION AT 1170 Mev/c

Frank S. Crawford, Jr., Fernand Grard, and Gerald A. Smith

April 2, 1962

PRODUCTION OF Σ^+K^+ BY (π^+, p) INTERACTION AT 1170 Mev/c

Contents

Abstract v

I. Introduction 1

II. Experimental Details 1

III. Results 2

IV. Partial-Wave Analysis 5

Acknowledgments 11

References 12

PRODUCTION OF Σ^+K^+ BY (π^+, p) INTERACTION AT 1170 Mev/c

Frank S. Crawford, Jr., Fernand Grard, and Gerald A. Smith

Lawrence Radiation Laboratory
University of California
Berkeley, California

April 2, 1962

ABSTRACT

The reaction $\pi^+ + p \rightarrow \Sigma^+ + K^+$ has been studied at 1170 Mev/c in the 72 inch liquid hydrogen bubble chamber. An analysis of 251 events gave a total cross section of 0.205 ± 0.014 mb and values of $\alpha^0\overline{P}_{\Sigma^+} = 0.62 \pm 0.19$, $\alpha^+\overline{P}_{\Sigma^+} = -0.18 \pm 0.13$. Based on the differential cross section and Σ^+ polarization, the s-, p-, and d-wave amplitudes were determined:

$$s_{1/2} \equiv a = 0.061 \pm 0.016 \ (\mathrm{mb/sr})^{1/2},$$

$$\frac{p_{1/2} + 2p_{3/2}}{s_{1/2}} \equiv \frac{b \exp i\chi_b}{a} = (2.14 \pm 1.13) \exp (34.7 \text{ deg} \pm 29.2 \text{ deg})i,$$

$$\frac{p_{3/2} - p_{1/2}}{s_{1/2}} \equiv \frac{c \exp i\chi_c}{a} = -(1.46 \pm 1.78) \exp (20.5 \text{ deg} \pm 26.1 \text{ deg})i,$$

$$\frac{3d_{5/2} + 2d_{3/2}}{s_{1/2}} \equiv \frac{d \exp i\chi_d}{a} = -(1.02 \pm 0.97) \exp (-7.4 \text{ deg} \pm 60.2 \text{ deg})i,$$

and

$$\frac{d_{5/2} - d_{3/2}}{s_{1/2}} \equiv \frac{e \exp i\chi_e}{a} = -(0.72 \pm 0.63) \exp (91.9 \text{ deg} \pm 25.8 \text{ deg})i.$$

An alternative set of solutions exists for the transformations $\chi_b \rightarrow -\chi_b$, $\chi_c \rightarrow 180 \text{ deg} - \chi_c$, $\chi_d \rightarrow -\chi_d$, $\chi_e \rightarrow 180 \text{ deg} - \chi_e$. Based on extrapolation of s and p amplitudes obtained near threshold in a previous experiment in which angular distribution, polarization, and energy dependence were used, an attempt has been made to resolve the Minami ambiguity in the amplitudes of

this experiment. Currently available data of Σ^- and Σ^0 production by (π^-, p) interaction at 1220 Mev/c were used to test the charge-independence hypothesis. Within the limits of statistical accuracy, no evidence of a violation of charge independence was observed.

PRODUCTION OF Σ^+K^+ BY (π^+, p) INTERACTION AT 1170 Mev/c*

Frank S. Crawford, Jr., Fernand Grard,† and Gerald A. Smith

Lawrence Radiation Laboratory
University of California
Berkeley, California

April 2, 1962

I. INTRODUCTION

As a continuation of a previous analysis of the reaction $\pi^+ + p \rightarrow \Sigma^+ + K^+$ near threshold (1020 Mev/c)[1] the same reaction has been studied for an average incident laboratory-system momentum of 1170 Mev/c. From a scan of 35 000 pictures obtained with the 72-inch liquid hydrogen bubble chamber, approximately 400 events have been identified. After removal of events that would have led to experimental biases, a sample of 251 events was used for a determination of the s-, p-, and d-wave production amplitudes based on the observed angular distribution and sigma polarization. By comparing the s- and p-wave amplitudes obtained near threshold with those at 1170 Mev/c, an attempt has been made to remove the Minami ambiguity in the identification of these amplitudes. The present data, along with those available for the production of Σ^- and Σ^0 at about the same incident pion momentum, have been compared in order to test the charge-independence hypothesis.

II. EXPERIMENTAL DETAILS

The conditions under which the pictures have been obtained were very similar to those of the previous experiment.[1] The bubble chamber was operated in a magnetic field of 18 kgauss. The incident momentum was such that by energy loss by ionization in the hydrogen the events were produced in a momentum interval of 1153 to 1183 Mev/c. The beam resolution (half width

at half-maximum) was about 3 Mev/c. A precise knowledge of the beam resolution was not required, as it is in threshold analysis, since the energy dependence of the amplitudes has not been taken into account.

Using the same technique as described previously,[1] the beam contamination was found to be 3.0± 1.6% for the combined positron and muon background, and 2.1± 1.2% for the proton background. The kinematical upper limits of the energy of the δ ray as produced by 1170-Mev/c pions, muons, and positrons are 68, 113, and 1170 Mev/c respectively. The theoretical cross sections for producing δ rays of energy 68 through 113 Mev/c by 1170-Mev/c muons and positrons are 0.33 mb and 1.38 mb respectively.[2] The corresponding cross section for δ rays with energy greater than 113 Mev/c produced by positrons is 1.53 mb. To determine the proton contamination, the following total cross sections have been used: 15.3± 1.5 mb for the elastic (π^+, p) scattering,[3] and 25.2±0.8 mb for the elastic (p, p) scattering.[4] In order to remove the forward-scattering ambiguity, the two-prong events were accepted for a c.m. angle of the proton greater than 36 deg. The corresponding cross sections were evaluated, using the above elastic cross sections and elastic differential angular distributions available for neighboring incident energies.[3,4]

III. RESULTS

For an average pion energy of 1170 Mev/c, the laboratory-system momentum of the Σ^+ varies from about 550 Mev/c for those sigmas produced in the backward direction in the c.m. system, to about 950 Mev/c for those produced in the forward direction. In order to avoid errors arising from a bias against short sigmas, only events in which the sigma track length exceeds a certain limit were accepted for the final analysis, and an appropriate

correction factor was introduced for the interpretation of the experimental results. From a preliminary analysis of the data as a function of the various parameters upon which the scanning and measurement efficiencies can depend, the following selection criteria were adopted: (a) all acceptable beam tracks were required to enter through the bubble chamber thin window ($-4.0 \leq x_w \leq 10.0$ cm, $13.0 \leq z_w \leq 19.0$ cm), where x_w and z_w are the two coordinates normal to the incident beam direction, measured at the window; (b) it was required that the beam meet criteria in azimuth (108.5 deg $\leq \phi_w \leq 111.5$ deg) and dip ($-1.0 \leq \lambda_w \leq 2.5$ deg) at the window, so that acceptable beam tracks were well collimated in the chamber; (c) the fiducial volume was furhter defined by $-70.0 \leq y \leq 55.0$ cm, where y is the dimension parallel to the beam with y=0 at the center of the chamber; (d) the angle of the sigma decay plane with respect to the optic axes of the bubble chamber cameras was restricted to be greater than 20 deg for the $\Sigma^+ \rightarrow p + \pi^0$ decay mode, and greater than 10 deg for the $\Sigma^+ \rightarrow n + \pi^+$ decay mode, since events with decay angles smaller than these were observed to be lost by scanners because they were edge-on with the line of sight of the cameras and interpretable as two-prong events; (e) the forward-backward distribution of the decay pion in the Σ^+ rest frame for the $\Sigma^+ \rightarrow n + \pi^+$ decay mode showed no evidence of scanning loss for events satisfying the above criteria. For the $\Sigma^+ \rightarrow p + \pi^0$ decay mode, a cutoff of $+0.90 \leq \cos\psi$ (proton) ≤ -0.97 was required, where ψ is measured in the Σ^+ rest frame; this included all events in which the Σ^+ decayed at an angle of less than about 5 deg in the laboratory system; (f) the distribution of the lengths of the sigma tracks for the events satisfying the above criteria indicated significant loss for a length less than 3.5 mm in the $\Sigma^+ \rightarrow n + \pi^+$ decay mode and 7.5 mm for the $\Sigma^+ \rightarrow p + \pi^0$ decay mode. Accordingly, each accepted event was weighted with a factor equal to

$$+ \exp\left[\frac{\ell_{min} M_{\Sigma^+}}{3p_{\Sigma^+} \tau_{\Sigma^+}}\right] ,$$

representing the inverse of the probability for the observation of a sigma track with a length greater than ℓ_{min}. Here $\bar{\tau}_{\Sigma^+}$, the mean lifetime of the sigma (in 10^{-10} sec), has been taken equal to 0.75,[1] p_{Σ^+} is the laboratory-system momentum of the Σ^+, and M_{Σ^+} its mass (1189 Mev).

Figure 1 shows the production c.m. angular distribution, corrected for short sigma tracks. Figure 2 shows the angular distribution of the proton in the rest frame of the Σ^+ with respect to the normal to the production plane for the $\Sigma^+ \rightarrow p + \pi^0$ decay mode. A similar distribution for $\Sigma^+ \rightarrow n + \pi^+$ is given in Fig. 3. Figure 4 gives the decay asymmetry in $\Sigma^+ \rightarrow p + \pi^0$, measured with respect to the normal to the production plane as a function of the sigma c.m. production angle. The production angular distribution has been fitted satisfactorily within the statistical limits of the data by a polynomial of order five in $\cos\theta$. No significant improvement in the fit could be achieved with a polynomial of higher order. The data of Figs. 2 and 3 have been fitted to the function $(1 + a \cos\phi)$, using the maximum-likelihood method. Results obtained for the polarization are $\alpha^0 \bar{P} = + 0.62 \pm 0.19$, and $\alpha^+ \bar{P} = -0.18 \pm 0.13$. The total cross section for $\pi^+ + p \rightarrow \Sigma^+ + K^+$ at 1170 Mev/c was measured to be 0.205 ± 0.014 mb, using the information contained in Table I. The numbers of corrected events corresponding to the two decay modes were determined separately. By adding all events, each corrected by the weighting factor

$$+ \exp\left(\frac{\ell_{min} M_{\Sigma^+}}{3p_{\Sigma^+} \tau_{\Sigma^+}}\right) ,$$

the number of events corrected for short sigma tracks was obtained. For

the $\Sigma^+ \rightarrow p + \pi^0$ decay mode, the number of events has been further corrected by assuming that the forward-backward angular distribution in the rest frame of the Σ^+ is isotropic. No such correction was required for the $\Sigma^+ \rightarrow n + \pi^+$ decay mode. Correction for the events whose decay planes made too small an angle with the optic axes was made by assuming a continuous distribution of the events as a function of that angle. Finally, the correction for scanning efficiency, determined separately for the two decay modes, was made. The resultant final corrected numbers of events correspond to a decay branching ratio of

$$\frac{(\Sigma^+ \rightarrow p + \pi^0)}{(\Sigma^+ \rightarrow p + \pi^0) + (\Sigma^+ \rightarrow n + \pi^+)} = 0.49 \pm 0.05,$$

in good agreement with the current best value of 0.50 ± 0.02.[5]

IV. PARTIAL-WAVE ANALYSIS

Baltay et al. have studied $\pi^+ + p \rightarrow \Sigma^+ + K^+$ at neighboring incident momenta of 1110 and 1220 Mev/c.[6] The angular distribution obtained at 1220 Mev/c by these workers was fitted satisfactorily within the accuracy of the statistics by Legendre polynomials P_0, P_1, P_2, and P_3 consistent with the presence of at least d waves. We fitted the data of Baltay et al. at 1220 Mev/c to a third-order polynomial in $\cos\theta$, with the following results: $d\sigma/d\Omega \propto 1 + (1.63 \pm 0.10)\cos\theta - (0.17 \pm 0.18)\cos^2\theta - (1.71 \pm 0.17)\cos^3\theta$ with $\chi^2 = 2.7$ for seven degrees of freedom. The data of Fig. 1 at 1170 Mev/c have been fitted with polynomials of orders three, four, and five with the following results: third order, $d\sigma/d\Omega \propto 1 + (1.58 \pm 0.25)\cos\theta + (0.21 \pm 0.21)\cos^2\theta - (1.50 \pm 0.40)\cos^3\theta$, with $\chi^2 = 14.1$ for seven degrees of freedom; fourth order, $d\sigma/d\Omega \propto 1 + (1.71 \pm 0.32)\cos\theta + (0.74 \pm 0.72)\cos^2\theta - (1.62 \pm 0.43)$

$\cos^3\theta - (0.64\pm0.81)\cos^4\theta$, with $\chi^2 = 13.7$ for six degrees of freedom; fifth order, $d\sigma/d\Omega \propto 1 + (2.44\pm0.41)\cos\theta + (0.23\pm0.65)\cos^2\theta - (5.45\pm1.55)\cos^3\theta - (0.14\pm0.76)\cos^4\theta + (3.59\pm1.40)\cos^5\theta$ with $\chi^2 = 5.6$ for five degrees of freedom. No additional powers of $\cos\theta$ were required for a better fit. Considering the results of the χ^2 tests stated above on the data in Fig. 1, the presence of at least f waves cannot be excluded conclusively. Since only 95 events in the mode $\Sigma^+ \rightarrow p + \pi^0$ were observed, the polarization distribution vs $\cos\theta$ of Fig. 4 does not permit any further comment on the possible presence of f waves. Nevertheless the following analysis has been done under the assumption that only s, p, and d waves are present in the production mechanism. A determination of the s, p, and d partial-wave production amplitudes was carried out, using the production c.m. angular distribution and the distribution of the decay pion with respect to the normal to the production plane. The differential cross section $d\sigma/d\Omega$ and the sigma polarization P have been expressed in terms of the nonspin flip amplitude, $g(\theta)$, and the spin flip amplitude $h(\theta)$[7] as

$$d\sigma/d\Omega = g(\theta)^2 + h(\theta)^2 , \tag{1a}$$

and

$$P\, d\sigma/d\Omega = 2\mathrm{Im}h^*(\theta)\, g(\theta) \tag{1b}$$

where

$$g(\theta) = \sum_\ell \left[(\ell+1)f_\ell^+ + \ell f_\ell^-\right] P_\ell(\cos\theta) , \tag{1c}$$

$$h(\theta) = \sin\theta \sum_\ell \left[f_\ell^- - f_\ell^+\right] \frac{dP_\ell(\cos\theta)}{d\cos\theta} . \tag{1d}$$

Here $f_\ell^\pm$ are the partial-wave amplitudes for total angular momentum $J = \ell \pm 1/2$. In our analysis, the contribution of angular momentum ℓ larger than two has been neglected; that is, the nonspin flip and spin flip amplitudes take the form

$$g(\theta) = S_{1/2} + (2P_{3/2} + P_{1/2})\cos\theta + 1/2\,(3D_{5/2} + 2D_{3/2})(3\cos^2\theta - 1), \quad (2a)$$

$$h(\theta) = (P_{3/2} - P_{1/2})\sin\theta + 3(D_{5/2} - D_{3/2})\sin\theta\cos\theta. \quad (2b)$$

The s, p, and d wave amplitudes were determined by the maximum-likelihood method. Considering for each event the c.m. production angle, θ, and the angle ϕ between the decay pion and the normal to the production plane, $\hat{n}$, defined by

$$\hat{n} = \frac{\underset{\sim}{P}_\pi \times \underset{\sim}{P}_\Sigma}{|P_\pi|\,|P_\Sigma|\sin\theta_{\pi\Sigma}}, \quad (3)$$

the probability for the observation of an individual event is

$$f = A_0 + A_1\cos\theta + A_2\cos^2\theta + A_3\cos^3\theta + A_4\cos^4\theta$$
$$+ \alpha\cos\phi\sin\theta\,(A_5 + A_6\cos\theta + A_7\cos^2\theta + A_8\cos^3\theta), \quad (4)$$

where α is the decay asymmetry parameter of the Σ^+ (α^0 for $p\pi^0$ and α^+ for $n\pi^+$). The coefficients A_n are related to the partial-wave amplitudes by the expressions:

$$\begin{aligned}
A_0 &= a^2 + d^2/4 - ad\cos\chi_d + c^2, \\
A_1 &= 2ab\cos\chi_b - bd\cos(\chi_d - \chi_b) + 6ce\cos(\chi_c - \chi_e), \\
A_2 &= b^2 - c^2 + 9e^2 - 3d^2/2 + 3ad\cos\chi_d, \\
A_3 &= 3bd\cos(\chi_d - \chi_b) - 6ce\cos(\chi_c - \chi_e), \\
A_4 &= 9d^2/4 - 9e^2, \\
A_5 &= -2ac\sin\chi_c + cd\sin(\chi_c - \chi_d), \\
A_6 &= -2cb\sin(\chi_c - \chi_b) - 6ea\sin\chi_e + 3ed\sin(\chi_e - \chi_d), \\
A_7 &= -3cd\sin(\chi_c - \chi_d) - 6eb\sin(\chi_e - \chi_b), \\
A_8 &= -9ed\sin(\chi_e - \chi_d),
\end{aligned} \quad (5)$$

where

$$
\begin{aligned}
S_{1/2} &\equiv a \\
2P_{3/2} + P_{1/2} &\equiv b \exp[i\chi_b] \quad , \\
P_{3/2} - P_{1/2} &\equiv c \exp[i\chi_c) \, , \\
3D_{5/2} + 2D_{3/2} &\equiv d \exp[i\chi_d] \\
D_{5/2} - D_{3/2} &\equiv e \exp[i\chi_e] \, .
\end{aligned}
\tag{6}
$$

Using the probability function f, given in Eq. (4), the production amplitudes have been determined by maximizing the likelihood function,

$$
\mathcal{L} = \prod_{i=1}^{N=251} \frac{f^{(i)} \exp\left[\dfrac{\ell_{min} \times M_{\Sigma^+}}{3\, p_{\Sigma^+}^{(i)} \bar{\tau}_{\Sigma^+}}\right]}{\displaystyle\int_{-1}^{+1}\int_{-1}^{+1} f \exp\left[\dfrac{\ell_{min} \times M_{\Sigma^+}}{3\, p_{\Sigma^+} \times \bar{\tau}_{\Sigma^+}}\right] d\cos\theta \, d\cos\phi} \, , \tag{7}
$$

The decay asymmetry parameters α^0 and α^+ have been taken equal to +1 and zero respectively. Table II gives the set of amplitudes that fits our data. These results do not change appreciably if α^0 is assumed to have a value somewhat different from + 1.00. For comparison, Table II includes also the results obtained for $\alpha^0 = +0.8$. The differences from the previous results are small compared with the errors. It is interesting to mention that the value of $\alpha^0 = +1.0$ fits the experimental data better than $\alpha_0 = +0.8$, as also was the case in the previous experiment. In Table III are found the A_n, as computed from Eq. (5). An alternative solution exists also for χ_b replaced by $-\chi_b$, χ_c replaced by 180 deg - χ_c, χ_d by - χ_d, and χ_e by 180 deg $-\chi_e$. The absence of an a priori energy-dependent representation of the amplitudes outside the threshold region has led to a determination of values of the amplitudes averaged over the energy interval investigated. The angular distribution based on the amplitudes is in complete agreement with the solid curve of Fig. 1,

obtained by an independent maximum-likelihood fit. In the same manner, the average polarization, $a^0\overline{P}$, has been computed from the amplitudes of Table II. The following value, $a^0\overline{P} = 0.55$, has been obtained, also in agreement with the average polarization, $a^0\overline{P} = 0.62^+ - 0.19$, determined from the separate maximum-likelihood fit to the form $(1+a\cos\phi)$, with the data of Fig. 2. The solid curve of Fig. 4 represents the s, p, and d amplitude prediction for $P(\theta)$, in agreement with the data. The form of the curve is given by the following ratio.

$$P(\theta) =$$

$$\sin\theta\,[-(0.0064\pm0.0055) - (0.0157\pm0.0167)\cos\theta - (0.0211\pm0.0191)\cos^2\theta + (0.0243\pm0.0312)\cos^3\theta\,,$$

divided by

$$(0.0164\pm0.0086) + (0.0266\pm0.0151)\cos\theta + (0.0094\pm0.0064)\cos^2\theta - (0.0256\pm0.0156)\cos^3\theta - (0.0087\pm0.0155)\cos^4\theta,$$

with $a^0 = +1.0$.

It must be noted that the interpretation of the amplitudes of Table II in terms of individual partial-wave amplitudes is ambiguous. It is well known that the set of amplitudes obtained by a Minami transformation followed by the complex conjugation operation corresponds to the same angular distribution and polarization (including the sign);[7,8] in this experiment this means that the amplitudes $s_{1/2}$, $p_{1/2}$, $p_{3/2}$, $d_{3/2}$, $d_{5/2}$, as obtained by the angular distribution and polarization, could equally represent amplitudes $p_{1/2}^*$, $s_{1/2}^*$, $d_{3/2}^*$ $p_{3/2}^*$, and $f_{5/2}^*$ respectively. Only with additional dynamical information can one attempt to find the correct set of solutions. That is, assuming that the amplitudes are reasonably well behaved and continuous functions of energy, one could eliminate one set of solutions if that set were in considerable disagreement with those extrapolated from a neighboring energy. In an

attempt to do this, the s- and p-wave amplitudes determined near threshold in the previous experiment[1] have been extrapolated to 1170 Mev/c, using the assumed energy-dependence of the amplitudes in that analysis, including the factor p_{th}^{*}/p_{1170}^{*}, where p^{*} represents the production c.m. momentum of the colliding pion. Since energy dependence was taken into account in the threshold experiment, no Minami ambiguity exists in the amplitudes that have been extrapolated to 1170 Mev/c. The results of this comparison are indicated in Table IV; here the extrapolated s and p amplitudes, along with the regular and Minami transform solutions at 1170 Mev/c, have been given. In each of the three cases, two sets of solutions are given corresponding to the transformations $\chi_b \rightarrow -\chi_b$ and $\chi_c \rightarrow 180$ deg $- \chi_c$. According to the previously adopted convention, the $S_{1/2}$ phases of the Minami solutions have been set to zero; phases of the amplitudes have been rotated so thatthe amplitudes are all positive. Since the errors on the amplitudes and phases are rather large, it is not possible to select that set of amplitudes determined at 1170 Mev/c, either regular or Minami transform, which unambiguously agrees with the extrapolated amplitudes.

Figure 5 represents the final corrected production differential cross section of this experiment (at 1170 Mev/c), along with the corrsponding minimum values as allowed by the charge-independence hypothesis[9] from the data obtained for Σ^- and Σ^0 at 1220 Mev/c by Crawford et al.[10] Based on this comparison, there is no evidence for a violation of the charge-independence hypothesis. A similar conclusion has been arrived at by Baltay et al. at 1220 Mev/c[6]

ACKNOWLEDGMENTS

The authors are pleased to acknowledge the encouragement of Professor Luis Alvarez in this experiment. We also wish to thank Jared Anderson, Dr. Hugo Camerini, Dr. Jack Fry, Dr. Myron L. Good, and Lester J. Lloyd for their aid in the experiment.

REFERENCES

* This work was done under the auspices of the U. S. Atomic Energy Comission.

† On leave from the Institut Interuniversitaire des Sciences Nucléaires, Belgium.

1. F. Grard and G. A. Smith, A Study of the Reaction $\pi^+ + p \rightarrow \Sigma^+ + K^+$ Near Threshold, Lawrence Radiation Laboratory Report, UCRL-10025 Jan. 19, 1962 (submitted to the Physical Review).
2. H. I. Bhabha, Proc. Roy. Soc. (London), A164, 257 (1938); B. Rossi, High Energy Particles (Prentice-Hall, Inc., Englewood Cliffs, 1952)
3. J. K. Kopp, A. M. Shapiro, and A. R. Erwin, Phys. Rev. 301, 123 (1961).
4. N. P. Bogachev and I. K. Vzorov, Doklady Akad. Nauk S.S.S.R. 99, 931 (1954); N. G. Meshcheryakov, N. P. Bogachev, and B. S. Neganov, Izvestia Akad. Nauk S.S.S.R. 19, 548 (1955); L. W. Smith, A. W. McReynolds, and G. Snow, Phys. Rev. 97, 1186 (1955).
5. D. A. Glaser, Proceedings of the Ninth International Annual Conference High-Energy Physics, Kiev, 1959 (Acadamy of Sciences, Moscow, 1959), p. 242.
6. C. Baltay, H. Courant, W. J. Fickinger, E. C. Fowler, H. L. Kraybill, J. Sandweiss, J. R. Sanford, D. I. Stonehill, and H. D. Taft, Revs. Modern Phys. 33, 374 (1961).
7. M. Nauenberg and A. Pais, Phys. Rev. 123, 1058 (1961).
8. S. Minami, Progr. Theoret. Phys. (Kyoto) 11, 213 (1954); S. Hayakawa, M. Kawaguchi, and S. Minami, ibid 332 (1954); ibid 12, 355 (1954).
9. J. J. Sakurai, Phys. Rev. 107, 908 (1957).
10. F. Crawford, R. Douglass, M. Good, G. Kalbfleisch, M. Stevenson, and H. Ticho, Phys. Rev. Letters 3, 394 (1959).

Table I. Scanning information and total cross section for $\pi^+ + p \rightarrow \Sigma^+ + K^+$ at 1770 Mev/c.

1.	Total number of pictures scanned.	34 738
2.	Total track length scanned, corrected for contamination, window criteria, and attenuation due to strong interactions. The latter correction is based on a total cross section of 29.7±0.44 mb, as given by Kopp et al. (reference 3), and made at the center of the chamber.	$(611.91 \pm 14.56) \times 10^3$ m
3.	Scanning efficiency for events meeting selection criteria on fiducial volume, decay plane orientation, window coordinates of beam track, and proton decay angle of 5 deg.	$p\pi^0$ = 94±3% $n\pi^+$ = 95±3%
4.	Number of observed events	$p\pi^0$ = 96 $n\pi^+$ = 155
5.	Number of corrected events	$p\pi^0$ = 217±22 $n\pi^+$ = 223±17
6.	Total cross section, based on above (2) and (5), and a target density of 0.0586 g=/cm^3.	0.205±0.014 mb

Table II. Maximum-likelihood solution to s-, p-, and d-wave amplitudes as defined by Eq. (6) for $a^0 = +1$, $a^0 = +0.8$, $a^+ = 0$, $\ell_{min}(\Sigma^+ \rightarrow p + \pi^0) = 0.75$ cm, and $\ell_{min}(\Sigma^+ \rightarrow n + \pi^+) = 0.35$ cm, with alternate solutions obtained by the transformations $\chi_b \rightarrow -\chi_b$, $\chi_c \rightarrow 180$ deg $-\chi_c$, $\chi_d \rightarrow -\chi_d$ and $\chi_e \rightarrow 180$ deg $- \chi_e$, and including the complete error matrix (angles are expressed in radians unless otherwise indicated).

	$a^0 = +1.0$	$a^0 = +0.8$
b/a =	$+2.14 \pm 1.13$	$+2.06$
χ_b =	$+34.7 \pm 29.2$ deg	$+35.8$ deg
c/a =	-1.46 ± 1.78	-1.31
χ_c =	$+20.5 \pm 26.1$ deg	$+26.9$ deg
d/a =	-1.02 ± 0.97	-1.11
χ_d =	-7.4 ± 60.2 deg	-11.9 deg
e/a =	-0.72 ± 0.63	-0.74
χ_e =	$+91.9 \pm 25.8$ deg	$+98.9$ deg
$\ell n\ \mathcal{L}$ =	-59.58	-59.81
a =	0.061 ± 0.016 (mb/sr)$^{1/2}$	

Error matrix

d/a	0.948	0.962	0.333	0.255	-0.165	0.187	-0.173	0.319
χ_d		1.105	0.389	0.288	-0.270	0.256	-0.002	0.361
e/a			0.396	0.242	-0.530	0.138	0.746	0.094
χ_e				0.203	-0.300	0.096	0.398	0.108
b/a					1.272	-0.282	-1.879	0.007
χ_b						0.264	0.342	0.146
c/a							3.130	-0.157
χ_c								0.206

Table III. The coefficients of Eq. (4) computed from Eq. (5) and the s, p, and d amplitudes of Table II: (all values in units of mb/sr).

$A_0 = 0.0164 \pm 0.0086$
$A_1 = 0.0266 \pm 0.0151$
$A_2 = 0.0094 \pm 0.0064$
$A_3 = -0.0256 \pm 0.0156$
$A_4 = -0.0087 \pm 0.0155$
$A_5 = -0.0064 \pm 0.0055$
$A_6 = -0.0157 \pm 0.0167$
$A_7 = -0.0211 \pm 0.0191$
$A_8 = 0.0243 \pm 0.0312$

Table IV. The s and p amplitudes of this experiment compared with those obtained by extrapolation from threshold. The regular solutions plus the Minami-transform solutions of this experiment are given; solution Regular I corresponds to the amplitudes and phases given in Table II, while Regular II corresponds to these amplitudes with the transformations $\chi_b \rightarrow -\chi_b$, $\chi_c \rightarrow 180$ deg $- \chi_c$ already made. The corresponding Minami-tranform solutions are given as Minami I and Minami II respectively. Solutions Regular A and Regular B correspond to the solutions obtained at threshold and these with $\chi_b \rightarrow -\chi_b$ and $\chi_c \rightarrow 180$ deg $- \chi_c$ respectively, extrapolated to 1170 Mev/c, using the assumed energy dependence of that experiment. The $S_{1/2}$ phases of the Minami solutions have been set to zero according to previously adopted convention. Amplitude phases have been rotated so that the amplitudes are all positive. There exists no Minami ambiguity in the threshold extrapolation solutions. Errors shown on the Minami I solutions are typical of errors on all Minami solutions.

	This experiment				Extrapolation	
	Regular I	Regular II	Minami I	Minami II	Regular A	Regular B
$a\left(\frac{mb}{sr}\right)^{1/2}$	0.061±0.016		0.102±0.094	0.036	0.083±0.014	
b/a	2.14±1.13		0.65 ±0.65	1.72	1.88 ±0.65	
χ_b^0	34.7±29.2	325.3±29.2	328.8±40.0	301.3	52.1±8.0	307.9±8.0
c/a	1.46±1.78		0.77	2.14	1.55 ±0.58	
χ_c^0	200.5±26.1	339.5±26.1	227.5	192.4	260.8±27.2	279.2±27.2



MU-26262

Fig. 1. Production c.m. angular distribution for the observed events corrected for sigma minimum-length cutoff. The solid curve represents the prediction of the s-, p-, and d-wave amplitudes of Table II, with the form $1+(1.71\pm0.32)\cos\theta+(0.74\pm0.72)\cos^2\theta-(1.62\pm0.43)\cos^3\theta-(0.64\pm0.81)\cos^4\theta$, with $\chi^2=13.7$ for six degrees of freedom. The dashed curve represents an independent maximum-likelihood fit to a fifth-order polynomial in $\cos\theta$ normalized to 336 events with the form $1+(2.44\pm0.41)\cos\theta+(0.23\pm0.65)\cos^2\theta-(5.45\pm1.55)\cos^3\theta-(0.14\pm0.76)\cos^4\theta+(3.59\pm1.40)\cos^5\theta$ with $\chi^2=5.6$ for five degrees of freedom.



MU-26263

Fig. 2. Decay proton angular distribution for $\Sigma^+ \rightarrow p + \pi^0$, measured in the Σ^+ rest frame with respect to $\hat{n}$. The curve represents a maximum-likelihood fit to the form $1 - \alpha^0 \bar{P} \cos\phi_p$ normalized to 148 events, including the correction for short sigma tracks.



MU-26264

Fig. 3. Decay neutron angular distribution for $\Sigma^+ \rightarrow n + \pi^+$, similar to that of Fig. 2. The fit to $1 - \alpha^+ \bar{P} \cos \phi_n$ has been normalized to 188 corrected events.



MU-26265

Fig. 4. Decay asymmetry with respect to $\hat{n}$ for $\Sigma^+ \rightarrow p + \pi^0$ based on 95 observed events. The solid curve represents the prediction of the s, p, and d amplitudes of Table II, with $\chi^2 = 1.6$ for one degree of freedom.



MU-26266

Fig. 5. The corrected absolute differential cross section of this experiment. The cross-hatched areas represent minimum Σ^+K^+ cross sections and their errors, as allowed by charge independence and the data of Crawford et al. at 1220 Mev/c.[10]

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.