

University of California

**Ernest O. Lawrence
Radiation Laboratory**

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

THE DECOMPOSITION OF NINE AMINO ACIDS DURING
CHROMATOGRAPHY ON PAPER

V. Moses

April 4, 1962

The Decomposition of Nine Amino Acids during Chromatography on Paper

It has recently been shown¹ that some decomposition of glycine takes place during chromatography on Whatman No. 4 paper using phenol-water and n-butanol-propionic acid-water as developing solvents. Losses of up to 38% were obtained in this system; these losses could be minimized by distillation of the phenol before use, and by pre-treatment of the paper with oxalic acid. Filter paper pre-treated in this way had a pH of 4.9, compared with 6.7 for untreated paper. This study has now been extended to include eight other amino acids.

Before use, the following amino acids were purified chromatographically on oxalic acid-treated Whatman No. 4 paper, using distilled phenol²-water followed by n-butanol-propionic acid-water³; DL-[2-¹⁴C]alanine (8.2 $\mu\text{c./mg.}$); DL-[4-¹⁴C]aspartic acid (10.7 $\mu\text{c./mg.}$); DL-[1-¹⁴C]glutamic acid (9.0 $\mu\text{c./mg.}$); [2-¹⁴C]glycine (8.9 $\mu\text{c./mg.}$); L-[¹⁴C]leucine (8.0 $\mu\text{c./mg.}$); DL-[3-¹⁴C]phenylalanine (21.1 $\mu\text{c./mg.}$); DL-[3-¹⁴C]serine (1.7 $\mu\text{c./mg.}$); DL-[2-¹⁴C]tyrosine (2.3 $\mu\text{c./mg.}$); and D-[4,4'-¹⁴C]valine (3.1 $\mu\text{c./mg.}$).

Each amino acid was eluted with water, evaporated to dryness under reduced pressure at 40°, and redissolved in water (0.1 N-HCl in the case of tyrosine). Aliquots of each amino acid solution containing 0.1 $\mu\text{c.}$ of ¹⁴C were spotted onto oxalic-acid treated, or untreated, Whatman No. 4 paper, and developed either in one or in two dimensions as shown in Table I. The positions of radioactive substances were determined by radioautography (Kodak Single-Coated Blue-Sensitive Medical X-Ray Film, exposed for 36 days). The radioactive areas were excised from the chromatograms and the radioactivity in each spot was counted automatically in the apparatus described by MOSES AND LONBERG-HOLM⁴.

The values reported in Table I are percentages of the total ^{14}C on each chromatogram present in substances other than the corresponding amino acid being investigated; the chemical nature of none of these other substances has been determined. Each of the amino acids investigated showed some breakdown during chromatography; in some cases as many as twenty decomposition products were detected. The extent of decomposition varied from zero with tyrosine in two instances, to over 14% with leucine in one system. Generally, glycine, leucine and phenylalanine showed the greatest degree of breakdown, and tyrosine the least; the other five substances were intermediate between these extremes.

Unlike the earlier findings with glycine¹, distillation of the phenol before use had no detectable effect on amino acid breakdown. Even glycine, the breakdown of which was earlier found to be much less extensive after distillation of the phenol, was unaffected. Presumably this is related to the quality of the undistilled phenol used on the two occasions: HUGGINS AND MOSES¹ employed British Drug Houses (Poole, England) Phenol Detached Crystals B.P.C., while in the present study the source was Mallinkrodt (St. Louis, Mo., U.S.A.) Phenol Liquefied Analytical Reagent.

Pretreatment of the chromatography paper with oxalic acid reduced the amount of decomposition by an average of 30% compared with untreated paper. The greater acidity of the oxalic acid-treated paper was probably responsible for much of this effect. However, oxalic acid specifically might also be of importance, by virtue of its binding of calcium and magnesium ions in the paper; there was some reduction in the extent of breakdown with oxalic acid-treated paper even when butanol-propionic acid-water (pH 2.5) alone was used as a solvent. The beneficial effect of oxalic acid treatment agrees with the earlier studies on the chromatographic breakdown of glycine¹.

-3-

The work described in this paper was sponsored by the United States Atomic Energy Commission.

Bio-Organic Chemistry Group,

V. MOSES

Lawrence Radiation Laboratory

University of California

Berkeley 4, Calif. (U.S.A.)

¹A.K. HUGGINS AND V. MOSES, Nature, 191 (1961) 668.

²O.J. DRAPER AND A.L. POLLARD, Science, 109 (1949) 448.

³A.A. BENSON, J.A. BASSHAM, M. CALVIN, T.C. GOODALE, V.A. HAAS AND W. STEPKA,
J. Am. Chem. Soc., 72 (1950) 1710.

⁴V. MOSES AND K.K. LONBERG-HOLM, submitted to Anal. Biochem.

TABLE I

BREAKDOWN OF NINE AMINO ACIDS DURING CHROMATOGRAPHY

Pretreatment of paper	Dimensions	1st Solvent	2nd Solvent	Extent of breakdown of amino acid (%)									
				Alanine	Aspartic Acid	Glutamic Acid	Glycine	Leucine	Phenyl-alanine	Serine	Tyrosine	Valine	Average for all amino acids
None	1	Ph.	---	2.0	3.4	4.8	10.4	5.7	3.0	2.4	0.6	1.6	3.8
"	1	Redist. Ph.	---	4.8	4.4	5.9	10.6	6.4	3.7	2.0	1.3	1.8	4.6
"	1	But-Prop.	--	3.5	3.2	3.7	9.2	10.7	7.7	1.0	2.5	3.3	5.0
"	2	Ph.	But.-Prop.	3.0	1.9	3.8	11.1	14.2	9.0	2.7	4.0	6.8	6.3
"	2	Redist. Ph.	But.-Prop.	3.4	2.3	4.9	12.6	12.9	11.2	2.2	4.6	6.3	6.7
"	2	But.-Prop.	Ph.	7.1	3.8	3.6	6.4	10.0	7.1	1.9	0.0	0.4	4.5
"	2	But.-Prop.	Redist. Ph.	3.9	3.7	4.3	8.9	9.8	5.5	0.8	0.5	1.6	4.3
Oxalic acid	1	Ph.	---	2.4	3.7	5.7	8.7	4.0	1.5	2.1	0.5	1.0	3.3
"	1	Redist. Ph.	---	1.9	3.2	5.2	8.0	3.2	2.8	1.3	0.4	0.7	3.0
"	1	But.-Prop.	---	2.8	2.3	3.7	6.7	8.4	7.6	1.2	3.3	2.2	4.3
"	2	Ph.	But.-Prop.	2.5	3.5	4.8	6.7	11.2	10.5	1.3	1.7	3.0	5.0
"	2	Redist. Ph.	But.-Prop.	1.5	2.2	2.4	6.6	8.4	10.4	1.4	3.3	2.8	4.3
"	2	But.-Prop.	Ph.	0.6	0.7	5.2	8.1	5.4	2.8	0.9	0.6	0.2	2.7
"	2	But.-Prop.	Redist. Ph.	0.9	7.3	1.9	9.0	2.8	2.3	1.1	0.0	0.8	2.9

Abbreviations: Ph., phenol-water; Redist.-Ph., redistilled phenol-water; But.-Prop., n-butanol-propionic acid-water.