

**University of California**  
**Ernest O. Lawrence**  
**Radiation Laboratory**

**TWO-WEEK LOAN COPY**

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

**Berkeley, California**

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

CERN Int. Conf. on High Energy Physics

UCRL-10220

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory  
Berkeley, California

Contract No. W-7405-eng-48

TEST OF ISOTOPIC SPIN CONSERVATION IN STRONG INTERACTIONS  
FROM AN EXPERIMENT LIMIT ON  $\sigma(d + d \rightarrow \text{He}^4 + \pi^0)$

John A. Poirier and Morris Pripstein

May 25, 1962

TEST OF ISOTOPIC SPIN CONSERVATION IN STRONG INTERACTIONS FROM  
AN EXPERIMENT LIMIT ON  $\sigma(d + d \rightarrow He^4 + \pi^0)^*$

John A. Poirier and Morris Pripstein

Lawrence Radiation Laboratory  
University of California  
Berkeley, California

May 25, 1962


To test the validity of isotopic spin (I-spin) conservation, the reaction



was looked for at the Berkeley 184-inch cyclotron. This strong interaction requires a change in I-spin from zero to one (since the I-spin of  $d$  and  $He^4$  are zero and the ordinary  $\pi^0$  is one) and is therefore forbidden by the conservation law. Preliminary results of our investigation have been published.<sup>1</sup> On the basis of a more comprehensive data analysis to be described below, we obtain an upper limit for the differential cross section of this reaction, in the center-of-mass system, of

$$\frac{d\sigma^{c.m.}}{d\Omega} (d + d \rightarrow He^4 + \pi^0) < 1.5 \times 10^{-34} \text{ cm}^2/\text{sr}. \quad (2)$$

at about  $\Theta_{c.m.} = 90$  deg. The data are consistent with no  $\pi^0$  production. By comparing this upper limit with the theoretical prediction of the cross section obtained by assuming that I-spin need not be conserved,<sup>2</sup> we conclude that I-spin is at least 99.6% conserved. The reaction



was also looked for at  $\Theta_{c.m.} = 65$ -deg; we obtain an upper limit for this

cross section in the center of mass system of

$$\frac{d\sigma}{d\Omega}^{\text{c.m.}} (d + d \rightarrow \text{He}^4 + \gamma) < 3.5 \times 10^{-34} \text{ cm}^2/\text{sr}. \quad (4)$$

The kinematics for reactions (1) and (3) for incident deuterons of 460 Mev laboratory kinetic energy are shown in Fig. 1. The experimental arrangement is shown in Fig. 2. Slits of lead were placed between the target and the first quadrupole to fix the lab angle of the alpha particles at  $8.7 \pm 0.6$  deg (indicated in Fig. 1) and to prevent the alpha-selecting magnet system from seeing the deuteron beam as it passed through the windows of the gas target. In order to accept the alphas from reaction (1) which are produced in the above lab angle interval, the magnet system was designed to have a momentum bite of approximately  $\pm 6\%$ . To minimize energy loss and multiple scattering, the entire path length of the alphas was in vacuum except for regions in the immediate vicinity of the target and the counters.

The target was 37.5-in. long, 6-in. in diameter, with 1/32-in. thick aluminum entrance and exit windows. It was filled with gas to a pressure of 390 psi (absolute) at liquid-nitrogen temperature. Fillings of hydrogen and deuterium were alternated.

The alpha particles were identified by means of effective momentum selection ( $p/Z$ ), time of flight and range (i.e.,  $T_1$ ,  $T_2$ ,  $T_4$ , Abs.,  $\bar{A}_1$ ), and  $dE/dx$  (i.e. lower level discrimination of the pulse heights of  $T_3$  and  $T_4$ ). The pulses from every counter of the telescope were recorded on a four-gun oscilloscope for each event. The electronics were tuned up by means of an alpha beam from the cyclotron and consistency checks were made periodically during the experiment. The R counter, shown in Fig. 2, was not used during the data-taking runs but was allowed to remain in position as part of the absorber arrangement.

The cross-section for deuterons on carbon,

$$\frac{d^2\sigma}{d\Omega dp} (d + C^{12} \rightarrow He^4 + \text{residue}) \quad (5)$$

was also measured by using the  $CH_2-H_2$  subtraction technique. The result is  $0.38 \times 10^{-30} \text{ cm}^2/\text{sr-Mev/c}$  for alpha particles of 1060 Mev/c lab momentum and lab angle of  $8-2/3$  deg. With the  $CH_2$  target we could show that our electronics did not saturate even though the production cross section for alpha particles and the background of deuterons were larger than with the deuterium target in place.

The data runs were cycled among four settings: the system set for 1275 Mev/c alphas produced at target center to observe reaction (1), and then for 1427 Mev/c alphas to observe reaction (3), first with deuterium and then with hydrogen in the target. The setting for 1275 Mev/c alphas was sensitive to the production of a neutral particle with a mass of 120 to 155 Mev while that for 1427 Mev/c was sensitive to a mass of 0 to 110 Mev. The calculations of the cross sections for reactions (1), (3), and (5) were, in reference 1, based on the data counts recorded by the electronics and included an error in the calculation of the solid angles. Since then, a comprehensive analysis of the oscilloscope film was undertaken. From these pictures the pulse height distribution and the timing of each counter output was measured. The traces were calibrated by triggering the oscilloscope from time of flight coincidence circuits set for (a) protons, (b) deuterons, and (c)  $He^3$ 's while using a scattered deuteron beam from the cyclotron; alpha calibration was obtained while using a scattered alpha beam from the cyclotron. A scan of the data film shows that all the events have a timing corresponding to that of alphas or deuterons (for a given magnetic rigidity, these particles have the same velocity); moreover, no pulses from the anti-counter,  $A_1$ , are seen. Thus, all the events have the timing and range of alphas.

However, an analysis of the pulse heights of the counters showed that only a small fraction of these events were, in fact, alphas. These pulse height distributions for the system set for reaction (1), are shown in Fig. 3. The spectra for the  $D_2$  and  $H_2$  target data were the same, within statistics, and have been combined in Fig. 3. The pulse height data are based on a scan of approximately one-third of all the events. The spectra of the data for reaction (3) show the same effect. The pulse height spectrum for counter  $T_1$  shows that it was sagging badly since it was located in a region of high background. For purposes of time-of-flight coincidences, however,  $T_1$  was shown to be counting alphas efficiently. The  $T_2$  spectrum has two well separated peaks with the higher pulse-height peak position coinciding with that of the alpha calibration peak, and the lower peak coinciding with calibrated deuteron spectrum. The outputs of  $T_1$ ,  $T_3$ , and  $T_4$  are correlated with that of  $T_2$ : for the events in which the  $T_2$  pulse height corresponds to that of alphas, the outputs of  $T_3$  and  $T_4$  also correspond to that for alphas, while for those events which have a deuteron pulse-height in  $T_2$  the pulse-height distribution for  $T_3$  and  $T_4$  are peaked about a pulse-height lower than that for alphas. The low pulse height edge of the distributions in  $T_3$  and  $T_4$  correspond to the lower-level discriminator cut-off.

The pulse height information indicates that most of the events which the electronics recorded as alphas were really deuterons. Since the timing of all the events was the same, the deuterons must have interacted near the final counters to produce particles which gave a large pulse-height in  $T_3$  and  $T_4$  and which had insufficient range to reach the anti counter,  $A_1$ . The only material in the beam upstream from and near the final counters was the R counter (0.063-in. plastic scintillator) and a 0.010-in. thick mylar window of the vacuum system. Further analysis shows that this effect must be due to singly charged particles

( $H^3$ , protons or inelastic deuterons) produced in a deuteron-carbon reaction in this material. Because of the very large number of deuterons scattered down our channel (see Table I) this effect dominated the alpha counting rate determined by the electronics.

The number of true alpha counts, determined from a scan of all the film data, is listed in Table I.

Table I. Counts per  $10^{13}$  incident deuterons.

P( $He^4$ ) Mev/c	Alpha Counts* (Oscilloscope Film Data)		Deuteron Counts** (Electronics Data)	
	D <sub>2</sub> Target	H <sub>2</sub> Target	D <sub>2</sub> Target	H <sub>2</sub> Target
1275 ( $\pi^0$ )	12.7 ± 0.77	9.1 ± 0.65	63 x 10 <sup>3</sup>	44 x 10 <sup>3</sup>
1427 ( $\gamma$ )	15.5 ± 1.10	10.5 ± 0.82	62 x 10 <sup>3</sup>	44 x 10 <sup>3</sup>

\*Errors are statistical

\*\*Counter telescope was not plateaued for counting deuterons;

therefore, numbers should be considered only as indicative.

Since the conservation of baryon number forbids the production of alpha particles from d-p collisions, the hydrogen data were treated as background and subtracted from the deuterium data to yield the net  $He^4$  signal from d-d collision. However, this signal is to be considered an upper limit for the following reasons:

1. It was experimentally determined that an alpha particle beam scatters into our channel preferentially from deuterium. Thus alpha contamination of the beam from the cyclotron or from deuteron interactions before the target would provide a net D<sub>2</sub>-H<sub>2</sub> counting rate.

2. Deuterons also scatter into our system preferentially from deuterium (see Table I). Thus if some of the deuterons interacted at the target walls to produce alphas, a net positive yield of alphas would result from a D<sub>2</sub>-H<sub>2</sub> subtraction.

3. Further data at different momenta (not shown in Table I) provide additional information on the background. The results of an analysis of these data are consistent with zero yield from reaction (1). We find, therefore, no evidence of non-conservation of isotopic spin. This also means that no evidence was found for the existence of the reaction  $d + d \rightarrow \text{He}^4 + \pi^0$ , where the  $\pi^0$  is a neutral meson of zero isotopic spin with a rest mass in the interval 120 to 155 Mev. If the  $\pi^0$  exists, then this reaction, which is a strong interaction, would not be forbidden by I-spin conservation and therefore should have been detected.

To get a firm upper limit for reaction (1), we assumed that the alpha counts determined from the oscilloscope film and listed in Table I for the  $D_2$  target minus those from the  $H_2$  target were due entirely to reaction (1). A new limit for the differential cross section in the center-of-mass angle interval  $\Theta_{c.m.} = 97 \pm 14$  deg was then calculated and the value is

$$\frac{d\sigma^{c.m.}}{d\Omega} (d + d \rightarrow \text{He}^4 + \pi^0) < 1.5 \times 10^{-34} \text{ cm}^2/\text{sr}. \quad (2)$$

for incident deuterons of 460 Mev laboratory kinetic energy. This value is a two standard deviation limit, the standard deviation being  $0.27 \times 10^{-34} \text{ cm}^2/\text{sr}$ . This value is also an upper limit on the cross section for the production of a  $\pi^0$  of the same mass as that of the ordinary  $\pi^0$ . To get an upper limit on the degree of non-conservation of I-spin in strong interactions, we compare the value in (2) with the theoretical prediction<sup>2</sup> for the cross section at our energies,

$$\frac{d\sigma^{Th.}}{d\Omega} = (380 \pm 50) \times 10^{-34} \text{ cm}^2/\text{sr},$$

which was computed with the assumption that

I-spin need not be conserved. From this ratio, increased by two standard deviations we find that the probability of non-conservation of I-spin is  $< 0.4\%$ , or alternatively, that I-spin in strong interactions is at least  $99.6\%$  conserved. The limit for reaction (3), mentioned previously, was obtained in the same manner

from the oscilloscope film data in Table I and is a two standard deviation limit, the standard deviation being  $0.61 \times 10^{-34} \text{ cm}^2/\text{sr}$ .

\*Work done under the auspices of the U. S. Atomic Energy Commission.

<sup>1</sup>John A. Poirier and Morris Pripstein, Phys. Rev. 122, 1917 (1961).

<sup>2</sup>K. R. Greider, Phys. Rev., 122, 1919 (1961).

#### Figure Captions

Fig. 1. Angle vs. momentum of  $\text{He}^4$  (lab).

Fig. 2. Experimental arrangement.

Fig. 3. Pulse height spectra determined from oscilloscope film.

■ - Experimental Data.

⊕ - Spectrum for events in which  $T_2$  pulse height occurs in lower peak (i.e.  $T_2$  p.h.  $\leq 0.9$  cm).

Histogram is the spectrum for events in which  $T_2$  pulse height occurs in higher peak (i.e.  $T_2$  p.h.  $\geq 1.0$  cm). Arrows indicate medians of the calibration pulse height spectra for d's and  $\text{He}^4$ 's obtained with d and  $\text{He}^4$  beams, respectively, from the cyclotron.




Fig. 1

MU - 22451





Fig. 2

MU-22452



EVENTS PER BIN VS. PULSE HEIGHT (cm)

Fig 3

MU-26879

UCRL-10220