UCRL-10337

University of California

Ernest O. Lawrence Radiation Laboratory

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

International Conference on High Energy Physics
CERN-July 1962.

UCRL-10337

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

THE REACTION $K^* + P \rightarrow K^* + N_{33}^*$

W. Chinowsky, G. Goldhaber, S. Goldhaber, W. Lee, and T. O'Halloran

June 27, 1962

UCRL-10337
June 27, 1962

Paper to be presented by Gerson Goldhaber at the
1962 High Energy Conference in CERN

THE REACTION $K^+ + P \rightarrow K^* + N^*_{33}$

W. Chinowsky, G. Goldhaber, S. Goldhaber, W. Lee, and T. O'Halloran

Lawrence Radiation Laboratory
University of California
Berkeley, California

We are in the process of studying the interaction of K^+ mesons on hydrogen at 1.96 Bev/c. The experiment was carried out in the 20-inch BNL hydrogen bubble chamber in the separated beam tuned to K^+ mesons. The momentum spread of the beam was about ±1%.

From our preliminary analysis we find a total cross section of

$$\sigma_{K^+H} = 18 \pm 3 \text{ mb.}$$

Of the interactions which are observed, we have concentrated on the reactions

$$K^+ + P \rightarrow K^+ + P + \pi^+ + \pi^- \qquad \text{(a)}$$

$$\rightarrow K^0 + P + \pi^+ + \pi^0 \qquad \text{(b)}$$

$$\rightarrow K^0 + N + \pi^+ + \pi^+ \qquad \text{(c)}$$

$$\rightarrow K^0 + P + \pi^+ \quad . \qquad \text{(d)}$$

The observed ratios for these, based on 151 measured and fitted events of type (a) and ~ 200 events of types (b), (c), and (d) combined, are

$$a:b:c:d: \;=\; 10:7:1:20$$

where we took account of the undetectable decay modes of the K^0. The partial cross section for processes (a), (b), (c), and (d) is

$$\sigma \;=\; 7 \pm 2 \text{ mb} .$$

If we ascribe the events occurring between $M_{K\pi} = 860 - 920$ to K^* production and $M_{P\pi} = 1180 - 1290$ to N^*_{33} production, then we find that out of 151 identified events

25 are in the "non-resonant" region
78 are in the K^* region
106 are in the N^*_{33} region,

and finally 62 of the above events are in the common K^* and N^*_{33} region. Thus about half or more of the interaction appears to go through the double resonant state.

In Fig. 1 we plot $M_{P\pi^+}$ $(T = 3/2)$ against $M_{K^+\pi^-}$ $(T = 3/2, 1/2)$, while in Fig. 2 we plot $M_{P\pi^-}$ $(T = 3/2, 1/2)$ against $M_{K^+\pi^+}$ $(T = 3/2)$, all from reaction (a). The projections on the various effective mass axes are also shown. The difference between these is very striking and demonstrates clearly that the K^* is a $T = 1/2$ resonance.

In Fig. 3 we plot $M_{P\pi^+}$ against $M_{K^o\pi^o}$, and in Fig. 4 $M_{P\pi^o}$ against $M_{K^o\pi^+}$ from reaction (b). This differs from reaction (a) in that the K^* can be formed with either pion. It is remarkable that the K^* seems to have disappeared—within the limited statistic—perhaps due to interference effects. In Fig. 5 we present the C.M. angular distribution of the K^* from reaction (a) with the incoming K^+, where the events were selected to correspond to the double resonance as discussed above. The K^* distribution is strongly forward peaked and consequently (C.M.) the N^*_{33} is backward peaked, suggestive of production in a peripheral collision.

In Figures 6 and 7 we plot the final two mass combinations M_{KP} against $M_{\pi\pi}$ for reactions (a) and (b). These do not appear to show any particular structure.

Figures 8 and 9 give two Dalitz plots for the 3-particle reaction (d). Here again the presence of both the K^* and the N^*_{33} are clearly observed, but here the π^+ is shared between them, giving rise to constructive interference effects. Figures 10, 11, and 12 give the corresponding mass plots for $M_{P\pi^+}$, $M_{K^o\pi^+}$, and M_{PK^o}.

We wish to take this opportunity to thank the many members of the staff of the Brookhaven National Laboratory for their very helpful attitude in making this experiment possible. In particular we would like to express our appreciation to Dr. H. Blewett, Dr. R. Shutt, Dr. J. Sanford, and Mr. Spiro and the AGS crew.

* This work was supported by the U. S. Atomic Energy Commission.


K⁺+p ⟶ (K⁺+π⁻)+(P+π⁺)
T=3/2 or 1/2, T=3/2
1500
1400
1300
1200
1100
Mpπ⁺ (MeV)
1238 MeV
890(MeV)
30
20
10
Number of events
40
30
20
10
Number of events
600
700
800
900
1000
1100
MK⁺π⁻ (MeV)
MUB-1175

Fig. 1


K⁺ + p
(K⁺ + π⁺) + (P + π⁻)
T = 3/2
T = 3/2 or 1/2
1500
1400
1300
1200
1100
M P π⁻ (MeV)
25
20
15
10
5
Number of events
600
700
800
900
1000
1100
M K + π⁺ (MeV)
15
10
5
Number of events
M p π⁻ (MeV)
MUB-1171

Fig. 2


$K^+ + p \rightarrow (K^0 + \pi^0) + (p + \pi^+)$
$T = \frac{3}{2}$ or $\frac{1}{2}$ $T = \frac{3}{2}$
1500
1400
1300
1200
1100
$M_{p\pi^+}$ (MeV)
Number of events
15
10
5
600
700
800
900
1000
$M_{K^0\pi^0}$ (MeV)
15 10 5
Number of events
MUB-1173

Fig. 3


K⁺+p→(K⁰+π⁺)+(p+π⁰)
T=3/2 or 1/2 T=3/2 or 1/2
1500
1400
1300
1200
1100
M pπ⁰ (MeV)
Number of events
10
5
600
700
800
900
1000
1100
M K⁰π⁺ (MeV)
10 5
Number of events
MUB-1174

Fig. 4



MU-27187

Fig. 5


$K^+ + p \rightarrow (K^+ + p) + (\pi^+ + \pi)$
1900
1800
1700
1600
1500
M_{K^+p} (MeV)
Number of events
15
10
5
0
300
400
500
600
700
$M_{\pi^+\pi^-}$ (MeV)
1800
1700
1600
1500
1400
M_{K^+p} (MeV)
20
15
10
5
0
Number of events
MUB-1172

Fig. 6.


K⁺+p → (p+K°) + (π⁺+π°)
1900
1800
1700
1600
1500
1400
M pK⁺ (MeV)
1800
1700
1600
1500
1400
M pK° (MeV)
15
10
5
Number of events
10
5
300
400
500
600
700
Mπ⁺π° (MeV)
MUB-1170

Fig. 7



MU-27188

Fig. 8



MU-27189

Fig. 9


890 MeV
K*
K⁺+p → K⁰+π⁺+p
Number of events
0
2
4
6
8
10
12
14
600
700
800
900
1000
1100
1200
M*Kπ (MeV)
MU-27190

Fig. 10



MU-27191

Fig. 11



MU-27192

Fig. 12

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.