UCRL 10645

University of California

Ernest O. Lawrence Radiation Laboratory

NEW ISOMERS OF ASTATINE-212

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

To be published in Phys. Review

UCRL-10645

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

NEW ISOMERS OF ASTATINE-212

W. Barclay Jones

January 28, 1963

New Isomers of Astatine-212 *

W. Barclay Jones

Lawrence Radiation Laboratory
University of California
Berkeley, California

January 28, 1963

The $_{83}Bi^{209}$ (α, n) $_{85}At^{212}$ reaction was investigated at the Crocker Laboratory 60-inch cyclotron of the University of California. Previous work on this reaction[1-4] reports that At^{212} has a half-life of 0.20 sec, and emits alpha-particles of 7.6 or 7.88 MeV.

In this experiment, the alpha decay energies were measured with a phosphorus-diffused-junction counter having an energy resolution of 30 keV. The spectra were observed at selected intervals between cyclotron beam bursts. The spectrum taken at a bombarding energy of 24 MeV is shown in Fig. 1 (the small peaks at 5.63, 6.04, 6.28, 6.78, and 8.78 MeV are due to the calibration source, Th^{228}). The results of the experiment are summarized in Table I. The half-life associated with each of these alpha energies was measured individually by time analysis of each pulse height.

A search was made for a gamma transition between the states responsible for the 7.82- and 7.60-MeV alpha-groups by means of detecting the conversion electrons; however, no such transition was observed. Less than 1% of the alpha activity could have a gamma decay in the energy range from 100 to 600 keV; however, a ≈63-keV transition was observed with a half-life of ~0.13 sec.[5] The data indicate that the level structure might be that shown in Fig. 2.

The 4+ and 5+ assignments for Bi^{208} are from theoretical computations by E. Wahlborn,[6] and are justified by experimental work.[7] The

shell-model configuration for nuclei in the region of At^{212} suggests that the neutron outside the closed shell[8] should be $2g_{9/2}$ and that the protons should be $(1h_{9/2})^3$; however, by the predictions of Pryce[9] this would lead to a ground state of 0- and an isomeric state of 9-. The 0- to 4+ transition is forbidden; so the assignment of 0- or 4+ does not appear to be justified. Alternately, as in Bi^{210}, the lower spin state for At^{212} could be 1-.

Figure 3 shows the relative excitation functions for these two isomers, as well as the absolute cross section for $Bi^{209}(\alpha, 2n)At^{211}$.[10] However, neither, the (α, n) cross sections relative to each other nor the absolute cross sections have been determined.

The hindrance factors were calculated from the empirical relation: $\log_{10} F = \log_{10} t_{1/2} - A_Z \omega_{eff}^{-1/2} - B_Z$, where A_Z and B_Z are the arithmetical means of corresponding values for the two adjacent even atomic numbers.[11]

FOOTNOTES AND REFERENCES

*Work supported by the U. S. Atomic Energy Commission.

1. M. Weissbluth, T. M. Putnam, and E. Segrè (unpublished work) reported by D. Strominger, J. M. Hollander, and G. T. Seaborg, Revs. Modern Phys. 30, 585 (1958).
2. M. M. Winn, Proc. Phys. Soc. (London) A67, 949 (1954).
3. J. C. Ritter and W. G. Smith, Phys. Rev. 128, 1778 (1962).
4. R. G. Griffioen and R. D. Macfarlane, in Lawrence Radiation Laboratory Report UCRL-10023, 1962 (unpublished).
5. F. S. Stephens and R. M. Diamond, Lawrence Radiation Laboratory, private communication.
6. S. Wahlborn, Nuclear Phys. 3, 644 (1957).
7. P. Mukherjee and B. L. Cohen, Phys. Rev. 127, 1284 (1962).
8. H. J. Mang, Phys. Rev. 119, 1069 (1960).
9. M. H. L. Pryce, Proc. Phys. Soc. (London) A65, 773 (1952).
10. E. L. Kelly and E. Segrè, Phys. Rev. 75, 999 (1949).
11. C. J. Gallagher and J. O. Rasmussen, J. Inorg. Nucl. Chem 3, 333 (1957).

Table I. Alpha-decay energies and half-lives for At^{212}.

Alpha decay energy (MeV)	Half-life (sec)	Approximate relative abundance (%)	Hindrance factor
7.60	0.305	20	6200
7.66	0.305	80	1700
7.82	0.120	80	1600
7.83	0.120	20	9500

FIGURE LEGENDS

Fig. 1. Spectrum of alpha energies for 24-MeV alpha particles on bismuth.

Fig. 2. Proposed energy-level diagram for alpha decay of At^{212}.

Fig. 3. Excitation functions for $Bi^{209}(\alpha, xn)At$. Units for the cross sections for the two isomers of At^{212} are not the same. Data for the At^{211} cross section is taken from reference 9.



MUB-1585

Fig. 1.



Fig. 2.

MU-29163


At 212 cross section (arbitrary units)
At 211 cross section (mb)
Lab energy of incoming alpha particles (MeV)
(α,n)
(α,2n)
0
1
2
3
4
5
6
7
8
9
0
100
200
300
400
500
600
700
17
18
19
20
21
22
23
24
25
26
27
MU-29162

Fig. 3.