

University of California
Ernest O. Lawrence
Radiation Laboratory

**ALPHA DECAY PROPERTIES OF SOME
ERBIUM ISOTOPES NEAR THE 82-
NEUTRON CLOSED SHELL**

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA
Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

ALPHA DECAY PROPERTIES OF SOME ERBIUM ISOTOPES
NEAR THE 82-NEUTRON CLOSED SHELL

Ronald D. Macfarlane and Roger D. Griffioen

April 1963

ALPHA DECAY PROPERTIES OF SOME ERBIUM ISOTOPES
NEAR THE 82-NEUTRON CLOSED SHELL[†]

Ronald D. Macfarlane* and Roger D. Griffioen**

Lawrence Radiation Laboratory

University of California, Berkeley, California

April 1963

ABSTRACT

New alpha-emitting isotopes of erbium, lying near the 82-neutron closed shell, were produced by Nd¹⁴² + O¹⁶ bombardments at incident energies up to 151 MeV. The nuclides studied and their alpha decay properties are:

Nuclide	Q_{α} (MeV)	Half-Life	Alpha Branching Ratio
Er ¹⁵²	4.93 ± 0.02	10.7 ± 0.5 sec	$0.90 + 0.05$ - 0.20
Er ¹⁵³	4.80 ± 0.02	36 ± 2 sec	$0.95 + 0.05$ - 0.20
Er ¹⁵⁴	4.26 ± 0.02	4.5 ± 1.0 min	---

Alpha decay from erbium isotopes with $A < 152$ was not observed indicating that the alpha particle binding energies of these isotopes are considerably higher than those for the 84-, 85- and 86-neutron isotopes.

Alpha reduced widths, calculated from the results for Er¹⁵² and Er¹⁵³, were found to be higher than the corresponding dysprosium,

[†]This research was performed under the auspices of the U. S. Atomic Energy Commission.

*Present Address: McMaster University, Hamilton, Ontario, Canada

**Present Address: Calvin College, Grand Rapids, Michigan

holmium and terbium isotopes. Relative excitation functions were obtained for some Nd^{142} (^{16}O , x n) and Nd^{142} (^{16}O , p x n) reactions.

I. INTRODUCTION

Alpha radioactivity has been observed for a number of nuclides which lie near the 82-neutron closed shell in the rare earth region. It has been found previously that for a given element the alpha decay energy has a minimum value at $N = 82$, increases very sharply with neutron number to a maximum value at $N = 84$ and gradually decreases with increasing neutron number beyond $N = 84$. The alpha decay properties of the 84-neutron isotopes of the elements from $Z = 60$ to $Z = 67$ have been reported and for some elements alpha activities from isotopes with $N = 85$ to $N = 88$ have been observed^{1,2,3,4,5,6,7,8}.

The purpose of this paper is to report on results which were obtained for the 84-, 85- and 86-neutron isotopes of erbium ($Z = 68$). A preliminary report of these results has been given earlier⁹. Because of their proximity to the 82-neutron shell, there is strong evidence that these isotopes as well as most of the other known alpha emitters in the rare earth region possess a stable spherical or near-spherical

¹J. O. Rasmussen, Jr., S. G. Thompson and A. Ghiorso, Phys. Rev. 89, 33 (1953).

²K. S. Toth and J. O. Rasmussen, Nucl. Phys. 16, 474 (1960).

³M. Karrass, Ann. Acad. Sci. Fennicae A VI, 65 (1960).

⁴R. D. Macfarlane and T. P. Kohman, Phys. Rev. 121, 1758 (1961).

⁵R. D. Macfarlane, J. Inorg. Nucl. Chem. 19, 9 (1961).

⁶R. D. Macfarlane, Phys. Rev. 126, 274 (1962).

⁷K. Nurmi, P. Kauranen and A. Siivola, Phys. Rev. 127, 943 (1962).

⁸R. D. Macfarlane and R. D. Griffioen, Phys. Rev. (In Press).

⁹R. D. Macfarlane and R. D. Griffioen, Bull. Am. Phys. Soc. 6, 451 (1961).

ground state shape. It is hoped that a careful study of the alpha decay properties of nuclides in this region will yield useful information on detailed features of alpha decay in the absence of the effects of spheroidal deformation.

II. EXPERIMENTAL DETAILS

The nuclides Er^{152} , Er^{153} and Er^{154} were produced by Nd^{142} (O^{16} , x n) reactions using 75 to 151 MeV O^{16} ions from the Berkeley heavy-ion accelerator (Kilac). Samples for alpha particle analysis were prepared using two different techniques.

In one method, which was used for the longer-lived activities, reaction recoils ejected from the target were thermalized by helium and collected on a charged plate. The plate was then placed in a Frisch-grid ionization chamber for alpha particle analysis. The target assembly employed in these experiments was also used in a modified form to electrostatically collect recoils from the alpha decay of the erbium isotopes in order to establish parent-daughter relationships. The following is a general description of the procedure which was used. The Nd^{142} target was bombarded with O^{16} ions and recoils ejected from the target were collected on a charged plate as before. After bombardment, the helium was quickly pumped out of the target assembly and replaced by a new supply at 1 atm pressure. This was found to be necessary because it was observed that some recoils from the target were retained in the helium present during the bombardment for as long as a few minutes after bombardment. The electrode containing the target recoils was then relocated to a position in front of a second electrode in the target assembly which had been shielded during the bombardment so that

no target recoils would be collected on it. A negative voltage was then applied to this electrode and recoils were collected from the alpha decay of the activity produced in the bombardment. In vacuum, it was found that beta decay recoils could be electrostatically collected from the sample but alpha decay recoils could not because of their longer range. At a pressure of 1 atm of helium, alpha decay recoils were collected with good efficiency but it was no longer possible to collect beta decay recoils. Particular attention was given to insuring that no deposition of target recoils on the electrode used to collect alpha decay recoils took place since this would invalidate the results of the "recoil-milking" experiments. It was found that if the precautions mentioned above were taken, less than 10^{-6} of the level of activity collected during the bombardment appeared on the electrode used to collect the alpha decay recoils. Specific details of the target assembly and arrangement of the collecting electrodes are described in earlier papers^{6,8}.

Because of the short half-life of Er^{152} , a faster method for collecting and analyzing the activity was also used. This method has been useful for studying alpha activities whose half-lives are greater than 0.03 sec. Details of this procedure are reported elsewhere¹⁰ and only the general features will be given here. Reaction recoils ejected from the target were slowed down in helium at 1 atm pressure in the target assembly. They were then swept through a small orifice into a chamber under vacuum adjacent to the target assembly where they were deposited on the surface of a collector. Alpha particle spectra of the recoil samples were recorded at the same time the activity was collected

¹⁰R. D. Macfarlane and R. D. Grifficen, Nucl. Inst. and Meth. (In Press).

using a gold-surface barrier detector. For the short-lived activities ($t_{1/2} \leq 2$ min), the level of activity was allowed to build up to an equilibrium value before recording alpha particle spectra. This was convenient for the cross section measurements since no corrections for decay were required.

Targets consisted of rare earth oxides (~ 2 mg/cm²) deposited on a 0.006 mm thick aluminum backing. The neodymium oxide target that was used was enriched in Nd¹⁴² to 93.93%. The other neodymium isotopes were present in the following amounts: Nd¹⁴³, 2.46%; Nd¹⁴⁴, 2.59%; Nd¹⁴⁵, 0.29% and Nd¹⁴⁶, 0.72%. Aluminum absorbers of varying thicknesses were used to degrade the energy of the heavy ion beam. The data of Northcliffe were used to convert range in aluminum to energy¹¹. For O^{16} ions within the range of 75 to 160 MeV, the energy loss in the rare earth oxide target was estimated to be 5 MeV. This was obtained by extrapolating the range-energy curves for various elements given by Hubbard to the rare earth elements and oxygen¹².

Relative excitation functions were obtained from the intensities of the various groups in the alpha particle spectra after normalizing to a constant integrated beam current. The counting data were also corrected for recoil collection efficiency which was found to be dependent upon bombarding energy for the electrostatic collection method but not for the gas sweeping technique.

The gold-surface barrier alpha particle detector that was used in the gas sweeping experiments was made from n-type 1800 ohm-cm silicon and operated at a reverse bias of 20 v. The active surface area was

¹¹L. C. Northcliffe, Phys. Rev. 120, 1744 (1960).

¹²E. L. Hubbard (Unpublished Results).

9 cm². Energy calibration was obtained from the alpha particles of Po²¹⁰ (5.30 MeV)¹³, Eo¹⁵¹ (4.51 MeV)⁸, Dy¹⁵⁰ (4.23 MeV)¹⁴ and Tb¹⁴⁹ (3.95 MeV)¹.

III. RESULTS

Alpha particle spectra were obtained on samples collected from Nd¹⁴² + O¹⁶ bombardments at incident energies ranging from 75 to 151 MeV. In addition to the known alpha emitters of dysprosium and holmium, three new alpha groups were observed - two intense groups at 4.80 and 4.68 MeV alpha particle energy and a weak group at 4.15 MeV. None of these alpha groups was observed in Pr¹⁴¹ + O¹⁶ bombardments which produce isotopes of the rare earth elements up to holmium⁸. Excitation functions were obtained for these activities. In each case, the variation of yield with incident bombarding energy was found to correspond very closely with results previously obtained for heavy-ion induced compound nucleus reactions where only neutrons are evaporated^{8,15}. From these measurements it was concluded that the three new alpha groups are due to isotopes of erbium. Additional results were obtained which confirmed that these are erbium alpha emitters and information on mass assignments was also obtained. These results are discussed below.

Pr¹⁵²

Figure 1 shows alpha particle spectra of activity collected during Nd¹⁴² + O¹⁶ bombardments at various incident O¹⁶ energies using

¹³I. Perlman and F. Asaro, Lawrence Radiation Laboratory Report UCRL - 9524 (Unpublished).

¹⁴R. D. Macfarlane and D. W. Seegmiller (Unpublished Results).

¹⁵G. W. Simonoff and J. M. Alexander, Lawrence Radiation Laboratory Report UCRL - 10099 Rev (To Be Published).

the gas sweeping technique for collecting the recoils from the target. A strong group was observed at 4.80 MeV alpha particle energy which decays with a half-life of 10.7 ± 0.5 sec (Figure 2). The peak of the excitation function for this activity (Figure 3 A) falls at an excitation energy of 85 MeV which is very close to the value of 84 MeV previously observed for the $\text{Pr}^{141} (\text{O}^{16}, 6n) \text{Ho}^{151}$ reaction leading to the high spin isomer⁶. This suggests that the 10.7 sec activity is produced by the same type reaction which would yield Er^{152} as the product.

In our previous work on the holmium alpha emitters, mass assignments were made by identifying beta or alpha decay daughters which were collected electrostatically as recoils from a thin sample of the parent⁸. However, because of the short half-life of the 4.80 MeV erbium alpha activity, it was not possible to use this technique. Also, the alpha decay daughter of Er^{152} is Dy^{148} which, on the basis of alpha decay energy systematics, is not expected to exhibit a measurable alpha decay branch.

Efforts were made to establish a parent-daughter relationship through a possible Er^{152} beta decay branch. The beta decay daughter, Ho^{152} , has previously been studied and it has been shown that it exists as an isomer pair⁸. Alpha decay from both the ground and isomeric states has been observed. The high spin Ho^{152} isomer ($E_{\alpha} = 4.45$ MeV) was detected as a product in the $\text{Nd}^{142} + \text{O}^{16}$ bombardments as shown in Figure 1 but measurements of the decay rate of the 4.45 MeV Ho^{152} group in the presence of the 10.7 sec erbium activity showed that most if not all of this activity is produced directly by an $\text{O}^{16}, p, \frac{1}{2}n$ reaction. The low spin isomer of Ho^{152} was also detected (Figure 1) and an

excitation function for this activity was obtained. These results were reported in an earlier paper⁸. The yield curve was found to peak at an excitation energy of approximately 82 MeV and it fairly closely parallels the shape of the excitation function of the 10.7 sec erbium activity. This suggests that the low spin Ho¹⁵² isomer is formed for the most part by the beta decay branch of the 10.7 sec erbium activity which must, therefore, be due to Er¹⁵². The direct formation of the low spin Ho¹⁵² isomer by an O¹⁶, p 5 n reaction is expected to have a very low cross section with a maximum value at an excitation energy approximately 70 MeV. This is 15 MeV lower than the peak cross section for the high spin Ho¹⁵² isomer. These predictions are based on previous results obtained for the production of isomer pairs by heavy-ion-compound-nucleus reactions in this region^{6,8}.

Since Er¹⁵² is an even-even nuclide with ground state spin and parity of 0+, it is reasonable to expect that the beta decay branch would preferably populate the low spin isomeric state of Ho¹⁵². If it is assumed that the beta decay branch of Er¹⁵² populates only the low spin Ho¹⁵² isomer and the peak cross section for the low spin Ho¹⁵² is due solely to the beta decay of Er¹⁵², it is possible to calculate the alpha branching ratio of Er¹⁵² from the relative intensities of the Er¹⁵² and Ho¹⁵² (low spin) alpha groups. A value of 0.90 ± 0.05
 $- 0.20$ was obtained.

Er¹⁵³

The second prominent erbium alpha activity that was observed has an alpha particle energy of 4.67 MeV (Figure 1) and a half-life of 36 ± 2 sec (Figure 2). A yield curve for this activity was obtained which was found to peak at an excitation energy of 71 MeV (Figure 3 A).

This compares favorably with a value of 72 MeV previously observed for the $\text{Pr}^{141} (\text{O}^{16}, 5 \text{ n}) \text{Ho}^{152}$ reaction leading to the high spin isomer⁸. From these results, it was concluded that the probable mass assignment for the 36 sec erbium alpha activity is Er^{153} .

The alpha decay daughter of Er^{153} , Dy^{149} is not conveniently detectable because it does not possess an alpha decay branch. However, the beta decay daughter of Dy^{149} , Tb^{149} , has an alpha branch which has been well characterized¹. In order to determine whether the 36 sec erbium activity is due to Er^{153} , alpha decay recoils were electrostatically collected from a thin sample of this activity and analyzed for the presence of Tb^{149} . The samples were prepared by electrostatic collection of recoils from the Nd^{142} target at two incident O^{16} bombarding energies. At 103 MeV, the level of the 36 sec erbium activity is almost a maximum value and the 10.7 sec Er^{152} is barely detectable. Alpha decay recoils collected from samples at this bombarding energy clearly showed the presence of Tb^{149} alpha activity (Figure 4 A) which was identified by alpha particle energy (3.95 MeV) and half-life (4.1 h)^{1,2}. When samples were collected at a bombarding energy of 84 MeV where the level of the 36 sec erbium activity is reduced by a factor of 60, no Tb^{149} was detected in the alpha spectrum of the recoils from alpha decay. At the 103 MeV bombarding energy, only the 36 sec erbium alpha group has sufficient intensity to have produced the level of Tb^{149} alpha activity that was observed (Figure 1 D). The 36 sec activity must, therefore, be due to Er^{153} . Ho^{153} which is produced in $\text{Nd}^{142} + \text{O}^{16}$ bombardments can also form Tb^{149} as a product of alpha decay but the intensity of this alpha group is so small that only an immeasurably small amount of Tb^{149} can be formed from the alpha decay of this nuclide.

In order to obtain an estimate of the alpha branching ratio of Er^{153} , assumptions were made with regard to the expected peak cross section for the $\text{Nd}^{142} (\text{O}^{16}, 5 \text{ n})$ reaction relative to the $\text{Nd}^{142} (\text{O}^{16}, 6 \text{ n})$ reaction. It was previously observed that the ratio of the $\text{O}^{16}, 5 \text{ n}$ to $\text{O}^{16}, 6 \text{ n}$ peak cross sections for the $\text{Co}^{140} + \text{O}^{16}$ reaction is 1.22⁸. If the same value is used for the corresponding products of the $\text{Nd}^{142} + \text{O}^{12}$ reaction, it is possible to calculate the alpha branching ratio of Er^{153} . A value of 0.95 ± 0.05 was obtained.

Er^{154}

When the Nd^{142} target was bombarded with O^{16} ions at incident energies between 80 and 110 MeV, a weak alpha group was observed at 4.15 MeV alpha particle energy which decays with a half life of 4.5 ± 1.0 min. Figure 5 shows alpha particle spectra of target recoils collected by the electrostatic method at incident energies of 105 and 92 MeV where the 4.15 MeV group is clearly indicated. A yield curve was obtained for this alpha group which was found to peak at an excitation energy of 55 MeV (Figure 3). Previous results with the $\text{Ce}^{140} + \text{O}^{16}$ and $\text{Fr}^{141} + \text{O}^{16}$ reactions showed that the $\text{O}^{16}, 4 \text{ n}$ excitation function in this region peaks in the vicinity of 50 to 55 MeV excitation energy⁸ so that on the basis of its excitation function the 4.15 MeV erbium alpha activity is most likely due to Er^{154} .

Proof that this activity is due to Er^{154} was obtained from the experiments where alpha decay recoils were collected from samples containing the erbium alpha emitters. Figure 4 A shows the alpha particle spectrum of recoils collected from the alpha activity of samples produced by $\text{Nd}^{142} + \text{O}^{16}$ bombardments at an incident energy of

103 MeV and Figure 4 B at an energy of 8 1/2 MeV. In both spectra an alpha group with the characteristic decay properties of Dy¹⁵⁰ ($E_{\alpha} = 4.25$ MeV, $t_{1/2} = 7.4$ min)^{2,14} was observed. The 4.15 MeV alpha group is the only alpha activity produced at both of these energies with sufficient intensity to have yielded the amount of Dy¹⁵⁰ activity observed. This activity must, therefore, be associated with Er¹⁵⁴. The sample corresponding to the higher bombarding energy also showed the presence of Tb¹⁴⁹ which, as discussed above, is a grand-daughter of Er¹⁵³ alpha decay. The specific activity of Dy¹⁵⁰ is approximately a factor of 60 higher than Tb¹⁴⁹. This is the reason why the intensity of the Dy¹⁵⁰ alpha group is greater than the Tb¹⁴⁹ group even though the alpha decay rate of Er¹⁵³ is much higher than for Er¹⁵⁴. The level of the Dy¹⁵⁰ activity observed as alpha decay recoils is higher at the 8 1/2 MeV bombarding energy than at 103 MeV. This is consistent with the excitation function data and assignment of Er¹⁵⁴ for the 4.15 MeV alpha group.

It was not possible to obtain a meaningful estimate of the alpha branching ratio of Er¹⁵⁴ by comparing excitation function data from Ce¹⁴⁰ + O¹⁶ and Pr¹⁴¹ + O¹⁶ reactions with that obtained for the Nd¹⁴² + O¹⁶ reaction. The reason for this is that at the incident O¹⁶ energy where the O¹⁶, 4 n excitation function peaks, the compound nucleus formation cross section is very sensitively dependent upon bombarding energy. A slight shift of the O¹⁶, 4 n excitation function to higher energies for the Nd¹⁴² + O¹⁶ reaction due to increased neutron binding energies will greatly affect the magnitude of the peak cross section of the O¹⁶, 4 n reaction when compared with the O¹⁶, 5 n and O¹⁶, 6 n excitation functions.

IV. DISCUSSION

It would be more meaningful to defer a detailed discussion of the alpha decay properties of the erbium isotopes in relation to the systematics of the other rare earth alpha emitters until the study of alpha decay in the higher numbers of the rare earth elements has been completed. We can, however, mention some observations which refer specially to the erbium isotopes.

A. Alpha Decay Energies

These first comments refer to the variation of alpha decay energy with mass number. This was found to progressively increase with decreasing mass number from $N = 86$ to $N = 84$. In our experiments, although sufficient excitation energy was available to have yielded some Er^{151} and possibly a small amount of Er^{150} , there was no evidence of alpha decay from these nuclides. This indicates that the alpha decay energies of these nuclides are considerably less than for the 84-neutron isotope, Er^{152} , and is consistent with results from precise mass measurements of the lighter rare earth nuclides which show that the alpha decay energies of the 82- and 83-neutron isotopes are much lower than the values for the 84-neutron isotopes¹⁶.

Another observation concerning the alpha decay energies of the erbium isotopes is the relatively small difference between the alpha decay energies of Er^{152} and Er^{153} (130 kev) and the much larger difference between Er^{153} and Er^{154} (540 kev). These differences are observed to a somewhat lesser extent with the 84-, 85- and 86-neutron

¹⁶L. A. König, J. H. E. Mattauch and A. H. Wapstra, Nucl. Phys. 31, 18 (1962).

isotopes of dysprosium where they differ by 170 and 400 kev, respectively. For the nuclides below $Z = 64$, however, the alpha decay energies of 84-, 85- and 86-neutron isotopes for a given element differ by approximately the same amount (~ 500 kev). It appears that this change in the differences in the alpha decay energies of neighboring isotopes begins at $\sim Z = 66$ which suggests that it may be associated with the structural change which occurs between $Z = 64$ and $Z = 66$ where there is evidence of a minor closed proton shell^{1,2}.

B. Alpha Reduced Widths

Alpha reduced widths (δ^2) were calculated for Er^{152} and Er^{153} from the experimental results using the method of Rasmussen¹⁷. The alpha reduced width is defined in the following expression:

$$\lambda = \frac{\delta^2 P}{h}$$

where λ is the alpha decay constant, P is the barrier penetrability and h is Planck's constant. The nuclear potential used in the calculation of P is the sum of the coulomb potential, the centrifugal potential and the real part of the alpha-nucleus potential given by¹⁸

$$V(r) = -1100 \exp \left\{ - \left[r - \frac{1.17 A^{1/3}}{0.574} \right] \right\} \text{ MeV}$$

A value of 0.091 MeV was obtained for the reduced width of Er^{152} and 0.13 MeV for Er^{153} assuming $l = 0$ α -waves only. These values are a factor of 1.6 to 9 times higher than the corresponding isotopes of holmium, dysprosium and terbium which means that alpha decay is more

¹⁷J. O. Rasmussen, Phys. Rev. 113, 1593 (1959).

¹⁸G. Igo, Phys. Rev. Letters 1, 72 (1958).

favored for the erbium isotopes. A possible qualitative explanation for this is as follows. In the earlier work of Rasmussen and co-workers on the gadolinium, terbium and dysprosium alpha emitters, they observed a break in the alpha decay energy curve at $Z = 64$ which they interpreted as being due to a minor closed shell at $Z = 64^{1,2}$. This idea received support from the proton level sequence scheme proposed by Mottelson and Nilsson which shows a substantial energy gap between the $d_{5/2}$ proton level, a level which is presumably completely filled at $Z = 64$, and the next level, an $h_{11/2}$ state¹⁹. Additional experimental support for this particular level sequence has been obtained from the observation of long-lived isomeric states for Tb^{149} , Ho^{151} and $Ho^{152}^{6,8}$. The smaller values of δ^2 for dysprosium, holmium and terbium [$\delta^2 (Tb^{149}) < \delta^2 (Dy^{150}) < \delta^2 (Ho^{151}) < \delta^2 (Er^{152})$] could be the result of significant differences in the proton part of the nuclear wave functions of the initial and final states involved in the alpha decay process. As more protons are added to the $h_{11/2}$ level, however, these differences would tend to become smaller and as a result the reduced widths would be expected to become larger. Further work in progress on thulium, ytterbium, lutetium and hafnium alpha emitters near the 82-neutron closed shell may indicate more clearly how δ^2 varies as more protons are added beyond $Z = 64$.

The results obtained on the erbium alpha emitters are summarized in Table I.

¹⁹B. R. Mottelson and S. G. Nilsson, *Kgl. Danske Videnskab Selskab, Mat. Fys. Skrifter* 1, No. 8 (1959).

TABLE I

Summary of Results

Nuclide	Q_α (MeV)	Half-Life	Alpha Branch	δ^2 (MeV)
Er ¹⁵²	4.93 ± 0.02	10.7 ± 0.5 sec	$0.90 \begin{matrix} + 0.05 \\ - 0.20 \end{matrix}$	0.091
Er ¹⁵³	4.80 ± 0.02	36 ± 2 sec	$0.95 \begin{matrix} + 0.05 \\ - 0.20 \end{matrix}$	0.13
Er ¹⁵⁴	4.26 ± 0.02	4.5 ± 1.0 min	---	--

ACKNOWLEDGMENTS

We would like to thank Dr. J. Alexander and Dr. G. Simonoff for their contributions by way of informative discussions and also C. Corum, J. Johnson, R. Latimer, W. Goldsworthy, A. Wydler and M. Nakamura who assisted in developing the experimental system used in this work. The co-operation of the Hilac personnel is gratefully acknowledged.

FIGURE CAPTIONS

Figure 1 Alpha particle spectra of holmium and erbium isotopes produced by $\text{Nd}^{142} + \text{O}^{16}$ at (a) 151 MeV incident O^{16} energy, (b) 136 MeV, (c) 119 MeV and (d) 101 MeV. Those spectra were recorded under conditions of continuous bombardment, collection, and counting and correspond essentially to the equilibrium level of activity.

Figure 2 Alpha decay curves for Er¹⁵², Er¹⁵³ and Er¹⁵⁴.

Figure 3 $\text{Nd}^{142} + \text{O}^{16}$ excitation functions for the production of (a) Er^{152} , Er^{153} and Er^{154} and (b) the high spin isomers of Ho^{151} and Ho^{152} . The yields are corrected for alpha branching ratios with the exception of Er^{154} . The cross sections are in arbitrary units but the relative values are as shown in the figure. The yield scales for (a) and (b) are the same.

Figure 4 Alpha particle spectra of alpha recoils collected from samples containing Er^{153} and Er^{154} produced from $\text{Nd}^{142} + \text{O}^{16}$. For spectrum (a) the bombarding energy was 103 MeV and for spectrum (b) 84 MeV.

Figure 5 Alpha particle spectra of nuclides produced by $\text{Nd}^{142} + \text{O}^{16}$ at (a) 103 MeV O^{16} energy and (b) 92 MeV O^{16} energy. Samples were prepared by electrostatic collection of target recoils. Bombardment time was 2 min and samples were counted for a 2 min period 5 min after bombardment.




Fig. 1

MU-29383




Fig. 2

MU-29387




Fig. 3




Fig. 4

MU-29385




Fig. 5

MU-29384

