UCRL-11000

University of California

Ernest O. Lawrence Radiation Laboratory

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

SU(4) ASSIGNMENTS FOR THE VECTOR RESONANCES

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

For Phys. Rev. Letters only

UCRL-11000

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

SU(4) ASSIGNMENTS FOR THE VECTOR RESONANCES

Pekka Tarjanne and Vigdor L. Teplitz

September 3, 1963

SU(4) ASSIGNMENTS FOR THE VECTOR RESONANCES*

Pekka Tarjanne† and Vigdor L. Teplitz

Lawrence Radiation Laboratory
University of California
Berkeley, California

September 3, 1963

Two possible interpretations of the 730-MeV $K\pi$ resonance (κ)[1] have been proposed. Nambu and Sakurai[2] considered a scheme into which the κ could naturally be inserted should its spin and parity be 0^+. Minami considered a 1^- spin and parity assignment for the κ and discussed a scheme in which there are two octuplets of vector mesons.[3] The purpose of this note is to consider an alternative scheme, based on the 1^- assignment, in which all the vector resonances are interpreted as forming one multiplet corresponding to the adjoint (regular) representation of SU(4).

I. In the case of the stable baryons, the stable scalar bosons, and, to a lesser extent, the baryon resonances, it seems fairly well established that the dominant symmetry, if any, is that of SU(3). For the vector resonances, however, the problem of $\varphi - \omega$ mixing complicates the phenomenological assignment of a symmetry. From a theoretical point of view, of the known particles the vector resonances seem the nearest to forming a self-contained (bootstrap) system.[4] It therefore seems possible that this system should exhibit a different, and even greater symmetry. Here, we consider the possibility of that symmetry being that of the group SU(4).

II. Since SU(4) is of rank three, the interactions of the vector mesons, in our scheme, conserve three additive quantum numbers. We take them to be hypercharge Y and the third component of isotopic spin T_3, as in the SU(3) case, and a third quantity, which we call supercharge, Z. Arguments have been given that require the assignment of the vector resonances to the adjoint representation, which, for SU(4), is fifteen-dimensional.[4]

We construct the 15 particles by the method of Gell-Mann.[5] We consider four fictitious basic fields: p and n with $Z = 0$, $Y = 1$, $T_3 = \pm 1/2$, Λ with $Z = Y = T_3 = 0$, and X with $Z = 1$, $Y = T_3 = 0$. With this assignment for X, the members of the 15-dimensional representation are

$$
\begin{aligned}
\rho^+ &= p\bar{n} \\
\rho^o &= (p\bar{p} - n\bar{n})/\sqrt{2} \\
\rho^- &= n\bar{p} \\
A &= (\Lambda\bar{\Lambda} - X\bar{X})/\sqrt{2} \\
B &= (p\bar{p} + n\bar{n} - 2\Lambda\bar{\Lambda})/\sqrt{6} \\
K^{*+} &= p\bar{\Lambda} \qquad \overline{K^{*+}} = \Lambda\bar{p} \\
K^{*o} &= n\bar{\Lambda} \qquad \overline{K^{*o}} = \Lambda\bar{n} \\
\kappa^+ &= p\bar{X} \qquad \overline{\kappa^+} = X\bar{p} \\
\kappa^o &= n\bar{X} \qquad \overline{\kappa^o} = X\bar{n} \\
\lambda &= \Lambda\bar{X} \qquad \bar{\lambda} = X\bar{\Lambda}
\end{aligned}
\tag{1}
$$

The quantum numbers of the resonances are easily inferred from those of

p, n, Λ, and X . The ω and φ are linear combinations of A and B ; in contrast with the SU(3) case, both appear in the adjoint representation. The only undetected resonances in the scheme are λ and $\bar{\lambda}$, $Z = \pm 1$ singlets with charge and strangeness zero. It should be noted that the three $Z = +1$ ($Z = -1$) particles form the SU(3) - $3(\bar{3})$ multiplet, which in the eightfold way cannot be physically realized with integral strangeness.[5] In the SU(4) case, however, introduction of the quantity Z allows the existence of the 3 and $\bar{3}$, with $Z = \pm 1$, as we have seen by explicit construction.

III. We expect the new quantity supercharge to be a property only of the vector mesons; i.e. the baryons and pseudoscalar mesons have $Z = 0$. We therefore expect small cross sections for producing κ and λ singly, compared with those for the other vector mesons, since these processes do not conserve supercharge. The observed cross section for κ production is about one-tenth that for K^* production.[1] Since κ and λ decays do not conserve supercharge either, we expect smaller widths for them. Again this is in agreement with the experimental result for κ , which is[1]

$$\Gamma_\kappa < \frac{1}{3} \Gamma_{K^*} .$$

These properties make difficult the problem of observing the λ . In addition, because λ decay does not conserve supercharge, we are not able to estimate the branching ratio between $\lambda \to 2\pi$ and $\lambda \to 3\pi$. In spite of nonconservation of supercharge in both production and decay processes it is not necessarily fruitless to introduce the quantity. Presumably, it is

conserved in the interactions of the vector mesons among themselves and is thus meaningful to the extent that the vector mesons form a self-consistent (bootstrap) system.[4]

IV. From Eq. (1) we may derive mass formulae analogous to the Gell-Mann-Okubo (GMO) results in the SU(3) case. Assuming p and n to be degenerate, we have

$$2(\varphi + \omega + \rho) = 3(\kappa + K^*) \tag{2}$$

$$\lambda = \kappa + K^* - \rho \tag{3}$$

$$\rho + 3B = 4K^* \tag{4}$$

where we have used the particle symbol for the square of the particle mass. We note that Eq. (2) is independent of the amount of mixing of A and B necessary to produce ω and φ, while Eq. (4) is just the GMO formula for the SU(3)-8 contained in the SU(4)-15. The three mass relations are the result of neglecting three mass-splitting terms consistent with T_3, Y, and Z conservation occurring in the 20 and in the 84, in the decomposition of $15 \times \overline{15}$, which is

$$15 \times \overline{15} = 84 + 45 + \overline{45} + 20 + 15_S + 15_A + 1 \quad .$$

Similarly the GMO formula arises from neglecting a term in the 27, in the decomposition of $8 \times \overline{8}$.[5,6]

Inserting the experimental values for the resonance energies in Eq. (2) gives a discrepancy of 11(± 3)% between the left- and right-hand sides.[7] Using Eq. (3), we obtain for the λ energy

$$M_\lambda = 870 \text{ MeV} \quad , \tag{5}$$

while inserting Eq. (2) in Eq. (3) gives

$$M_\lambda = 950 \text{ MeV} \quad . \tag{6}$$

If the κ should be found to be 1^-, a search in this area for the λ would seem reasonable.

We are grateful to Professor Geoffrey F. Chew and Professor Stanley Mandelstam for helpful comments. We wish to thank Dr. David L. Judd for his hospitality at the Lawrence Radiation Laboratory.

REFERENCES

* Work done under the auspices of the U. S. Atomic Energy Commission.

† Present address: Tekn. fys. laitos, Otaniemi, Finland

1. G. Alexander, G. R. Kalbfleisch, D. H. Miller, and G. A. Smith, Phys. Rev. Letters 8, 447 (1962).

2. Y. Nambu and J. J. Sakurai, Phys. Rev. Letters, 11, 42 (1963).

3. S. Minami, Some Considerations about the K^* (730 MeV). Louisiana State University preprint 1963.

4. R. E. Cutkosky, J. Kalckar, and P. Tarjanne, in Proceedings of the International Conference on High-Energy Physics, Geneva 1962 (CERN, Geneva, 1962), p. 653; R. E. Cutkosky, Phys. Rev. 131, 1888 (1963).

5. M. Gell-Mann, Phys. Rev. 125, 1067 (1962); Y Ne'eman, Nucl. Phys. 26, 222 (1961).

6. S. Okubo, Progr. Theor. Phys. 27, 949 (1962).

7. M. Roos, Rev. Mod. Phys. 35, 314 (1963).

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.