UCRL-11137

University of California

Ernest O. Lawrence Radiation Laboratory

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Phys. Rev.

UCRL-11137

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

A SEARCH FOR Y_2^* RESONANCES

Margaret H. Alston, Angela Barbaro-Galtieri,
Arthur H. Rosenfeld, and Stanley G. Wojcicki

January 7, 1964

A Search for Y_2^* Resonances*

Margaret H. Alston, Angela Barbaro-Galtieri,
Arthur H. Rosenfeld and Stanley G. Wojcicki†

Department of Physics and Lawrence Radiation Laboratory
University of California, Berkeley, California

January 7, 1964

ABSTRACT

A search has been made for a doubly charged $\Sigma\pi$ resonance produced in $\overline{K}N$ interactions. Data in hydrogen with incident K^- momentum of 1.51 and 1.70 BeV/c and in deuterium with a K^- momentum of 1.49 BeV/c show no evidence for any $T = 2$ resonance below a mass of 1900 MeV.

All these final states are complicated and hard to analyze. Bumps were indeed observed in the mass spectra of $\Sigma^{\pm}\pi^{\pm}$; however, we interpret them as the result of known $Y_{0,1}^{*}$ resonances and explain our data fairly simply without requiring any $T = 2$ resonance.

II. EXPERIMENTAL METHOD

For this experiment we used film of the 72-inch Bubble Chamber exposed between September 1961 and June 1962 to incident 1- to 2-BeV/c K^- mesons from the Bevatron.[5] The momentum spread at each setting is typically ±3%. Details of the exposure are given in Table I. The path length in hydrogen was obtained by counting (a) τ decays and (b) the total number of interactions and normalizing to the known total cross sections.[6] Results from the two methods agreed to within 10% at all momenta. In deuterium the path length was obtained by finding the number of 3-prong events that fit the kinematics of a τ decay and by checking the ionization of these events on the scanning table.

In hydrogen the interactions of interest appeared in the chamber as events with four outgoing prongs in which one prong showed a "kink". To make the scanning as uniform as possible, the event was accepted if the kink was more than 2mm from the vertex on the scanning table, whose magnification with respect to the bubble chamber is 2/3. All events were re-examined, and in addition, in the 1.70-BeV/c film all 4-prongs found during scanning were examined very carefully in all three views for kinks. In this way some spurious events were rejected and additional events found. In deuterium the acceptable event types were four or three outgoing prongs with a kink. The

event was classified as a "3-prong" if the spectator proton was less than 1mm long on the scanning table and was thus unmeasurable. Again the 3- and 4-prong events were reexamined for kinks. In this way we obtained a sample of events in deuterium and at 1.70 BeV/c in hydrogen, in which we consider the scanning efficiency to be 100%. The scanning efficiency at 1.51 BeV/c in hydrogen is estimated to be 92 ± 1%. The number of events found is shown in Table I. In deuterium about 60% of the events have three prongs.

After scanning, the events were measured on a Franckenstein measuring projector and processed through the PANAL, PACKAGE, EXAMIN, and SUMMEX programs.[7] All constrained kinematic fits were attempted in PACKAGE. The 4-prong events in hydrogen are subject to four equations of constraint at the production vertex, and can be unambiguously identified. However, if the spectator proton in deuterium is unmeasurable, the fit has only one constraint, and, if one further variable is unmeasurable, the event cannot be fitted. The 3-prong events were arbitrarily accepted if χ^2 was less than 10 and the proton momentum was less than 220 MeV/c.

III. RESULTS

A. Σ 3π Events in Hydrogen at 1.51 BeV/c

Mass histograms of the $\Sigma\pi$ systems in reactions (1) and (2) are shown in Fig. 1. In the $T_3 = 0$ distributions each event appears twice because two possible pions can be paired with the Σ. It is clear that the reactions are dominated by the production of $Y_0^*(1405)$ and $Y_0^*(1520)$. As a first attempt at explaining the data, we simply assume that each event will exhibit either a

$Y_0^*(1405)$ or $Y_0^*(1520)$ production. From the $(\Sigma\pi)^0$ mass distributions, we estimate that the observed ratio is about 3:1, with no background events. Of course, of all the $Y_0^*(1520)$ events produced, only 2/3 of 55% decay via $\Sigma^\pm\pi^\pm$,[8] so that the total $Y_0^*(1520)$ production is comparable with the total $Y_0^*(1405)$ production. Curves of the expected distributions were calculated by using the H4 program.[9] The solid curve shows the expected distribution if a $Y_0^*(1405)$ with a width (Γ) of 50 MeV is produced in 3/4 of the observed events, and in 1/4 a $Y_0^*(1520)$ with a width (Γ) of 20 MeV is produced. This gives a tolerable fit to the data for $T_3 = 0$. However, in the $T_3 = \pm 2$ distribution the fit is poor, and in particular the deviations of the observations from the predicted distribution is different in the different charge states. In the $\Sigma^+\pi^+$ state there is an excess of 40 to 50 events at about 1450 MeV, whereas the excess is at about 1530 MeV in the $\Sigma^-\pi^-$ distribution. To explain this effect we looked for other resonances in the final state.

A plot of the $\pi\pi$ and 3π mass distributions showed no "bumps". Figure 2 shows the mass distributions for $\Sigma\pi\pi$. In the $\Sigma^+\pi^+\pi^-$ and $\Sigma^-\pi^+\pi^-$ distributions, each event occurs twice. Again the distribution predicted by our previous assumption is shown as a solid curve. It can be seen that in the $T_3 = -1$ distributions, agreement is quite good, whereas in the $T_3 = +1$ states there is a pronounced excess of events at 1660 MeV, showing the production of $Y_1^{*+}(1660)$[10] and its subsequent 3-body decay. We estimate that the number of these events is 145 for the $\Sigma^+\pi^+\pi^-$ decay mode, and 73 for $\Sigma^-\pi^+\pi^+$, showing a branching ratio of 2:1. Note that charged Y^* decay into $\Sigma\pi\pi$ need not give 1:1 even in the absence of background.

We now investigate the effect of these 218 Y_1^{*+} events on the $\Sigma\pi$ mass distributions. The three possible modes of decay of the $Y_1^*(1660)$ to give $\Sigma + \pi + \pi$ are

(a) $Y_1^*(1660) \rightarrow \Sigma + \pi + \pi$, according to 3-body phase space

(b) $Y_1^*(1660) \rightarrow Y_0^*(1405) + \pi$
$\qquad\qquad\qquad\qquad\qquad\qquad \hookrightarrow \Sigma + \pi$ (100%)

and

$$\text{(c)}\quad Y_1^*(1660) \rightarrow Y_1^*(1385) + \pi$$
$$\hookrightarrow \Sigma + \pi\ (4\pm4\%)$$
$$\text{or}\quad \Lambda + \pi\ (96\%)\ .$$

First we want to show that channel (c) does not make an appreciable contribution to $\Sigma\pi\pi$ (π) final states. The total decay of $Y_1^*(1660)$ into the dominant $\Lambda\pi^+\pi^0$ mode is about 400 events at this incident momentum.[10, 11] It is not possible to determine how many of these $Y_1^*(1660)$ events decayed via the sequence (c). However, even if they all decay that way, the subsequent $Y_0^*(1385) \rightarrow \Sigma\pi/\Lambda\pi$ ratio is so small that the total contribution to the Σ channel is only about 16 events.[12, 13]

Possibilities (a) and (b) remain. The relevant kinematics are shown in Table II. Consider the sequence of events $K^- + p \rightarrow Y_1^*(1660) + \pi_1^-$. The Y_1^* then decays in one of two ways:

$$\text{(a)}\quad Y_1^*(1660) \rightarrow \Sigma + \pi_2 + \pi_3;\ P_\Sigma < 328\ \text{MeV/c}$$

$$\text{(b)}\quad Y_1^*(1660) \rightarrow Y_0^*(1405) + \pi_2;\ P\ \text{decay} = 196\ \text{MeV/c in the } Y_1^*(1660)\ \text{frame}$$
$$\hookrightarrow \Sigma + \pi_3;\ P\ \text{decay} = 144\ \text{MeV/c in the } Y_0^*(1405)\ \text{frame.}$$

Since the $Y_1^*(1660)$ is produced in a two-body reaction, the energy of pion 1 will be strongly peaked, the width of the distribution being the width of the Y_1^* combined with that of the incident K^- momentum distribution. In addition, since the Σ is moving very slowly, the invariant mass of the $\Sigma\pi_1$ combination will also be strongly peaked because the motion of the Σ will hardly smear the distribution. The central value of this enhancement will vary with the incident K^- momentum, but its central value and shape for a particular K^- momentum are almost independent of the decay mode of the $Y_1^*(1660)$. The central value is 1540 MeV for an incident K^- momentum of 1.51 MeV/c. The distributions of $M(\Sigma\pi_2)$ and $M(\Sigma\pi_3)$ of course depend upon the decay mode of the $Y_1^*(1660)$. If the decay follows phase space, π_2 and π_3 are indistinguishable, and both $M(\Sigma\pi)$

distributions peak at about 1430 MeV. For the case of $Y_1^*(1660) \rightarrow Y_0^*(1405) + \pi_2$, the π_2 will have a total energy of about 240 MeV. Since again the slowness of the Σ minimizes smearing, the $M(\Sigma\pi_2)$ distribution will be peaked at about 1440 MeV, with a width of about 100 MeV. The $M(\Sigma\pi_3)$ distribution will of course be that of the $Y_0^*(1405)$.

We next consider which charge states will exhibit the effects described. Since the $Y_1^*(1660)$ is produced only in the positive charge state, π_1 must be negative. Also for decay mode (b), π_2 must be positive, since Y_0^* is neutral. The charge states where enhancements are possible are shown in Table II. For the $T = 2$ channels, an enhancement should appear at 1540 MeV for $\Sigma^-\pi^-$ but not $\Sigma^+\pi^+$, and at 1430 MeV for $\Sigma^+\pi^+$ but not $\Sigma^-\pi^-$. The expected enhancements for $\Sigma^-\pi^-$ and $\Sigma^+\pi^+$ are shown in Fig. 1, each normalized to the observed number of $Y_1^{*+}(1660)$. The excess events observed in the $\Sigma^+\pi^+$ and $\Sigma^-\pi^-$ distributions agree in general with the predictions. The dotted curves show the expected $T_3 = \pm 2$ distributions when $Y_1^*(1660)$ production and decay via $Y_0^*(1405)$ is included. The ratio of $Y_0^*(1405)$ to $Y_0^*(1520)$ to $Y_1^*(1660)$ observed via $\Sigma\pi$ decay modes is 7:3:2. The modifications to the $T_3 = 0$ distributions are small and are not shown. One can see that, in general, agreement with the observed data is surprisingly good considering the simplicity of the model. Further, <u>no</u> $T = 2$ resonance is required to describe the data.

No branching ratio can be given for the decay of $Y_1^*(1660)$ into the two modes considered because $M(\Sigma\pi)$ distributions for the two cases are similar and also because of the large number of background events, many of which show the production of $Y_0^*(1405)$. The displacement of the $\Sigma^+\pi^-$ peak at 1405 and the broadening of the $\Sigma^-\pi^+$ peak is not understood, but is probably due to interference effects.

B. $\Sigma 3\pi$ Events in Hydrogen at 1.70 BeV/c

The mass histograms for $\Sigma\pi$ and $\Sigma\pi\pi$ in reactions (1) and (2) are shown in Figs. 3 and 4; Unfortunately, there are far less data at this momentum. These histograms show the same general features as at 1.51 BeV/c. Again the interaction is dominated by the production of $Y_0^*(1405)$ and $Y_0^*(1520)$. The solid curve shows the production and $\Sigma\pi$ decay of $Y_0^*(1405)$ and $Y_0^*(1520)$ in the ratio 3:1. The $M(\Sigma\pi\pi)$ distributions again show the production of $Y_1^{*+}(1660)$--about 20 events in both $T_3 = +1$ channels and no effect in the $T_3 = -1$ channel. The enhancement these events produce in the $M(\Sigma\pi)$ distributions for $T_3 = \pm 2$ is shown in Fig. 3. These center at 1440 MeV again for $\Sigma^+\pi^+$ and 1630 MeV for $\Sigma^-\pi^-$. It can be seen that the general shape of the distributions can be explained by our simple model in which $Y_0^*(1405)$; $Y_0^*(1520)$, and $Y_1^*(1660)$ productions and $\Sigma\pi$ decays are observed in the ratio of about 5:2:1. Again, no resonance in the $T = 2$ channel is required to explain the data.

C. $\Sigma\, 2\pi$ Events in Deuterium at 1.49 BeV/c

As already mentioned, the two reactions studied in deuterium were

$$K^- + \begin{matrix}(n)\\(p)\end{matrix} \rightarrow \Sigma^- + \pi^- + \pi^+ \ (+p) \qquad (3)$$

to study the $\Sigma^-\pi^-$ system, and for comparison,

$$K^- + \begin{matrix}(n)\\(p)\end{matrix} \rightarrow \Sigma^+ + \pi^- + \pi^- \ (+p)\ . \qquad (4)$$

The events of interest are those in which a Σ and two pions are produced by the interaction of the K^- with the neutron of the deuteron; the proton is merely a spectator. All 4-prong and 3-prong events that showed a kink in one track were analyzed, but only those peripheral events with a recoil proton momentum less

than 220 MeV/c were considered for the above reactions. A plot of the recoil momentum is a fairly good fit to the Hulthén distribution up to 220 MeV/c, although there are too many events at momentum greater than 100 MeV/c. However, a study of these events shows that the error is large and the visible range of the proton is always consistent with a lower momentum than that calculated. We therefore consider that all the selected events are examples of reactions (3) and (4), and that the impulse model is a good approximation at this incident K^- momentum.

The Dalitz plots for these reactions are shown in Fig. 5, and the projected histograms in Fig. 6. It can be seen that the production of $Y_0^*(1405)$ and $Y_0^*(1520)$ dominates this reaction also, and there appears to be some neutral $Y_1^*(1660)$ events. However we shall now show that there is <u>no</u> unexplained peaking in the $T_3 = -2$ $\Sigma\pi$ states.

In an attempt to parameterize these data, we assumed that resonances would be produced at 1405, 1520, 1660, and 1815 MeV,[14] and that there would also be background events. Our data appear to agree with the quoted widths (Γ) of 50, 50, and 120 MeV for $Y^*(1405)$, (1660), and (1815), respectively. However they indicate a width of at least 30 MeV for $Y_0^*(1520)$, and this value was used in the fit. No attempt was made to divide the (1815) resonance into two peaks at 1765 and 1815 MeV,[14] since any enhancement in this region is obviously small and is consistent with zero. Even with four resonances of fixed width and some background events, it is impossible to obtain a good fit to the data. Further, it is clear from the Dalitz plots and histograms that there are large interference effects in the final state because:

a. There is a paucity of events between the 1405- and 1520-MeV peaks in the $\Sigma^-\pi^+$ state.

b. The width of the 1520-MeV ($\Gamma = 30$) peak is broader than that found by other authors; i.e., we find $\Gamma = 16$ MeV.[8,14] This discrepancy is particularly pronounced in the $\Sigma^+\pi^-$ state and is too great to be entirely explained by the resolution, which is found to be about ±7 MeV for both Σ^+ and Σ^- events.

c. The branching ratios of the T = 0 resonances into $\Sigma^+\pi^-$ and $\Sigma^-\pi^+$ are not unity, which is the value expected for non interfering T = 0 or T = 1 resonances.

No attempt was made to least-squares fit the data, because the model used is obviously over-simplified, and there are many parameters that can be varied and rather few data. The solid curve shown in Fig. 6 is a typical calculated curve for no interference in the final state. The fit is poor; however, the general characteristics are correct, the deviations are presumably due to interference effects. No T = 2 resonance is apparent in the $\Sigma^-\pi^-$ mass spectrum. The ratios for the observed production and decay via the $\Sigma\pi$ mode of $Y_0^*(1405)$: $Y_0^*(1520)$: $Y_1^*(1660)$: $Y_0^*(1815)$: background is about 6 : 7 : 4 : 2 : 10.

Some indication of the severity of the interference effects can be obtained from the branching ratios ($\Sigma^+\pi^-/\Sigma^-\pi^+$) for the resonances since these should be unity if there is no interference. After corrections have been applied (see next section), this ratio is 2.0 ± 0.6 and 0.7 ± 0.2 for $Y_0^*(1405)$ and $Y_0^*(1520)$, respectively. The branching ratios for $Y_1^*(1660)$ and $Y_0^*(1815)$ are unity within the errors.

An interesting comparison can be made between our data and that of Galtieri et al.[14] They investigated the reaction $K^- + d \rightarrow K^- + p + \pi^-$ (+ p), where the interaction takes place on the neutron, and the proton is a spectator. The number of events can be compared directly to obtain the branching ratios

of the resonances into the $\Sigma\pi$ and $\bar{K}N$ channels, because the two experiments used the same film sample. The data are shown in Table III. The branching ratios are in fair agreement with values from other experiments. Namely, for $Y^*(1520)$ a value of 1.67 ± 0.2 was obtained by Ferro-Luzzi et al.[8] For $Y_1^*(1660)$, $\bar{K}N/\Sigma\pi$ branching ratios already reported are poorly known and somewhat contradictory; they are $< 1/6$,[10] 0.7 ± 0.35,[15] and 0.5 ± 0.25.[16] Inverting our value in Table III we find 0.14 ± 0.14, which tends to pull down the most likely estimate. As for $\Sigma\pi/\bar{K}N$ branching ratios in the $Y^*(1815)$ peak or peaks, Wohl[17] quotes $\leq 6\%$, based on data obtained in the K^-p reaction between 1.0 and 1.15 BeV/c. Our results (23 ± 25 or $31 \pm 33\%$) are consistent. For good values, we will have to await resolution of the problem of how many resonances are present.[14]

D. Cross Sections

The number of fitted events was corrected to take account of the mis-measured events and losses due to unobserved Σ decays. All events found were measured once if practical, and all events, that failed ($\approx 30\%$) were remeasured. By looking at the distribution of these remeasured events, we can estimate the number of events that would probably have fitted the "interesting" hypotheses had we continued remeasuring.

For the hydrogen events, the Σ lifetime and Σ-decay angular distributions were plotted. From these we calculated and corrected for the number of Σ decays that would not have been observed. The loss for Σ^+ is obviously greater than for Σ^- because of the small angles of the protonic decay and the shorter length of the Σ. Corrections at 1.7 BeV were difficult to estimate because of poor statistics. The total correction was estimated

as about 42% for Σ^+ and 12% for Σ^- at both incident momenta. A correction of 8% for scanning inefficiency was applied to the 1.5-BeV/c data.

It was impossible to estimate accurate corrections for the deuterium events because (a) the many unfittable background events made it hard for us to estimate a correction for badly measured events and (b) we could not calculate the correction for unobserved Σ events very well because of poor statistics. Since we feel that neither of these corrections should be very much different in deuterium than in hydrogen, we have assumed that the hydrogen corrections are adequate for our deuterium data. In addition, a correction of 3% was made for proton recoils with momentum greater than 220 MeV/c.

The cross sections obtained are shown in Table IV; the errors reflect statistics, the systematic errors discussed, and uncertainty in estimating the K^- pathlength (see Table I).

IV. CONCLUSION

An attempt has been made to find a $\Sigma\pi$ resonance with $T = 2$. Even though all the final states which are produced by K^-N interactions and which contain a doubly charged $\Sigma\pi$ state are extremely complicated, a relatively simple model not requiring a $T = 2$ resonance appears to explain all the available data.

ACKNOWLEDGMENTS

This work was done as part of the K72 experiment. We would like to thank Prof. L. W. Alvarez for his encouragement and support during this experiment and all the members of the group who assisted us. In particular we thank Prof. R. D. Tripp for useful discussions on the deuterium data, and G. Gilbert and B. Marten who did much of the scanning.

FOOTNOTES AND REFERENCES

*This work was done under the auspices of the U. S. Atomic Energy Commission.

†Now at L'École Polytechnique, Paris, France.

1. J. D. Dowell, W. Koch, B. Leontic, A. Lundby, R. Meunier, J. P. Stroot, and M. Szeptycka, Phys. Letters 1, 53 (1962).

2. G. R. Kalbfleisch, G. Alexander, O. I. Dahl, D. H. Miller, A. Rittenberg, and G. A. Smith, Phys. Letters 4, 225 (1963).

3. C. Baltay, J. Sandweiss, H. D. Taft, B. B. Culwick, W. B. Fowler, J. K. Kopp, R. I. Louttit, J. R. Sanford, R. P. Shutt, S. M. Thorndike, and M. S. Webster, Phys. Rev. Letters 11, 32 (1963).

4. M. H. Alston, L. W. Alvarez, M. Ferro-Luzzi, A. H. Rosenfeld, H. K. Ticho, and S. G. Wojcicki, in Proceedings of the 1962 International Conference on High Energy Physics at CERN (CERN, Geneva, 1962).

5. R. Hubbard, D. O. Huwe, G. R. Kalbfleisch, J. Kirz, D. H. Miller, J. B. Shafer, D. H. Stork, H. K. Ticho, and C. Wohl, A Variable Momentum, Separated K^- Beam, Lawrence Radiation Laboratory Report UCRL-10690, 1963 (unpublished).

6. V. Cook, B. Cork, T. F. Hoang, D. Keefe, L. T. Kerth, W. A. Wenzel, and T. F. Zipf, Phys. Rev. 123, 320 (1961).

7. For a summary of these programs and a list of references see A. H. Rosenfeld, Nucl. Instr. Methods 20, 422 (1963).

8. M. B. Watson, M. Ferro-Luzzi, and R. D. Tripp, Phys. Rev. 131, 2248 (1963).

9. M. W. Horovitz, Alvarez Group Memo randum 295, May 23, 1962 (unpublished).

10. A preliminary result from these data has been published: L. W. Alvarez, M. H. Alston, M. Ferro-Luzzi, D. O. Huwe, G. R. Kalbfleisch, D. M. Miller, J. J. Murray, A. H. Rosenfeld, J. B. Shafer, F. T. Solmitz, and S. G. Wojcicki, Phys. Rev. Letters 10, 184 (1963).

11. Darrell O. Huwe, Lawrence Radiation Laboratory, private communication.

12. P. Bastien, M. Ferro-Luzzi, and A. H. Rosenfeld, Phys. Rev. Letters 6, 702 (1961).

13. M. H. Alston, L. W. Alvarez, P. Eberhard, M. L. Good, W. Graziano, and S. G. Wojcicki, Phys. Rev. Letters 6, 698, 1961.

14. L. Barbaro-Galtieri, A. Hussein, and R. D. Tripp, Phys. Letters 6, 296 (1963).

15. P. L. Bastien and J. P. Berge, Phys. Rev. Letters 10, 188 (1963).

16. Gerald A. Smith in Proceedings of the Athens Topical Conference on Recently Discovered Resonant Particles, Athens, Ohio, 1963 (University of Ohio, Athens, Ohio, 1963).

17. Charles G. Wohl, Lawrence Radiation Laboratory, private communication.

Table I. Path length and number of events.

Momentum (BeV/c)	C. m. energy (MeV)	Target	Path length (events / μb)	No. of observed events Σ^+	No. of observed events Σ^-	No. of events fitting Reaction 1 and 2 or 4 or 3 Σ^+	No. of events fitting Reaction 1 and 2 or 4 or 3 Σ^-
1.51	2025	Hydrogen	5.2±0.4	816	749	676[a]	544[a]
1.70	2110	Hydrogen	1.1±0.1	261	209	201[a]	149[a]
1.49	2015	Deuterium	0.5±0.03	739[+]	1013[+]	265[b]	308[b]

[a]Reactions 1 and 2.

[b]Reactions 3 and 4.

Table II. Kinematics of Y_1^{*+} (1660) production and decay

Reaction	$\Sigma\pi$ combination	Possible $\Sigma\pi$ combinations		Central values	
		$Y_1^{*+} = \Sigma^+\pi^+\pi^-$	$Y_1^{*+} = \Sigma^-\pi^+\pi^+$	1.51 BeV/c (B^*=2025 MeV)	1.70 BeV/c (E^*=2110 MeV)
Y_1^*(1660) production	$\Sigma^\pm\pi_1^\mp$	$\Sigma^+\pi^-$	$\Sigma^-\pi^-$	1540	1630
Decay mode(a)	$\Sigma\pi_2$ or $\Sigma\pi_3$	$\Sigma^+\pi^+$	$\Sigma^-\pi^+$	1430	1430
		$\Sigma^+\pi^-$			
Decay mode(b)	$\Sigma^\pm\pi_2^+$	$\Sigma^+\pi^+$	$\Sigma^-\pi^+$	1440	1440
	$\Sigma^\pm\pi_3^\mp$	$\Sigma^+\pi^-$	$\Sigma^-\pi^+$	1405	1405

Table III. Branching ratio ($\Sigma\pi/\bar{K}N$) for decay of various Y^*'s produced in K^-d reactions. The errors here represent crude (rather than statistical) estimates.

Resonance	$K^- + n \rightarrow \Sigma^* + \pi^{\mp} + \pi^{\pm}$		$K^- + n \rightarrow K^- + p + \pi^-$		$\Sigma\pi/\bar{K}N$
	Corrected No. of $\Sigma^+\pi^- + \Sigma^-\pi^+$	Total $\Sigma\pi$ channels	Corrected No. of K^-p	Total $\bar{K}N$ channels	
$Y_0^*(1405)$	202±40	303±60	-	-	-
$Y_0^*(1520)$	238±45	358±68	150±30	300±60	1.2±0.3
$Y_1^*(1660)$	139±30	139±30	10±10	20±20	7.0±7.5
$Y_0^*(1815)$	30±30	45±45[a]	-	-	0.23±0.25[a]
$+ Y_1^*(1765)$		30±30[b]	96±20	192±40	0.31±0.33[b]

[a] Assuming all the resonance is in a $T = 0$ state.

[b] Assuming all the resonance is in a $T = 1$ state.

Table IV. Cross sections

	Hydrogen [Reactions (1) and (2)]				Deuterium [Reactions (3) and (4)]	
	1.51 BeV/c		1.70 BeV/c		1.49 BeV/c	
	Σ^+	Σ^-	Σ^+	Σ^-	Σ^+	Σ^-
Number of fitted events	676±27	544±23	201±14	149±12	265±16	308±18
Corrected No. of events	1130±57	745±45	320±24	194±17	485±42	505±52
σ (μb)	220±20	145±15	290±34	176±22	970±110	1020±120

FIGURE LEGENDS

Fig. 1. Mass plots of $\Sigma\pi$ systems in reactions 1 and 2 at 1.51 BeV/c incident K^- momentum (invariant mass of 2025 MeV). ——— Calculated distribution when 3/4 of the events exhibit a $Y_0^*(1405)$ and 1/4 a $Y_0^*(1520)$, —— — enhancements expected for $M(\Sigma^{\pm}\pi_1^-)$ from events with $Y_1^{*+}(1660)$ production, ----- enhancements expected for $M(\Sigma^{\pm}\pi_2^+)$ from events with $Y_1^{*+}(1660)$ production and decay via $Y_0^*(1405)$ (mode b), — — - - - enhancements expected for $M(\Sigma^{\pm}\pi_3^{\mp})$ from events with $Y_1^{*+}(1660)$ production and decay via $Y_0^*(1405)$ (mode b), — - - - enhancements expected for $M(\Sigma^{\pm}\pi_{2 \text{ or } 3})$ from events with $Y_1^{*+}(1660)$ production and decay via phase space (mode a), and — — — Total distribution for $M(\Sigma^+\pi^+)$ and $M(\Sigma^-\pi^-)$ where $Y_1^*(1660)$ production and decay via mode b are included.

Fig. 2. Mass plots for $\Sigma\pi\pi$ systems in reactions (1) and (2) at 1.51 BeV/c. The solid line indicates the distribution when 3/4 of the events exhibit $Y_0^*(1405)$, and 1/4 $Y_0^*(1520)$.

Fig. 3. Mass plots of $\Sigma\pi$ systems in reactions (1) and (2) at 1.70-BeV/c incident K^- momentum (invariant mass is 2110 MeV). ——— Distribution if 3/4 events exhibit $Y_0^*(1405)$ and 1/4 $Y_0^*(1520)$, — — enhancement expected in $M(\Sigma^+\pi^+)$ from the production of 20 $Y_1^{*+}(1660)$ and decay via mode b, and — . — enhancement expected in $M(\Sigma^-\pi^-)$ from the production of 20 $Y_1^{*+}(1660)$ and decay via mode b).

Fig. 4. Mass plots of $\Sigma\pi\pi$ systems in reactions (1) and (2) at 1.70-BeV/c incident K^- momentum. The solid line indicates the distribution as in Fig. 3, curve A.

Fig. 5. Dalitz plots of (a) $\Sigma^-\pi^-\pi^+$ events in reaction (3), and (b) $\Sigma^+\pi^-\pi^-$ events in reaction (4).

Fig. 6. Mass distributions of $\Sigma\pi$ and $\pi\pi$ systems in reactions (3) and (4). The curve is a rough fit explained in the text.


M(Σ⁺π⁺)
T₃=+2
676 points
▫ 1 points
M(Σ⁺π⁻)
T₃=0
676×2 points
▫ 1 point
M(Σ⁻π-)
T₃=-2
543 points
▫ 1 point
M(Σ⁻π⁺)
T₃=0
543×2 points
▫ 1 point
Number of events per 10-MeV interval
M (Σπ) (MeV)
M(Σπ) (MeV)
1320
1500
1700
MUB-2242

Fig. 1



Fig. 2

MUB-2241


M (Σ⁺π⁺)
T₃ = +2
201 points
□ 1 point
M (Σ⁺π⁻)
T₃ = 0
201×2 points
□ 1 point
M (Σ⁻π⁻)
T₃ = -2
149 points
□ 1 point
M (Σ⁻π⁺)
T₃ = 0
149×2 points
□ 1 point
Number of events per 10-MeV interval
0
10
20
30
1320
1400
1500
1600
1700
1800
M (Σ π) (MeV)
MUB-1755

Fig. 3



Fig. 4

MUB-1753



MU-33195

Fig. 5



Fig. 6

MUB-2243

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.