UCRL-11953 Rev.
c.2

University of California

Ernest O. Lawrence Radiation Laboratory

MEASUREMENTS OF TRANSITION PROBABILITIES OF ^2S-$^2P^o$ TRANSITIONS IN LITHIUM-LIKE IONS

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted for publication in the
Physics Letters Journal.

UCRL-11953 Rev.

UNIVERSITY OF CALIFORNIA
Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

MEASUREMENTS OF TRANSITION PROBABILITIES OF ^2S-$^2P^o$ TRANSITIONS IN LITHIUM-LIKE IONS

Klaus Berkner, William S. Cooper III,
Selig N. Kaplan, and Robert V. Pyle

March 26, 1965

MEASUREMENTS OF TRANSITION PROBABILITIES OF ^2S-$^2P^o$ TRANSITIONS IN LITHIUM-LIKE IONS*

Klaus Berkner, William S. Cooper III,
Selig N. Kaplan, and Robert V. Pyle

Lawrence Radiation Laboratory
University of California
Berkeley, California

March 26, 1965

We have used the foil-excitation technique recently employed by Kay[1] and Bashkin et al.[2] to measure directly the average of the lifetimes of the $1s^22p(^2P^o_{1/2})$ and $1s^22p(^2P^o_{3/2})$ states of C IV, N V, O VI, F VII, and Ne VIII ions in the lithium-like isoelectronic sequence. In each case a beam of ions accelerated to an energy of 0.97 MeV/nucleon by the Berkeley heavy-ion linear accelerator (Hilac) was further ionized and excited in an aluminum foil. The decay of radiation from the ^2S-$^2P^o$ transitions was then measured as a function of position downstream from the exciting foil; a vacuum-uv monochromator with a sodium-salicylate-covered photomultiplier was used as a photon detector.

Ions C^{+2}, N^{+3}, O^{+3}, F^{+3}, or Ne^{+3} were accelerated to $\beta = 0.0457$ by the "prestripper" section of the Berkeley Hilac. The ion beam was collimated, passed through a 140 $\mu g/cm^2$ Al foil, and monitored by two concentric Faraday cups. The foil was attached to a moveable shaft that permitted 35-cm displacement along the beam line. The central Faraday cup, 1.9-cm in diameter, received most of the beam (typically 80 to 90% for the greatest foil-cup separation). The other cup gave an indication of the angular spread of the beam.

The ions emerged from the foil with an equilibrium charge distribution and in various states of electronic excitation. The radiation from these

excited states was observed with a vacuum-uv scanning monochromator (McPherson 0.5-meter Seya-Namioka) with 1.0 by 20 mm entrance and exit slits. The monochromator collected light from 1 cm of beam path and had a roughly triangular bandpass 32 Å wide at half maximum. The detector was a photomultiplier tube (Amperex XP 1010) coated with sodium salicylate and cooled to -40°C to reduce thermal noise. Current pulses due to individual photons were registered by a scaler circuit; to correct for photomultiplier noise, data were also recorded on a second scaler for an equivalent time interval between beam pulses.

The monochromator was scanned in wavelength at a given foil position to locate the center of the doublet, and the photon counts were then recorded as a function of foil position. Peak counting rates were about 4000 counts per minute for a beam current of 1 μA. Figure 1 shows the number of counts as a function of foil position for the F VII doublet at 890.8 and 883.1 Å. The foil comes into the field of view of the monochromator at about 4.8 cm and leaves it at about 5.8 cm; beyond this position the intensity of the doublet is seen to decay exponentially with distance. Results for other ions in the lithium-like isoelectric sequence were similar.

The transition probabilities were obtained by least-squares fitting the data points with a computer program to an exponential plus a constant background. In every case a single exponential fitted the data very well, and the computer-selected background was consistent with estimates derived from the observed counting rate before the foil came into the field of view of the monochromator. Whether the data were normalized to the total charge collected by only the central Faraday cup or to the total charge collected by both Faraday cups made, on the average, a difference of only 2% in the value of the transition probability. In the final results these two values were

averaged and an uncertainty of 3% assigned, sufficient to include both the uncertainty in normalization and the uncertainty in curve fitting. The only other source of uncertainty is the calculation of the velocity from the accelerator parameters and the calculated energy loss in the foil ($\Delta E/E \approx 0.05$). An uncertainty of ± 1.5% is assigned to the calculated velocities.

The average transition probabilities for the $1s^2 2s(^2S_{1/2}) - 1s^2 2p(^2P^o_{1/2,3/2})$ transitions of C IV, N V, O VI, F VII, and Ne VIII are shown in Table I. Also shown are theoretical values calculated by Varsavsky[3] (screening approximation), Griem[4] (Coulomb approximation), Heroux[5] (Coulomb approximation), and Weiss[6] (Hartree-Fock method, a, and a presumably more accurate method, b, employing 45-term configuration interaction wave functions). The wavelengths and the various authors' estimates of the reliability of their values are included.

Within the estimated uncertainties, all the theoretical values, even those derived from the Coulomb approximation, agree well with our measurements.

We are grateful to Dr. C. M. Van Atta for supporting and encouraging this research. One of us (SNK) thanks Dr. A. C. Helmholz and Dr. B. J. Moyer for the support and interest that enabled him to participate. We also thank Dr. A. W. Trivelpiece and H. K. Forsen, University of California, Berkeley, for use of their monochromator.

FOOTNOTE AND REFERENCES

* Work performed under the auspices of the U. S. Atomic Energy Commission.

1. L. Kay, Proc. Phys. Soc. (London) 85, 163 (1965).
2. S. Bashkin, L. Heroux, and J. Shaw, Phys. Letters 13, 229 (1964).
3. C. M. Varsavsky, Astrophys. J. Suppl. Ser. No. 53, pp. 75-107 (1961).
4. H. R. Griem, Plasma Spectroscopy (McGraw-Hill Book Company, New York, 1964).
5. L. Heroux, Proc. Phys. Soc. (London) 83, 121 (1964).
6. A. W. Weiss, Astrophys. J. 138, 1262 (1963).

Table I. Average transition probabilities for $1s^2 2s(^2S_{1/2}) - 1s^2 2p(^2P^o_{1/2,3/2})$.

Spectrum	λ(Å)	Transition probability × 10^{-8} (sec^{-1})					
		Measured	Theoretical				
			Varsavsky	Griem	Weiss[a]	Weiss[b]	Heroux
C IV	1550.7 1548.2	2.97 ± 0.11	2.78	2.64	2.681	2.645	2.6
N V	1242.8 1238.8	3.34 ± 0.12	3.46	3.40	3.416	3.373	4.0
O VI	1037.6 1031.9	3.94 ± 0.14	4.37	4.14	4.144	4.055	4.2
F VII	890.8 883.1	4.48 ± 0.16	5.07	---	4.872	---	---
Ne VIII	780.3 770.4	5.55 ± 0.19	5.94	---	5.591	---	5.75
Estimated uncertainty		3.5 %	≥ 10 %	≥ 20%	5-15%	2-4%	

[a] Hartree-Fock method

[b] 45-term configuration interaction wave functions

FIGURE CAPTION

Fig. 1. Counts vs foil position for the F VII doublet at 890.8 and 883.1 Å. The line is the computer result for the best fit to the data.



MUB-5309

Fig. 1.