UCRL- 1209

UNIVERSITY OF CALIFORNIA — BERKELEY

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

RADIATION LABORATORY

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UCRL-1209
Unclassified Physics
Distribution

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

Elastic D-P Scattering at 190 Mev

Martin O. Stern and Arnold L. Bloom

April 5, 1951

Berkeley, California

UCRL-1209
Unclassified Physics
Distribution

Elastic D-P Scattering at 190 Mev

Martin O. Stern and Arnold L. Bloom

Radiation Laboratory, Department of Physics
University of California, Berkeley, California

April 5, 1951

As a result of recent theoretical interest, a comprehensive program has come into being, the purpose of which is to deduce some properties of the neutron-neutron interaction at high energies.

So far, p-p experiments have been carried out by Birge[1] at 100 Mev, and by Chamberlain, Segrè, and Wiegand[2] at, among other energies, 120 Mev. Professor Wilson M. Powell[3] is engaged in the study of n-d scattering using 90 Mev neutrons. We are measuring the scattering on protons, both elastic and inelastic, of the 190 Mev deuterons of the Berkeley 184-in. cyclotron. In all these experiments the relative velocities are comparable.

In this letter, we present our results to date on the elastic d-p scattering, first reported by one of us (ALB) at a recent meeting[4] of the American Physical Society. A more complete report, giving the details of the experiment, and possibly extending the data to larger and smaller angles, will be published in the near future.

The experimental arrangement was akin to that used by Chamberlain and Wiegand.[5] Deuterons scattered out of the cyclotron were collimated and made to impinge on a target. Coincidences between protons and deuterons were observed by means of stilbene crystals. Either one crystal, or two in

[1] R. W. Birge, Phys. Rev. 80, 490 (1950).

[2] Chamberlain, Segrè, and Wiegand, Phys. Rev. 81, 284 (1951).

[3] W. M. Powell, to be published.

[4] A. L. Bloom and M. O. Stern, Phys. Rev. 81, 660 A (1951).

[5] O. Chamberlain and C. Wiegand, Phys. Rev. 79, 81 (1950).

coincidence, detected the scattered protons, another crystal, in coincidence with the former, the deuterons. Suitable tests were performed to convince us that only the desired events were being observed.

Carbon and polyethylene targets were used alternately to obtain the scattering cross section by subtraction. The data accumulated so far are shown in Fig. 1. The differential scattering cross section, in 10^{-27} cm^2 sterad.$^{-1}$ in the center-of-mass system, is shown as a function of the angle of deflection of either particle in that system. Table I gives the weighted averages of the cross sections for a given angle. The slight relativistic correction has been taken into account.

The solid curve of Fig. 1 shows the theoretical cross section as given by Chew.[6] Evidently the fit is excellent from 60° to 110°, but at larger angles the measured cross section is lower than Chew's; this means that at this energy the pick-up process which causes a rise in the cross section in this region is less important than the Born approximation with Serber potentials for the n-p and p-p interaction would lead one to expect.

The errors shown in Fig. 1 are compounded of the standard deviations due to counting statistics and the estimated uncertainties in the factor which determines how much of the carbon count must be subtracted from that of the CH_2 target to yield the hydrogen effect. In addition, certain systematic errors, due to the uncertainties in the voltage plateaus, beam integration, and steadiness of the beam, etc., may occur. These have been estimated to amount to no more than 7 percent standard deviation, and have been taken into account in the errors quoted in Table I.

We wish to thank Drs. O. Chamberlain and E. Segrè for their constant assistance and interest in this work. This work was performed under the auspices of the Atomic Energy Commission.

[6] G. F. Chew, Phys. Rev. 74, 809 (1948).

Table I

Angle	Cross Section	Angle	Cross Section
38°	5.4 ± 0.4	98°	0.65 ± 0.05
48°	4.5 ± 0.4	108°	0.61 ± 0.05
58°	2.3 ± 0.2	118°	0.52 ± 0.09
68°	1.22 ± 0.09	128°	0.55 ± 0.05
78°	1.16 ± 0.10	138°	0.40 ± 0.05
82°	0.89 ± 0.08	149°	0.44 ± 0.09
88°	0.70 ± 0.06	159°	1.5 ± 0.5

Differential scattering cross sections in the center-of-mass, in 10^{-27} cm^2 sterad.$^{-1}$, have been averaged over all data for a given angle, to yield the values shown as a function of the angles of deflection in the center-of-mass.



Figure 1

Differential elastic scattering cross section of 190 Mev deuterons scattered from protons, in the center-of-mass system, as a function of the angle deflection in that system. The solid curve is the theoretical cross section as given by Chew (Ref. 6).