cy. 2

UCRL- 1242

UNCLASSIFIED

cy. 2

UNIVERSITY OF CALIFORNIA — BERKELEY

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

RADIATION LABORATORY

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UCRL-1242
Unclassified-Physics Distribution

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

SCATTERING OF π -MESONS BY DEUTERIUM

S. Fernbach, T. A. Green
and K. M. Watson

April 17, 1951

Berkeley, California

SCATTERING OF π -MESONS BY DEUTERIUM

S. Fernbach, T. A. Green
and K. M. Watson

Radiation Laboratory, Department of Physics
University of California, Berkeley, California

April 17, 1951

For sufficiently high meson energies the impulse approximation[1] may be used to express the scattering cross section for deuterium in terms of the free particle cross sections for protons and neutrons. The evaluation of such an expression is of interest since, under appropriate conditions, it may permit an estimate of the meson-neutron scattering cross section from the directly observable hydrogen and deuterium cross sections. The theoretical background for such a calculation has already been given by Chew[1], who has treated the very similar but more complicated problem of the scattering of neutrons by deuterium.

The starting point in the calculation is the use of the impulse approximation in which one assumes that the incident particle is scattered by a free nucleon wave packet whose momentum distribution is the same as that of one or the other of the bound nucleons. Then, following Chew, one argues that the contribution of each nucleon to the scattering amplitude is given to a good approximation by the product of the free particle scattering amplitude and an overlap integral which depends on the initial and final wave functions. One may simplify the calculation of the differential cross section by integrating it over the relative coordinates of the outgoing nucleons, thus considering only the scattering angle of the meson as the quantity to be observed. Then, according to the arguments of Gluckstern and Bethe[2], one can use plane waves

---

[1] G. F. Chew, Phys. Rev. 80, 196, 1950.
[2] Gluckstern and Bethe, Phys. Rev. 81, 761 (1951).

for the wave functions in the final state. The differential cross section for the scattering of a meson into the solid angle $d\Omega$, situated about the laboratory scattering angle θ, may then be written:

$$\frac{d\sigma}{d\Omega}(\theta) = \left[\frac{d\sigma_p}{d\Omega}(\theta) + \frac{d\sigma_n}{d\Omega}(\theta)\right] H_1(\theta)$$

$$+ 2\cos\omega\sqrt{\frac{d\sigma_p}{d\Omega}\frac{d\sigma_n}{d\Omega}} \quad H_2(\theta) \tag{1}$$

where $\frac{d\sigma_p}{d\Omega}(\theta)$ and $\frac{d\sigma_n}{d\Omega}(\theta)$ are the free particle cross sections, $\cos\omega$ is an unknown phase factor, and the weighting factors $H_1(\theta)$ and $H_2(\theta)$ are defined by:

$$H_1(\theta) = \frac{1}{J_o}\int d\vec{k}\, J(\vec{k})\, \phi_D^2\left[\vec{k} + \frac{1}{2}(\vec{q}(\vec{k}) - \vec{q}_o)\right]$$

$$H_2(\theta) = \frac{1}{J_o}\int d\vec{k}\, J(\vec{k})\, \phi_D\left[\vec{k} + \frac{1}{2}(\vec{q}(\vec{k}) - \vec{q}_o)\right]\phi_D\left[\vec{k} - \frac{1}{2}(\vec{q}(\vec{k}) - \vec{q}_o)\right] \tag{2}$$

In Eq. 2, $\vec{q}_o$ and $\vec{q}(\vec{k})$ are the momenta of the incident and outgoing mesons, respectively, and $\vec{k}$ is the relative momentum of the outgoing nucleons. ϕ_D is the Fourier transform of the deuteron wave function, $J(\vec{k})$ is the volume of momentum space available to the scattered meson, and J_o is the corresponding quantity for scattering by free particles. The integrations are to be extended over all values of $\vec{k}$ compatible with energy conservation.

Eq. 1 should best approximate the differential cross section for high meson energies and large scattering angles. In Fig. 1 we have plotted $H_1(\theta)$ and $H_2(\theta)$ for 70 Mev incident mesons and for scattering angles greater than 50 degrees.

One may obtain another approximate expression for the differential cross section, this one being best for small scattering angles, by neglecting the

restriction imposed on $\vec{q}(\vec{k})$ by energy conservation. If one replaced $\vec{q}(\vec{k})$ and $J(\vec{k})$ by appropriate average values and integrates over all $\vec{k}$ space there results the closure approximation for which

$$H_1^c(\theta) = 1 \qquad\qquad H_2^c(\theta) = \left\langle \frac{\sin |\vec{q} - \vec{q}_o| r}{|\vec{q} - \vec{q}_o| r} \right\rangle_{av.} \tag{3}$$

In Eq. 3, $\vec{q}$ is defined, for a given angle θ, by the free particle energy-momentum conservation laws, while the average is taken over the square of the deuteron wave function. As θ increases, $H_2^c(\theta)$ decreases from its initial value, unity. For purposes of comparison with the plane wave approximation, we have plotted $H_2^c(\theta)$ in Fig. 1. One sees that the two approximations give values of $H_1(\theta)$ and $H_2(\theta)$ which agree to within 15 percent. This indicates that the readily applied closure approximation may be used with some justification for the discussion of the cross section at other meson energies. On this basis we have calculated, for various incident meson energies, the scattering angle beyond which the correlation effects will be small. As a criterion of smallness we have taken $H_2^c(\theta) \quad \frac{1}{3} H_1^c(\theta)$. The results are shown in Fig. 2. They are of interest since they indicate under which experimental conditions one might estimate the meson-neutron cross section without a knowledge of the phase factor $\cos\omega$.

This work was performed under the auspices of the Atomic Energy Commission.

Information Division
5/21/51 nw

Caption

Fig. 1. Free particle and correlation term weighting factors for π-D scattering.

Fig. 2. Meson scattering angle beyond which correlation term is small.



FIG. 1 FREE PARTICLE AND CORRELATION TERM WEIGHTING FACTORS FOR π - D SCATTERING

MU 1934



FIG. 2 MESON SCATTERING ANGLE BEYOND WHICH CORRELATION TERM IS SMALL.

MU 1933