

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNCLASSIFIED

The Utilization of the Three Singly- C^{14} -marked Lactic Acids
by Escherichia coli.

Dean Fraser and Bert Tolbert

Virus Laboratory and the Radiation Laboratory of
the University of California, Berkeley, California

RECEIVED
AUG 8 1951
BERKELEY AREA
U. S. ATOMIC
ENERGY COMMISSION

As a preliminary to preparing bacteriophages marked with C^{14} , it was desirable to ascertain which singly-labelled lactic acid, when used as a carbon source, gives the highest incorporation of radioactive carbon in the cells of Escherichia coli. The results, while not unexpected, present several interesting features and serve to supplement the much more complete data obtained by Deudoroff (1951) with another bacterium under rather different conditions (see the accompanying article).

A synthetic medium containing sodium lactate as the energy source has been used widely in studies of the T-series of Escherichia coli bacteriophages (Adams, 1950). Since we have found that our usual glycerol medium gives better yields of T3 bacteriophage when fortified with casein hydrolysate, experiments on the utilization of lactates were conducted in the presence or absence of such a casein hydrolysate supplement, each set containing a control of the opposite type. The medium was prepared as usual (Adams, 1950).

The radioactive lactates were available as zinc salts. These were converted to the free acid by treatment with washed Dowex 50¹ ion exchange resin in the acid form. Each sample was dissolved in 1 ml water, mixed with 1 ml wet packed resin and the resin separated using a low speed centrifuge. The resin was washed with two half ml portions of water. The combined solutions in 1.5 ml water were made up to 50 ml with inactive lactate medium, the pH adjusted, and the solution autoclaved as usual. The specific radioactivity of each of the final mixtures was determined (Dauben et al: 1947).

Each culture was aerated with a simple bubbler in a test tube and the effluent CO_2 was collected in a scrubber of 1 M NaOH using an apparatus which passed the gas in a helical path. The bacteria were harvested at a concentration of 1 to 2 $\times 10^8$ cells/ml, washed by centrifugation twice

RECEIVED
AUG 8 1951
BERKELEY AREA
U.S. ATOMIC
ENERGY COMMISSION

with a 1.5 per cent phosphate buffer (pH 7.0) and once with distilled water. This suspension was plated directly for radioactivity (Dauben et al, 1947). The CO_2 in the scrubbers was precipitated as BaCO_3 on aluminum discs (Calvin et al, 1949). The radioactivity of the samples was determined either with a proportional counter (Nucleometer) or with thin end window GM tubes, depending on the specific activity. The final supernatant fluid from each sample was tested to show that no significant activity remained.

RESULTS

^{Expt. 1}
Table 1 shows the results for a set of cultures in which lactate was the sole carbon source except for sample 1, in which an amino acid supplement was added to a duplicate of sample 2. As expected, most of the utilized carbon from the ~~carboxyl-marked lactate~~ ^{lactate-1-C¹⁴} is discharged as CO_2 . Some, nonetheless, is retained as bacterial substance. With ~~1~~-marked ^{2-C¹⁴} lactate approximately three times as much of the carbon is retained in the bacteria and correspondingly less escapes as CO_2 ; with the ~~1~~-marked ^{3-C¹⁴} lactate the efficiency of utilization is slightly greater still.

^{Expt. 2}
Table 2 shows an experiment in which each of the three lactates was supplemented with casein hydrolysate (the culture with lactate-2 was paralleled by a similar unsupplemented culture). The results are similar to those shown in Table 1, except that in each case less of the radioactive carbon is used; i.e., the bacteria are supplying some of their requirements at the expense of the inactive casein hydrolysate.

^{In both experiments}
It is interesting to note that the total utilization of radioactive ^{labelled} carbon is greater for carboxyl-marked lactate than for the other isomers. Since the same total amount of lactate must be used in each case, this suggests ^{that} a portion of the C_2 fragment left after removal of the carboxyl is excreted into the medium ² (1.5 and 2.0 per cent excreted vs. 6.2 and

CHICAGO, ILL.
AUG 8 1951
U.S. DEPARTMENT OF COMMERCE
BUREAU OF ECONOMIC ANALYSIS

5.7 per cent utilized^{in Expt. 1}. The difference in total utilization with and without amino acid supplement shows that the bacteria draw on lactate as a carbon and energy source to a smaller extent when provided with amino acids. Since almost the same amount of carboxyl carbon is converted to CO_2 in the presence and absence of the supplement, it seems unlikely that the energy requirement of the bacteria is appreciably lowered by the provision of amino acids, the data suggesting rather that amino acids are used as a carbon and particularly as an energy source in preference to the two-carbon lactate fragment, which is apparently largely excreted (3.6 and 4 per cent vs. 2.5 to 2.1 per cent utilized^{in Expt. 2}). This is not surprising in view of the poor utilization of two carbon compounds as a sole carbon source by Escherichia coli (Stephenson, 1949).

It has been presumed that the carbohydrate of the Escherichia coli arises through reversal of the glycolytic cycle; i.e., through conversion of lactate to pyruvate, etc. The indication that lactate is used synthetically mostly as a two-carbon fragment rather than a three-carbon unit cannot, however, be interpreted as militating against this mechanism--as might appear at first consideration--because from the composition of Escherichia coli (Taylor, 1946) it can be seen that carbohydrate carbon is but some 10 per cent of the total carbon. The ratio of incorporation of carbon-1 to carbons 2 and 3 is thus still consistent with the hypothesis that carbohydrate is made from the intact three-carbon unit, and that the protein and nucleic acid constituents are synthesized from a smaller unit. The present results do not offer decisive evidence on this problem.

It may be of interest to note, incidentally, that the incorporation of the three lactates into T3 bacteriophage follows a parallel pattern.

RECEIVED
AUG 8 1951
BERKELEY AREA
U. S. ATOMIC
ENERGY COMMISSION

SUMMARY

Study of the utilization of the three C^{14} -singly-labelled lactic acids by Escherichia coli has shown that carboxyl carbon is largely, but not exclusively, converted to CO_2 . The alpha and beta carbons are mostly utilized in synthetic reactions but some is converted to CO_2 and an appreciable amount of the C_2 fragment left from decarboxylation of the lactate is excreted into the medium. Casein hydrolysate is apparently used synthetically and used as an energy source in preference to the C_2 fragment.

RECEIVED
AUG 8 1951
BERKELEY AREA
U. S. ATOMIC
ENERGY COM. D.

FOOTNOTES

¹ Don Chemical Co., Midland, Michigan.

² The sum of $\%C^{14}$ in CO_2 and $\%C^{14}$ in bacteria can differ for the three isomers only by C^{14} excreted into the medium. In either experiment subtraction of this total for parts b or c from part a thus gives un-used or excreted C^{14} .

RECEIVED
AUG 8 1951
BERKELEY AREA
U. S. ATOMIC
ENERGY COMMISSION

REGISTERED
AUG 8 1951
BERKELEY CALIF.
U. S. ATOMIC
ENERGY COMMISSION

TABLE 1

Utilization of the C¹⁴-singly-marked lactic acids by Escherichia coli

with and without casein hydrolysate supplement

Carbon Source	Original Medium		Disintegrations/ 10 ⁷ Ract./ml	% C ¹⁴ in Bacteria	% C ¹⁴ in CO ₂
	Disintegrations/ ml/min.	Ract./ml			
Expt. 1					
a. Lactate-1-C ¹⁴	5.63 x 10 ⁵	2.6 x 10 ⁹	2.4 x 10 ³	1.1	6.6
b. Lactate-2-C ¹⁴	7.58 x 10 ⁵	2.6 x 10 ⁹	10.4 x 10 ³	3.7	2.5
c. Lactate-3-C ¹⁴	4.80 x 10 ⁵	2.6 x 10 ⁹	8.0 x 10 ³	4.3	1.3
d. Lactate-2-C ¹⁴ plus casein hydrolysate	9.08 x 10 ⁵	2.6 x 10 ⁹	6.9 x 10 ³	1.9	0.50
Expt. 2					
a. Lactate-1-C ¹⁴ plus casein hydrolysate	5.68 x 10 ⁵	1.5 x 10 ⁹	2.4 x 10 ³	0.64	5.5
b. Lactate-2-C ¹⁴ plus casein hydrolysate	5.36 x 10 ⁵	1.8 x 10 ⁹	5.7 x 10 ³	1.9	0.58
c. Lactate-3-C ¹⁴ plus casein hydrolysate	4.36 x 10 ⁵	2.1 x 10 ⁹	3.9 x 10 ³	1.9	0.16
d. Lactate-2-C ¹⁴ unsupplemented	4.48 x 10 ⁵	1.5 x 10 ⁹	6.1 x 10 ³	2.1	1.3

REFERENCES

Adams, Mark, 1950, Methods in Medical Research, The Year Book Publishers, Inc., Chicago. Refer to II, 7.

Calvin et al, 1949, Isotopic Carbon, John Wiley and Sons, New York, 109-114.

Dauben, W. G., Ried, J. C., and Yankwich, P. S., 1947, Techniques in the use of carbon 14, Anal. Chem. 19, 828.

Stephenson, M., 1949, Bacterial Metabolism, Longmans, Green, London, 184.

Taylor, A. R., 1946, Chemical analysis of the T2 bacteriophage and its host, Escherichia coli (strain E), J. Biol. Chem., 165, 271.

Wiame, J. M., and Foudoroff, H., J. Bact. (accompanying paper).

1951 Oxidative assimilation by Pseudomonas saccharophila with C¹⁴-labelled substrates J. Bact. 62.

RECEIVED
AUG 8 1951
BERKELEY AREA
U. S. ATOMIC
ENERGY COMMISSION