UCRL-16850

University of California
Ernest O. Lawrence Radiation Laboratory

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

THE MASSES AND EXCITED STATES OF ^{24}Al AND ^{28}P

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to Physics Letters

UCRL-16850

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

THE MASSES AND EXCITED STATES OF ^{24}Al AND ^{28}P

N. Mangelson, M. Reed, C. C. Lu, and F. Ajzenberg-Selove

May 1966

THE MASSES AND EXCITED STATES OF ^{24}Al AND ^{28}P[†]

N. Mangelson, M. Reed, C. C. Lu, and F. Ajzenberg-Selove[††]

Lawrence Radiation Laboratory
University of California
Berkeley, California

May 1966

The ^{24}Mg(^{3}He,t)^{24}Al and ^{28}Si(^{3}He,t)^{28}P reactions have been studied at ^{3}He energies in the range 37.8 to 40.1 MeV. The mass excesses of ^{24}Al and ^{28}P are, respectively, -0.07±0.06 and -7.12±0.06 MeV; 18 excited states of ^{24}Al and 12 excited states of ^{28}P have been observed.

[†]This work was supported by the U. S. Atomic Energy Commission and by the National Science Foundation.

[††]J. Simon Guggenheim Fellow 1965-1966; now at Haverford College, Haverford, Pennsylvania.

There is very little known[1),†] about the proton-rich nuclei, $^{24}_{13}Al$ and $^{28}_{15}P$. The positron decay has been observed[1-3)] to various of the states in the corresponding $T_z = 0$ nuclei, $^{24}_{12}Mg$ and $^{28}_{14}Si$. The atomic mass excesses of ^{24}Al ^{28}P are [4),††] 0.09±0.3 MeV and -7.69±0.3 MeV. If the ^{24}Al $E_{\beta+}$ (max) reported by Scott and Polichar[3)] are assumed to be due to transitions to the ^{24}Mg states at 4.12 MeV[1)] and 8.44 MeV[4,5)], the atomic mass excess of ^{24}Al is -0.01±0.13 MeV. No energy levels of ^{24}Al and ^{28}P have been reported. By analogy with the mirror nucleus ^{24}Na, the ground state of ^{24}Al should have $J^{\pi} = 4+$. The corresponding $T = 1$ state in ^{24}Mg has been located[3)] at $E_x = 9.512 \pm 0.008$ MeV. The ground state of ^{28}P is possibly a $J^{\pi} = 3+$ state, by analogy with the 3+, first $T = 1$ state[1)] in ^{28}Si at $E_x = 9.324 \pm 0.011$ MeV[3)], although the close spacing of the next $T = 1$ state ($\Delta E_x = 31$ keV in ^{28}Al, between the $J^{\pi} = 3+$ and 2+; $E_x \approx 60$ keV in ^{28}Si, between the $J^{\pi} = 3+$ and 2+) would permit $J^{\pi} = 2+$ as an alternative.

The two types of reaction which can be used to study ^{24}Al and ^{28}P are the (p,n) and the (^3He,t) reactions on ^{24}Mg and ^{28}Si with Q values of -14 to -15 MeV. We will report on a study of the $^{24}Mg(^3He,t)^{24}Al$ and $^{28}Si(^3He,t)^{28}P$ reactions[†††] at $E(^3He) \approx 40$ MeV.

---

†C. Van der Leun, private communication. We are indebted to Dr. Van der Leun for information concerning more recent evidence on ^{24}Al and ^{28}P.

††These mass excesses have been calculated on the basis of the ^{24}Mg and ^{28}Si masses given by ref. 4, and the Q of the ^{24}Al and ^{28}P $\beta+$ decay given by ref. 1.

†††These reactions have been observed, but no results have been published: See M. Rickey and R. G. Matlock, Bull. Am. Phys. Soc. 10 (1965) 463.

A beam of 40 MeV ^{3}He particles, accelerated by the Berkeley 88-inch spiral ridge cyclotron, was used for the experiments. The experimental procedures were identical to those described in another paper[6]. The ^{24}Mg and Si targets† were self-supporting foils. The magnesium target was enriched to 99.96% ^{24}Mg and its thickness was 270±30 μg/cm^2. The silicon was not enriched and therefore contained 92.2% ^{28}Si: the target thickness was 160±15 μg/cm^2. ^{12}C foils used for calibration purposes ranged from 150 to 300 μg/cm^2.

Figure 1 shows parts of the triton spectra for the ^{24}Mg(^{3}He,t)^{24}Al reaction at E(^{3}He) = 39.3 MeV, θ = 20° and 25°. Additional spectra were also obtained at E(^{3}He) = 39.3 MeV (θ = 30°, 50°, 55°, and 60°) and at E(^{3}He) = 40.1 MeV (θ = 30°). The FWHM of peaks corresponding to single states is ≈ 85 keV. The Q value for the ground-state reaction was determined to be -13.88±0.06 MeV. From this and the masses given by Mattauch et al.[4] for ^{24}Mg, ^{3}He, and t, the mass excess of ^{24}Al is determined to be -0.07±0.06 MeV. The excitation energies of the observed states of ^{24}Al are displayed in table 1. The first excited state we observe in ^{24}Al at E_x = 0.47±0.03 MeV is in good agreement with the energy of the known[1] first state in the mirror nucleus ^{24}Na, E_x = 0.473±0.003 MeV. The higher states cannot be meaningfully compared. It is clear that some of our triton groups correspond to unresolved levels. The differential cross section for formation of the ground state of ^{24}Al decreases from 67±15 μb/sr at 23.4° c.m. to 6±2 μb/sr at 68.4° c.m.

---

†The targets were prepared by C. E. Ellsworth and A. Johns. We acknowledge with thanks their invaluable help. The separated ^{24}Mg was furnished by the Stable Isotopes Division of ORNL.

Figure 2 shows parts of the triton spectra for the $^{28}Si(^{3}He,t)^{28}P$ reaction at $E(^{3}He) = 37.8$ MeV, $\theta = 20^{\circ}$ and 35°. Additional spectra were obtained at $E(^{3}He) = 40.1$ MeV, $\theta = 30^{\circ}$ and 35°. The Q value for the ground state reaction was determined to be -14.38±0.06 MeV. From this the mass excess of ^{28}P was determined to be -7.12±0.06 MeV. The excitation energies of the observed ^{28}P states are shown in table 1. The ground state and first excited states of ^{28}P were resolved by fitting the triton peaks with two Gaussian peaks. It is clear here also, by comparison with ^{28}Al, that many of the triton groups are due to unresolved states. The differential cross section for the ground state of ^{28}P falls from 70±15 μb/sr at 22.8°c.m. to 25±6 μb/sr at 39.7°c.m.

By use of the method of Wilkinson[7] and the 1964 mass tables[4], the excitation energy of the first $T = 1$, $T_z = 0$ level was predicted for the $A = 24$ and $A = 28$ isobaric triplets. The first $T = 1$ level of ^{24}Mg is 9.512±0.008 MeV[3], compared with predicted values of 9.502±0.008 and 9.54±0.07 MeV based on the masses of ^{24}Na and ^{24}Al respectively. The first $T = 1$ level of ^{28}Si is 9.324±0.011 MeV[3] compared with predicted values of 9.378±0.008 and 9.35±0.07 MeV based on the masses of ^{28}Al and ^{28}P respectively.

We are grateful to the entire staff of the 88-inch cyclotron, and in particular to Joel Moss, Creve Maples, Jr., Donald A. Landis, and Fred S. Goulding. We are greatly indebted to Dr. Bernard G. Harvey and Dr. J. Cerny for their interest and many useful comments.

References

1) P. M. Endt and C. Van der Leun, Nucl. Phys. 34 (1962) 1.

2) A. Scott and R. Polichar, Bull. Am. Phys. Soc. 8 (1963) 85.

3) M. Rickey, E. Kashy, and D. Knudsen, Bull. Am. Phys. Soc. 10 (1965) 550.

4) J. H. E. Mattauch, W. Thiele, and A. H. Wapstra, Nucl. Phys. 67 (1965) 1.

5) E. L. Haase, H. A. Hill and D. B. Knudsen, Phys. Letters 4 (1963) 338.

6) N. Mangelson, F. Ajzenberg-Selove, M. Reed, and C. C. Lu, Nucl. Phys. (1966).

7) D. H. Wilkinson, Phil. Mag. 1 (1956) 1031.

Table 1

Energy levels of ^{24}Al and ^{28}P

Levels of ^{24}Al				Levels of ^{28}P			
Group no.	E_x [a]	Group no.	E_x [a]	Group no.	E_x [a]	Group no.	E_x [a]
1	0	11	3.92±30	1	0	11	4.17±70
2	0.47±30	12	4.34±40	2	0.125±30	12	4.53±50
3	1.12±30	13	4.53±40	3	1.14±40	13	4.94±30
4	1.28±50	14	4.77±50	4	1.31±30		
5	1.62±40	15	4.48±60	5	1.54±40		
6	2.38±40	16	5.93±40	6	2.12±30		
7	2.88±30	17	6.81±40	7	2.65±100		
8	3.06±40	18	7.07±50	8	3.24±30		
9	3.35±50	19	(8.25±50) [b]	9	3.59±30		
10	3.71±50			10	3.84±30		

[a] Energies given in MeV ± keV.

[b] Assignment to ^{24}Al not certain.

Figure Captions

Fig. 1. Spectra from the $^{24}Mg(^{3}He,t)^{24}Al$ reaction at $E(^{3}He) = 39.3$ MeV, $\theta = 20^{\circ}$ and 25°. N represents the number of counts per channel. The E_x scale gives the excitation energy in ^{24}Al. The binding energy ($E_b = 1.88$ MeV) for breakup into $^{23}Mg + p$ is indicated. The locations of possible contaminant groups from $^{12}C(^{3}He,t)^{12}N$ and $^{16}O(^{3}He,t)^{16}F$ are also shown.

Fig. 2. Spectra from the $^{28}Si(^{3}He,t)^{28}P$ reaction at $E(^{3}He) = 37.8$ MeV ($\theta = 20^{\circ}$ and 35°). The binding energy (2.03 MeV) for breakup into $^{27}Si+p$ is shown. See also caption for fig. 1.



MUB-10173

Fig. 1



MUB-10172

Fig. 2

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.