UCRL-16980

University of California

Ernest O. Lawrence Radiation Laboratory

DECAY PROPERTIES OF THE $A_2(1310)$ MESON

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to Physical Review Letters

UCRL-16980
Preprint

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

DECAY PROPERTIES OF THE $A_2(1310)$ MESON

Suh Urk Chung, Orin I. Dahl, Lyndon M. Hardy,
Richard I. Hess, Janos Kirz, and Donald H. Miller

November 28, 1966

Decay Properties of the $A_2(1310)$ Meson*

Suh Urk Chung,† Orin I. Dahl, Lyndon M. Hardy,‡
Richard I. Hess,** Janos Kirz, and Donald H. Miller

Lawrence Radiation Laboratory and Department of Physics
University of California, Berkeley, California

November 28, 1966

ABSTRACT

The properties of the A_2 enhancement are determined from the $K\bar{K}$ and $\pi\rho$ decay modes independently. The characteristics of both systems are consistent with the decay of a particle having $I^G J^P = 1^- 2^+$.

Existence of a strong $\pi\rho$ enhancement between 1.0 and 1.4 BeV was discovered by Goldhaber et al. in a study of the reaction $\pi^+p \rightarrow \pi^+\pi^+\pi^-p$ at 3.65 BeV/c.[1] The ABBBHLM Collaboration[2] and Chung et al.[3] demonstrated that the enhancement consisted of two peaks: the A_1 at 1080 MeV and the A_2 at 1310 MeV. In addition, Chung et al. reported evidence for existence of a $K\bar{K}$ peak at 1310 MeV; the assignment $I^GJ^P = 1^-2^+$ was deduced on the assumption that the $\pi\rho$ and $K\bar{K}$ peaks represented alternative decay modes of the $A_2(1310)$. In several recent studies of the $\pi\rho$ system alone, the assignment $J^P = 2^+$ has been favored for the $A_2(1310)$.[4-6] In others, however, the assignments $J^P = 1^+$ or 2^- have appeared more likely;[7-9] in this case, the $K\bar{K}$ peak represents the decay of a new particle. In the present Letter we attempt to resolve this question by determining quantum numbers independently for the $K\bar{K}$ and $\pi\rho$ peaks; the analysis supports the original assumption of Chung et al.[3]

The film was obtained in the course of a systematic study of π^-p interactions near 3.2 and 4.2 BeV/c in the Lawrence Radiation Laboratory's 72-inch hydrogen bubble chamber. The experimental details are given by Hess[10] and Chung.[11] The observed numbers of events and corresponding cross sections are given in Table I.

The $K\bar{K}$ system has been studied in both $pK^-K_1^0$ and $nK_1^0K_1^0$ final states where the decays $K_1^0 \rightarrow \pi^+\pi^-$ were observed; all successfully fitted events in the fiducial volume were used. In contrast, the $A_2(1310)$ represents less than 10% of the $\pi^+\pi^-\pi^-p$ final state, so that useful comparisons are possible only after imposition of stringent selection criteria. For subsequent analysis, $\pi^-\rho^0p$ events are defined as those

with at least one $M(\pi^+\pi^-)$ combination in the interval 0.66 to 0.84 BeV. Background due to the sequence $\pi^- p \to \pi^-\pi^- N^{*++}(1238) \to \pi^-\pi^-\pi^+ p$ has been minimized by rejecting events with $1.12 \text{ BeV} \leq M(\pi^+ p) \leq 1.32$ BeV and $\Delta^2_{p\pi^+} \leq 1.5 \text{ (BeV/c)}^2$. In addition, events were rejected if they fell into the region where the Deck mechanism[12] is strongest, $\Delta^2_{p\pi^-} \leq 0.55 \text{ (BeV/c)}^2$ and $\hat{p}_p \cdot \hat{p}_0 \equiv \cos\theta_p \leq -0.8$. Here $\underset{\sim}{p}_p$ is the momentum of the outgoing proton and $\underset{\sim}{p}_0$ is the beam direction in the $p\pi^-$ rest frame.[13]

The effective-mass distributions, $M(K\bar{K})$, for the $K\bar{K}$ systems are shown in Fig. 1a; the $M(\pi^-\rho^0)$ distribution is shown in Fig. 1d for events with $\Delta^2_p \leq 0.65 \text{ (BeV/c)}^2$. In both cases a good fit is provided by a Breit-Wigner resonance, with $M_0 = 1310 \pm 20$ MeV and $\Gamma = 65 \pm 20$ MeV, above a smooth background. The Δ^2_N distributions for events in the A_2 interval, 1.24 to 1.38 BeV, are shown separately for $K\bar{K}$ events (Fig. 1, b and c) and $\pi^-\rho^0$ events (Fig. 1, e and f). After comparison with control regions we conclude that, within statistics, contributions from the 1310-MeV peak are similar in all cases.

Possible quantum numbers for the $K\bar{K}$ peak at 1310 MeV are readily deduced. Since the decay $K_1^0K_1^0$ is observed, C is +1 and J^P is $(\text{even})^+$. Histograms of decay cosine ($\cos\theta_K$ in the $K\bar{K}$ rest frame) and Treiman-Yang angle are plotted in Fig. 2. The decay cosine distributions for both the $K^-K_1^0$ and $K_1^0K_1^0$ events contain strong $\cos^2\theta$ components, so that J is not equal to zero. Since I is 1 for $K^-K_1^0$, we conclude that $G = (-1)^{J+I} = -1$. Consequently, I^G is 1^- and J^P is 2^+, 4^+, etc. for the $K\bar{K}$ peak at 1310 MeV.

In deducing possible quantum numbers for the $\pi\rho$ system, we note first that the decay $A_2(1310) \rightarrow \pi\rho$ is allowed, consequently G is -1; in addition, Abolins et al. have shown that I = 1.[14] To determine the spin and parity, we consider the decay correlations in the A_2 rest frame; we define $\underset{\sim}{q}$ as the relative momentum of the $\pi^+\pi^-$ pair forming the ρ^0, $\underset{\sim}{p}$ as the momentum of the third pion, and $\cos\beta \equiv \hat{q} \cdot \hat{p}$. For collinear decays, corresponding to points on the boundary of the Dalitz plot $\cos\beta = \pm 1$. For these decays $\psi(3\pi)$ is proportional to $Y_J^M(\hat{q})$, so that P is $-(-1)^J$; consequently, for 3π systems with $P = (-1)^J$, collinear decays are <u>not allowed</u>.[15] Since the parity of the 3π system can be deduced <u>only</u> from the density on the Dalitz plot near the boundaries, a precise estimate of background is crucial; for systems with $P = (-1)^J$, a small residual background of collinear events can lead (erroneously) to the <u>opposite</u> parity assignment.

The $\cos\beta$ distributions are shown in Fig. 3, b, c, and d for events in the A_2 region and control regions; the strong contribution from the decay $A_2 \rightarrow \pi\rho$ produces the peak at $\cos\beta \simeq 0.2$ in Fig. 3b. Events in the interval 1.24 to 1.38 BeV may be identified as (1) $A_2 \rightarrow \pi\rho$, (2) $\pi\rho$ background, or (3) 3π background. We designate the fraction of events of each type by ϵ_i. The smooth curve in Fig. 1d suggests that $\epsilon_2 + \epsilon_3 = 0.6 \pm 0.1$; the $M(\pi^+\pi^-)$ distribution for the same events gives $\epsilon_3 = 0.4 \pm 0.1$.

For comparison with the experimental data, theoretical $\cos\beta$ distributions[16] for possible J^P assignments were modified by addition of noninterfering background. To examine the dependence of each J^P assignment on background, $\epsilon_2 + \epsilon_3$ was varied from 0 to 1. The $\pi\rho$ background was

calculated using the matrix element for an s-wave $\pi\rho$ interaction (i.e., $J^P = 1^+$ appropriately symmetrized);[17] a uniform distribution in $\cos\beta$ was assumed for the 3π background. The behavior of χ^2 (for 19 degrees of freedom) is shown in Fig. 3a as a function of the assumed background level; the slopes are discontinuous, since we have arbitrarily set $\epsilon_2 = 0$ for $\epsilon_1 \geq 0.6$. We note that when background is ignored, the most likely assignments are $J^P = 1^+(\ell = 0)$ and $J^P = 2^-(\ell = 1)$. However, for a realistic background level of 40 to 70%, $J^P = 2^+$ represents the only assignment (of those considered) compatible with the data: the fitted distribution is shown in Fig. 3b for $\epsilon_1 = 0.4$. Consequently, for a model with noninterfering background, parsimony requires that we identify the $\pi\rho$ and $K\bar{K}$ peaks as alternative decay modes of an $I^G J^P = 1^- 2^+$ state at 1310 MeV; production cross sections are given in Table II. The combined data give $\Gamma(A_2^- \to K\bar{K})/\Gamma(A_2^- \to \pi\rho) = 0.05 \pm 0.02$; a factor of 1/2 has been included for the unobserved $\pi^0\rho^-$ decays. The decay $A_2 \to \pi\eta$ is allowed; some evidence for a peak near 1310 MeV has been reported. We have examined the $M(\pi^-\eta)$ distribution from the $\pi^-(\pi^+\pi^-\pi^0)p$ final state and (after correcting for unobserved decays) estimate $[\Gamma(A_2^- \to \pi\eta)]/[\Gamma(A_2^- \to \pi\rho)] = 0.12 \pm 0.08$.

Since A_2 events are concentrated at low Δ_N^2 and the decay $A_2 \to \pi\rho$ is dominant, it is likely that production occurs through ρ exchange. Unmodified, this model predicts a $\cos^2\theta_K \sin^2\theta_K(1 + a\cos 2\phi)$ distribution for the $K\bar{K}$ decay mode, where ϕ is the Treiman-Yang angle; the $\cos\theta_K$ distributions in Fig. 2 are in strong disagreement with this prediction. Similarly, the model does not account for the observed correlation between the beam direction and the normal to the $\pi\rho$ decay

plane.[11] Analogous discrepancies in other reactions involving ρ exchange have been explained by absorption effects.[18]

We wish to thank the scanning and measuring staff for their untiring efforts in bringing this experiment to a successful completion. It is a pleasure to acknowledge the encouragement and support of Professor Luis Alvarez throughout the course of this experiment.

FOOTNOTES AND REFERENCES

*Work done under the auspices of the U. S. Atomic Energy Commission.

†Present address: Brookhaven National Laboratory, Upton, L. I., N. Y.

‡Present address: TRW Systems, Inc., 1 Space Park, Redondo Beach, California.

**Present address: Logicon, Inc., 205 Avenue I, Redondo Beach, Calif.

1. G. Goldhaber, J. L. Brown, S. Goldhaber, J. A. Kadyk, B. C. Shen, and G. H. Trilling, Phys. Rev. Letters 12, 336 (1964).

2. Aachen-Berlin-Birmingham-Bonn-Hamburg-London (I. C.)-München Collaboration, Phys. Letters 10, 226 (1964).

3. S. U. Chung, O. I. Dahl, L. M. Hardy, R. I. Hess, G. R. Kalbfleisch, J. Kirz, D. H. Miller, and G. A. Smith, Phys. Rev. Letters 12, 621 (1964).

4. J. Alitti, J. P. Baton, B. Deler, M. Neveu-René, J. Crussard, J. Ginestet, A. H. Tran, R. Gessaroli, and A. Romano, Phys. Letters 15, 69 (1965).

5. R. L. Lander, Maris Abolins, D. D. Carmony, T. Hendricks, Nguyen-huu Xuong, and P. M. Yager, Phys. Rev. Letters 13, 346 (1964) and Ref. 2 report that the $J^P = 2^+$ assignment is consistent with their data, although other J^P assignments cannot be ruled out.

6. G. Benson, L. Lovell, E. Marquit, B. Roe, D. Sinclair, and J. Vander Velde, Phys. Rev. Letters 16, 1177 (1966).

7. V. E. Barnes, W. B. Fowler, K. W. Lai, S. Orenstein, D. Radojicic, M. S. Webster, A. H. Bachman, P. Baumel, and R. M. Lea, Phys. Rev. Letters 16, 41 (1966).

8. N. M. Cason, Phys. Rev. 148, 1282 (1966).

9. Aachen-Berlin-CERN Collaboration, reported by D. R. O. Morrison at the XIIIth International Conference on High Energy Physics, Berkeley, 1966. See G. Goldhaber, rapporteur's talk in the proceedings of this conference (University of California Press, Berkeley to be published).

10. R. I. Hess, Production of $K\bar{K}$ Pairs in $\pi^- p$ Interactions (Ph. D. thesis), UCRL-16832, June 1966.

11. S. U. Chung, Multi-pion Final States from $\pi^- p$ Interactions at 3.2 and 4.2 GeV/c (Ph. D. thesis), UCRL-16981, July 1966.

12. R. T. Deck, Phys. Rev. Letters 13, 169 (1964).

13. This cut reduces the number of events in the mass region below the A_2, but eliminates few events in the A_2 region itself. The analysis was also carried out without this cut, and the conclusions remained unchanged. See Ref. 11 for a detailed account.

14. M. A. Abolins, D. D. Carmony, R. L. Lander, O. Piccioni, Ng. H. Xuong, and P. M. Yager, Phys. Letters 21, 584 (1966).

15. R. H. Dalitz, Strange Particles and Strong Interactions (Oxford University Press, London, 1962), p. 40.

16. We have used a computer program written by R. Diebold, CERN/TC/PROG 64-25 modified for our purpose. As an alternative method for taking background into account, varying amounts of the $\cos\beta$ distribution in the control region were subtracted from the distribution in the A_2 interval. A comparison of the resulting distribution with theoretical curves for various J^P assignments yielded similar results; see Ref. 11 for details.

17. We thank Dr. Vanya Cocconi (private communication) for informing us that calculated distributions for events produced through the Deck mechanism resemble closely those for a $J^P = 1^+$ system.

18. J. D. Jackson, J. T. Donohue, K. Gottfried, R. Keyser, and B. E. Y. Svensson, Phys. Rev. 139, B428 (1965).

Table I. Final states analyzed.

Final state	Number of events		Cross section (μb)	
	3.2 BeV/c	4.2 BeV/c	3.2 BeV/c	4.2 BeV/c
$p\pi^+\pi^-\pi^-$	6318	2986	1910 ± 80	1920 ± 100
pK^0K^-	228[a]	95[a]	65.1±5.3	65.7±7.9
$nK_1^0K_\ell^0$	201[b]	68[b]	45.3±4.1	36.6±5.1

a. $K_1^0 \rightarrow \pi^+\pi^-$ decay was observed for these events. The cross sections were corrected for this detection efficiency ($\epsilon \approx 1/3$).

b. Decay of both $K_1^0 \rightarrow \pi^+\pi^-$ was observed for these events. The cross sections were corrected for this detection efficiency ($\epsilon \approx 4/9$).

Table II. Cross section for A_2 production.

Reaction	Cross section (μb)	
	3.2 BeV/c	4.2 BeV/c
$\pi^- p \rightarrow A_2^- p;\ A_2^- \rightarrow K^0 K^-$	18 ± 4	17 ± 5
$\pi^- p \rightarrow A_2^0 n;\ A_2^0 \rightarrow K\bar{K}$	36 ± 10	18 ± 9
$\pi^- p \rightarrow A_2^- p;\ A_2^- \rightarrow \rho^0 \pi^-$	150 ± 50	175 ± 45

FIGURE LEGENDS

Fig. 1. (a) Effective-mass histogram for the $K\bar{K}$ systems at 3.2 and 4.2 BeV/c. (b) and (c) Histograms of Δ_N^2 at 3.2 and 4.2 BeV/c for $K\bar{K}$ events in the A_2 region. (d) Effective-mass histogram for the $\pi^- \rho^0$ system at 3.2 and 4.2 BeV/c. Selections are discussed in the text. (e) and (f) Histograms of Δ_p^2 at 3.2 and 4.2 BeV/c for the $\pi\rho$ events in the A_2 region.

Fig. 2. Histograms of decay cosine (= $\hat{p}_K \cdot \hat{p}_0$ in the A_2 rest frame) and the Treiman-Yang angle for $K\bar{K}$ events in the A_2 region. The A_2^- histogram is shown in (a) and (b) and the A_2^0 in (c) and (d). In (c) and (d) two points have been plotted per event.

Fig. 3. (a) Variations of χ^2(19 degrees of freedom) for various J^P assignments for the A_2 as a function of the background level. (b) The $\cos\beta$ distribution in the A_2 region. See text for explanation of the curves. (c) The $\cos\beta$ distribution for the region below the A_2(1.0 to 1.14 BeV). (d) The same distribution for the region above the A_2(1.48 to 1.62 BeV).


(a)
K_1^0 K_1^0 (265 events)
K_1^0 K^- (317 events)
Number of events
M (KK̄) (BeV)
(b)
K_1^0 K_1^0
K_1^0 K^-
3.2 BeV/c
(c)
4.2 BeV/c
Δ_N^2 (BeV/c)^2
(d)
1461 events
70%
40%
Number of events
M ($\pi^-\rho^0$) (BeV)
(e)
3.2 BeV/c
(f)
4.2 BeV/c
Δ_p^2 (BeV/c)^2
MUB-12320

Fig. 1


(a) A_2^- (105.38 events)
2.9-3.3 BeV/c
3.8-4.2 BeV/c
(b) A_2^-
(c) A_2^0 (104.12 events)
2.9-3.3 BeV/c
3.8-4.2 BeV/c
(d) A_2^0
Number of events
Decay cosine
Treiman-Yang angle (deg)

Fig. 2


(a)
χ^2
Background (%)
$2^-(\ell=3)$
2^+
$1^+(\ell=2)$
1^-
$1^+(\ell=0)$
$2^-(\ell=1)$
(b)
479 events
Number of events
Cos β
(c)
224 events
(d)
157 events
MUB 12319

Fig. 3

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.