

University of California
Ernest O. Lawrence
Radiation Laboratory

GAMMA-VIBRATIONAL BANDS IN ^{182}Os AND ^{184}Os

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Int. Conf. on Nucl. Phys.
Proceedings

UCRL-16986

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

GAMMA-VIBRATIONAL BANDS IN ^{182}Os AND ^{184}Os

T. Yamazaki and D. L. Hendrie

July 1966

GAMMA-VIBRATIONAL BANDS IN ^{182}Os AND ^{184}Os †

T. Yamazaki and D. L. Hendrie

Lawrence Radiation Laboratory
University of California
Berkeley, California

July 1966

ABSTRACT

The systematics of the gamma-vibrational bands in even Os nuclei ($A = 182 \sim 192$) has been obtained from the study of gamma rays in $^{185,187}\text{Re}(p,2n)^{184,186}\text{Os}$ and $^{185,187}\text{Re}(p,4n)^{182,184}\text{Os}$ reactions. The gamma-vibrational energies obtained are 892 keV in ^{182}Os and 942 keV in ^{184}Os . The present results reveal a unique correlation of $\hbar\omega_\gamma$ with the moment of inertia.

†Work performed under the auspices of the U. S. Atomic Energy Commission.

The gamma-vibrational bands in neutron-deficient even Os nuclei have been studied by observing interband gamma rays in $^{185,187}\text{Re}(p,2n)^{184,186}\text{Os}$ and $^{185,187}\text{Re}(p,4n)^{182,184}\text{Os}$ reactions. The previously known gamma-vibrational band of ^{186}Os , which was studied by Emery et al.¹ from the decay of ^{186}Ir , is considerably populated up to the 5^+ level in $^{187}\text{Re}(p,2n)^{186}\text{Os}$ reaction (see the spectrum in Fig. 1(a)). This information was utilized to assign gamma rays in $^{185}\text{Re}(p,2n)^{184}\text{Os}$ reaction (Fig. 1(b)). The assignment given in Fig. 1(b) seems to be suspicious, because in the strong-coupling limit the $2_{\gamma} \rightarrow 2_g$ line and $2_{\gamma} \rightarrow 0_g$ lines should agree with the $4_{\gamma} \rightarrow 4_g$ and $3_{\gamma} \rightarrow 2_g$ lines, respectively. If we regard the 840 keV peak as the $2_{\gamma} \rightarrow 2_g$, $4_{\gamma} \rightarrow 4_g$ doublet and the 936-keV peak as the $2_{\gamma} \rightarrow 0_g$, $3_{\gamma} \rightarrow 2_g$ doublet, then the 822 keV and 942 keV lines must be associated with another level at 942 keV. However, if they were so, such a level should also be populated in ^{186}Os as well as the gamma vibrational levels. Therefore we have made the assignments as shown in the Fig. 1(b), although they are still doubtful.

After having established the gamma-vibrational band in ^{184}Os , the population of the same states was examined in $^{187}\text{Re}(p,4n)^{184}\text{Os}$ reaction at $E_p = 35$ MeV, which showed lower population of the gamma-vibrational band compared with the ground band and relatively higher population of higher-spin gamma-vibrational states. Subsequently this information was utilized to identify gamma rays in $^{185}\text{Re}(p,4n)^{182}\text{Os}$ reaction.

The systematics of the ground and the gamma-vibrational bands has been obtained, as shown in Fig. 2. Remarkable facts are

1) The energy of gamma vibration is uniquely correlated with the moment of inertia. As the neutron number decreases, the moment of inertia

and the gamma-vibrational energy increase until they reach maxima at $A = 184$ ($N = 108$), then both start decreasing.

2) There is appreciable difference in the moment of inertia between the ground band and the gamma-vibrational band. The level spacing $\Delta_{32}^{\gamma} \equiv E(3_{\gamma}) - E(2_{\gamma})$ should be equal to $\Delta_{20}^g \equiv E(2_g)$ in the strong-coupling limit. The differences are presented in Table 1. The quantity, $\Delta_{32}^{\gamma} - \Delta_{20}^g$, is negative in ^{164}Er , ^{168}Yb , and ^{180}W according to a recent experiment by Graetzer et al.,² while in Os nuclei it is always positive. This quantity is roughly evaluated by perturbation as follows:

$$\frac{\Delta_{32}^{\gamma} - \Delta_{20}^g}{\Delta_{20}^g} = -\frac{4}{3} \left(\frac{E(2_g)}{E_{\beta}} \right)^2 + \frac{4}{9} \left(\frac{E(2_g)}{E_{\gamma}} \right)^2$$

The first term arises from the beta-vibration-rotation interaction. As far as only the interaction between the ground band and the $K = 2$ gamma-vibrational band, is taken in account, the second term should vanish. This is why in the Davydov-Chaban model this quantity is always negative. The second term appears when the interaction between the $K = 2$ gamma-vibrational state and the $K = 0$ (two-phonon) gamma-vibrational state is taken into account. Therefore the fact mentioned above may infer the presence of the two-phonon gamma-vibrational band which plays an important role in the Os region.

REFERENCES

1. G. T. Emery, et al., Phys. Rev. 129, 2597 (1963).
2. Graetzer, et al., Nucl. Phys. 76, 1 (1966).

Table I.


Nucleus	$E(2_g)$ (keV)	$E(2_\gamma)$ (keV)	$(\Delta_{32}^\gamma - \Delta_{20}^g) / \Delta_{20}^g$ (%)	$(\Delta_{42}^\gamma - \Delta_{42}^g) / \Delta_{42}^g$ (%)
^{182}Os	128	892	14.9	9.7
^{184}Os	122	942	17.2	7.6
^{186}Os	137	768	3.7	1.7
^{188}Os	155	633	1.3	2.8
^{190}Os	187	557	6.4	10.2
^{192}Os	206	489	-1.9	
^{158}Dy	98.7	945	1.3	1.0
^{164}Er	91.3	858	-3.6	-4.8
^{168}Yb	87.9	986	-4.4	-5.5
^{180}W	103.6	828	-9.3	-11.1

FIGURE CAPTIONS

Fig. 1(a). Gamma ray spectra in $^{187}\text{(p,2n)}^{186}\text{Os}$ reaction.


Fig. 1(b). Gamma ray spectra in $^{185}\text{(p,2n)}^{184}\text{Os}$ reaction.

Fig. 2. Systematics of the ground band and the gamma-vibrational bands in even Os nuclei.



NU-10185

Fig. 1(a)



CALIFORNIA COMPUTER PRODUCTS, INC. ANALYZER, L.A. 90000 CHEM. NO. 00

NU-10184

Fig. 1(b)



Ground and gamma-vibrational bands
in even Os nuclei




Fig. 2

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

