UCRL-17303

University of California

Ernest O. Lawrence Radiation Laboratory

MULTIPLE NUCLEAR EXCITATION OF ROTATIONAL LEVELS IN ^{152}Sm AND ^{154}Sm

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

To be submitted to Physics Letters

UCRL-17303
Preprint

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

MULTIPLE NUCLEAR EXCITATION OF ROTATIONAL LEVELS IN ^{152}Sm AND ^{154}Sm

B. G. Harvey, D. L. Hendrie, O. N. Jarvis,
Jeannette Mahoney and J. Valentin

December 1966

MULTIPLE NUCLEAR EXCITATION OF ROTATIONAL LEVELS IN ^{152}Sm AND ^{154}Sm*

B. G. Harvey, D. L. Hendrie, O. N. Jarvis,†

Jeannette Mahoney, and J. Valentin‡

Lawrence Radiation Laboratory
University of California
Berkeley, California

December 1966

Abstract

By observing the inelastic scattering of 50 MeV alpha particles from samarium, we have made a definite correlation between reversed phase angular distributions and double excitation of second excited rotational levels. For the first time, we have also seen triple excitation of third excited rotational levels.

*This work was performed under the auspices of the U. S. Atomic Energy Commission

†Permanent Address: AERE Harwell, Didcot, Berks., England

‡Permanent Address: CENG, University of Grenoble, Grenoble, France

The scattering of medium energy alpha particles from intermediate weight nuclei show distinctive angular distributions for elastic and inelastic scattering. These angular distributions are marked by distinct diffraction maxima and minima and result from the strong absorption of the incident alpha particles. Simple excitations of collective levels in even nuclei yield characteristic distributions for each angular momentum transfer (i.e., level spin) at small angles and asymptotically approach the Blair phase rule (BPRI) at large angles. Specifically, even parity states (2^+, 4^+) asymptotically produce diffraction maxima in phase with the elastic scattering minima. Such angular distributions can be fit quite well by the Fraunhofer model[1] modified by a slightly diffuse edge[2].

More recently, however, several examples of low lying states of known even parity have been observed to give rise to angular distributions with reversed phases[3]. Moreover these distributions were found to have flatter envelopes than those of the simple excitation. The combination of these two properties define the characteristics of Blair phase rule two (BPRII) distributions. It was found that the states giving rise to such distributions could often be thought of as two-surfon levels, such as second rotational states (4^+) or two-phonon vibrational states (0^+, 2^+, 4^+ triplet). Thus a correspondence was made between (BPRII) angular distributions and the "double excitation" of two surfon collective states. However, in no case was the collective nature of the level well established.

To explain these results, recourse was made to the extended optical model of Rost and Austern[4]. It is known that two second order terms of the

transition matrix can give rise to double excitation. The first involves the second order term in the expansion of the nuclear potential in powers of the deformation. Although this term can be simply handled in DWBA approximation, it has been shown that this term alone cannot account for the experimental distributions[5]. The second term, corresponding to an excitational cascade through the one-surfon level, has been calculated only after further approximations have been made. It was then shown that this term, with interference effects from the first, can explain BPRII distributions. Austern and Blair[6] have developed such a theory in detail in which, however, the two terms cannot be separated. Calculations using the coupled-channels method have given similar results[7]. All these calculations depend on the purity of the two-surfon states; admixtures of this state with states of different character lead to results that are more complicated to predict.

In order to investigate the validity of the correspondence between double excitation and BPRII, we chose to look at the inelastic scattering of 50 MeV alpha particle from ^{152}Sm and ^{154}Sm. The requirement of experimentally accessible pure two-surfon states are well met by the rotational levels of the rare earths. The deformations, β, of these two nuclei are known to be .304 and .351 respectively[8].

Self-supporting samarium target foils about 300 $\mu g/cm^2$ thick were obtained by heating isotopic samarium oxide in the presence of lanthanum metal[9]. Scattered alpha particles were detected in four cooled lithium-drifted silicon detectors; energy spectra were stored in a multichannel analyser. The total beam passing through the target was collected and measured with a Faraday cup and checked with a fixed angle

monitor counter. Target thicknesses were obtained by normalizing the elastic cross section to Rutherford scattering at small angles. Total experimental resolutions were about 45 keV, sufficient to resolve the 2^+ from the ground state at most angles. Particular care was taken to reduce backgrounds and low energy tails by using a highly polished cylindrical-edged gold plated analyzing slit located far from the target area. Similar care was taken with the counter collimators. Nevertheless, backgrounds proved to be our most severe experimental problem, and tails from the large elastic peaks at forward angles obscured much of the inelastic data below about 25 degrees. Extraction of the inelastic cross sections was aided by use of a computer program which approximated the spectra by a series of gaussian peaks plus background[10]. Sample samarium spectra at larger angles are shown in fig. 1.

The inelastic differential cross sections to the 2^+, 4^+ and 6^+ levels of ^{152}Sm and ^{154}Sm are shown in figs. 2 and 3. These levels were all clearly resolved from adjacent inelastic levels except for the 6^+ level of ^{152}Sm. This level could not be separated from the 0^+ level at 920 keV, but we do not expect significant contribution from the 0^+ level. Inspections of the 2^+ and 4^+ distributions for both target nuclei shows the expected BPRI distribution for the 2^+ levels and BPRII distributions for the 4^+ levels. Both the reversed phase and the flatter angular distribution of the 4^+ levels are apparent. In addition, the cross sections are roughly in the ratio to be expected from the additional factor β^2 which would multiply the double excitation cross section. Therefore we can establish a definite correspondence between known two-surfon levels and BPRII distributions for these nuclei.

We consider next the angular distributions of the 6^+ levels. These distributions have still another reversal of phase. The envelopes of their oscillations are even flatter than those of the 4^+ levels. The magnitude of their cross sections is reduced from that of the 4^+ levels again by about a factor of β^2. In addition, the ^{154}Sm data at about 25° indicates a minimum where one would not be expected for a direct $\ell=6$ transition. We denote these characteristics by BPRIII and, by analogy with the 4^+ levels, relate them to "triple excitation" of the previously known 6^+ three-surfon members of the ground state rotational bands.

Extension of the Austern-Blair model to include triple excitation in the Fraunhofer limit provides qualitative agreement with our results. However the Austern-Blair model does not include coulomb excitation, which is certainly important for the excitation of the 2^+ level, but probably not for the other states. More important, deformations as large as those in the samarium isotopes lead to such strong coupling of the 2^+ and elastic channels that the DWBA approach is inappropriate. A coupled-channels program with coulomb excitation is indicated for more quantitative calculations.

The precise measurement of small amounts of higher order multipole deformations awaits such an improved theory. The constancy of the phase reversals and depth of the minima shown by our data are evidence that any such higher order terms are small. By means of the Austern-Blair model, we can place upper limits on these higher multipole mixtures. These limits are 0.1 for β_4 with a positive sign and 0.03 for β_6 with a negative sign. These limits are smaller by a factor of 3 for β's with the opposite sign. The limits compare with those of reference 11.

The regularity of the multiple excitation distributions also argues for a simplification of the excitation process. In addition to the two paths available to the double excitation, a triple excitation might also proceed via a mixed single-excitation, double-excitation cascade through either the 2^+ or 4^+ states. We feel that such a complicated mixture of excitation mechanisms would result in an angular distribution more featureless than the sharply structured patterns we see. The derivative relationship of ref. 5 seems to apply to these results.

We wish to thank Claude Ellsworth for fabrication of the targets and Prof. J. S. Blair for stimulating and revealing discussion.

References

1) J. S. Blair, Phys. Rev. 115 (1959) 928

2) E. V. Inopin and Yu. A. Berezhnoy, Nucl. Phys. 63 (1965) 689

3) R. Beurtey, P. Catillon, R. Chaminade, M. Crut, H. Faraggi, A. Papineau, J. Saudinos, and J. Thirion, Compt. Rend. 252 (1961) 1756

4) E. Rost and N. Austern, Phys. Rev. 120 (1960) 1375

5) N. Austern, R. M. Drisko, E. Rost, and G. R. Satchler, Phys. Rev. 128 (1962) 733

6) N. Austern and J. S. Blair, Ann. Phys. 33 (1965) 15

7) B. Buck, Phys. Rev. 127 (1962) 940; J. Wills (Ph.D. Thesis) University of Washington (1963)

8) P. H. Stelson and L. Grodzins, Nuclear Data 1 (1965) 21

9) L. Westgaard and S. Bjørnholm, Nucl. Instr. and Methods 42 (1966) 77

10) A. Springer (Ph.D. Thesis) University of California Lawrence Radiation Laboratory Report UCRL-11681 (May 1965)

11) B. Elbek, M. Kregar, and P. Vedelsky, Nucl. Phys. 86 (1966) 385

Figure Captions

Fig. 1. Energy spectra for the inelastic scattering of 50 MeV alphas from ^{152}Sm and ^{154}Sm plotted on a logarithmic scale. Positions of excited states are indicated.

Fig. 2. Differential cross sections for the 2^+, 4^+, and 6^+ members of the ground state rotational band of ^{152}Sm. Relative errors are, in general, smaller than the data points.

Fig. 3. Differential cross sections for the 2^+, 4^+, and 6^+ members of the ground state rotational band for ^{154}Sm. Relative errors are, in general, smaller than the data points. We did not find excitation of the 0^+ state at 1.012 MeV.


Counts
^{152}Sm (α,α′)
50 MeV
θ = 52°
^{154}Sm (α,α′)
50 MeV
θ = 55°
1000
100
10
1
1.236 3+
1.087 2+
1.045 3+
.963 1−
.811 2+
.712 6+ + .685 0+
.366 4+
.122 2+
g.s. 0+
1.012 3−
.905 8+ + .912 1−
.545 6+
.267 4+
.082 2+
g.s. 0+
MUB 13965

Fig. 1


Q = - 0.122 MeV
(2+)
152Sm (α,α′)
Eα = 50 MeV
Q = -0.366 MeV
(4+)
Q = -0.712 MeV
(6+)
dσ/dΩ (mb/sr)
100
10.0
1.0
0.1
0.01
20
30
40
50
60
70
θc.m. (deg)
MUB-13733

Fig. 2


^{154}Sm (α,α′)
E_α = 50 MeV
Q = -0.082 MeV
(2+)
Q = -0.267 MeV
(4+)
Q = -0.545
(6+)
100
10
1.0
0.1
0.01
0.001
20
30
40
50
60
70
dσ/dΩ (mb/sr)
$\theta_{c.m}$ (deg)
MUB-13952

Fig. 3

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.