UCRL-18175
cy. 2

# University of California

# Ernest O. Lawrence Radiation Laboratory

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545*

## LEVEL AND ISOMER SYSTEMATICS IN EVEN Sn ISOTOPES

T. Yamazaki, G. T. Ewan, and S. G. Prussin

April 16, 1968

RECEIVED
LAWRENCE
RADIATION LABORATORY
MAY 15 1968
LIBRARY AND
DOCUMENTS SECTION

Berkeley, California

UCRL-18175
cy. 2

## DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UCRL-18175

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

# LEVEL AND ISOMER SYSTEMATICS IN EVEN Sn ISOTOPES

T. Yamazaki, G. T. Ewan, and S. G. Prussin

April 16, 1968

# Level and Isomer Systematics in Even Sn Isotopes

T. Yamazaki*, G.T. Ewan† and S.G. Prussin‡

Lawrence Radiation Laboratory, University of California
Berkeley, California, U.S.A.

ABSTRACT

Levels of even Sn isotopes (A = 108 - 118) have been studied in Cd($\alpha$, xn) reactions. Nanosecond time analysis of $\gamma$-rays has revealed the systematic presence of isomers.

A method of systematically studying the existence of isomeric states with nanosecond life-times populated in ($\alpha$, xn) reactions has been discussed earlier[1]. We have used this method to study isomeric states in even-Sn isotopes from A = 108 to 118. The present study has revealed the existence of new isomers in A = 112 to 118 in addition to the known negative-parity (5- and 7-) isomers in $^{116}$Sn, $^{118}$Sn, and $^{120}$Sn[2-5]. The energies of these new levels appear to vary systematically with A. Theoretical considerations suggest that new isomers from A = 114 to 118 may arise from the $(h_{11/2})^2$ neutron configuration.

In our experiments metallic targets of separated cadmium isotopes were bombarded with $\alpha$-particles from the Berkeley 88-inch

---

* Now at Department of Physics, University of Tokyo, Bunkyo-ku, Tokyo, Japan.

† Permanent address: Chalk River Nuclear Laboratories, Chalk River, Ontario, Canada.

‡ Now at Department of Nuclear Engineering, University of California, Berkeley, California, U.S.A.

cyclotron. The beam energy was varied from 28 MeV to 50 MeV to control the yield of the ($\alpha$, 2n), ($\alpha$, 3n), and ($\alpha$, 4n) reaction products. Gamma-rays from the targets were observed with Ge(Li) detectors of different active volumes and the time distributions of the gamma-rays were measured by the method described in reference 1. The spectra were initially surveyed either with 2048 channels for energy analysis and 2 time channels (prompt and $\sim$ 50 nsec delayed) or with 1024 channels for energy analysis and 4 time channels (prompt, 25, 50, and 75 nsec delayed). These experiments provided immediate evidence of the existence of isomers and a rough estimate of transition life-times. A typical spectrum observed in the $^{110}$Cd + 28 MeV $\alpha$ reaction is shown in Fig. 1. This was taken with a 7 cm$^2$ (surface area) x 1 cm (thickness) Ge(Li) detector.

Accurate estimates of half-lifes were obtained by recording spectra with 256 channels for energy analysis and 16 channels for time analysis. The lower limit of observable half-lives in these runs was about 4 nsec. (In the survey runs, the lower limit was about 7 nsec). The time spacing of the cyclotron beam bunches placed an upper limit of about 300 nsec for accurate life-time measurements. Longer-lived isomers were easily identified but their half-lives could not be determined with accuracy. The half-lives of isomers in $^{112}$Sn to $^{118}$Sn were measured with this technique. Typical time distributions for gamma rays from the $^{110}$Cd($\alpha$, 2n)$^{112}$Sn reaction are shown in Fig. 2. The transitions shown all have a half-life of 14 nsec and are presumably fed by the same isomeric decay.

The results of the present measurements are summarized in Fig. 3. This figure includes level assignments derived from previous studies[6]. The isomeric states implied from our results are shown by heavy lines. Their positions rely upon expected systematic trends in the level schemes and additional experiments are required to establish with certainty the energies and spin assignments of these levels. The detailed analysis of the experimental data will be described elsewhere[7].

There have been several theoretical calculations of the level structure of tin isotopes[8-11]. Kisslinger and Sorensen[8] used a pairing plus quadrupole - quadrupole interaction. More recently Arvieu and co-workers[9] have made extensive calculations using the pairing scheme and a Gaussian type of residual interaction and later a surface delta interaction[10]. The experimental data shown in Fig. 3 allow us to make the following qualitative comparisons with the theoretical predictions. For reference we show the quasi-particle energies for the $(h_{11/2})^2$, $(d_{5/2})^2$ and $(g_{7/2})^2$ configurations used by Arvieu[11].

1) The energy of the first 2+ level increases with N from N = 58 to N = 64, where it reaches a maximum, and then starts to decrease. This is the result of a slight shell effect at N = 64.

2) The energy of the 4+ level shows a dependence on N similar to that of the 2+ level energy, except at N = 64 where there is an appreciable depression. This is not reproduced in Arvieu's earlier calculations but is in qualitative agreement with later calculations for N $\geq$ 64 by Plastino, Arvieu and Moskowski[10]

in which a surface delta interaction was used. A qualitative understanding of the dip in energy may also be obtained by considering the structure of the 4+ state. Below N = 64 its structure is dominated by the $(d_{5/2})^2$, $(g_{7/2})^2$ and $d_{5/2}g_{7/2}$ components. Above N = 64 the structure is dominated by the $(h_{11/2})^2$ component. At N = 64 all components contribute approximately equally and lead to a lowering in the energy of the state. The 2+ state is less affected because more configurations are available.

3) The 5- state $(h_{11/2}, s_{1/2})$ or $(h_{11/2}, d_{3/2})$ is isomeric for A = 116, 118 and 120, but not for A = 114, where the energy difference between the 5- and the 4+ levels is too great for the E1 transition to have a life-time measurable in our experiments. The 7- state $(h_{11/2}, d_{3/2})$ is isomeric for A = 118 and 120 due to the highly hindered E2 decay, but not for A $\lesssim$ 116 as it feeds the 6- state via a fast M1 transition. The 7- level for A = 114 could not be identified, presumably because it lay too high for observation. The systematics of these negative parity states are consistent with the assumed behaviour of the $h_{11/2}$ orbit.

4) In addition to the negative-parity isomers, we observed at least one long-lived isomer (> 100 nsec) for each isotope in the region N $\gtrsim$ 64. On the other hand, no such isomers were seen in the region N < 64. The half-life of the isomer in $^{116}$Sn has been determined by Chang, Hageman and Yamazaki[12] to be 0.8 μsec. From the quasi-particle states available

these long-lived isomers are tentatively assigned to the 10+ state of the $(h_{11/2})^2$ configuration. For $N \geq 64$, the 8+ and 10+ states of the $(h_{11/2})^2$ configuration are expected to lie around 3 MeV and, as no other configuration contributes to these high spin states, they should remain relatively unperturbed. The energy spacing between the 8+ and 10+ levels is estimated to be $\sim$ 100 keV and the 10+ state could be expected to have a half-life of $\sim$ 1 μsec. In the region $N < 64$, the $(h_{11/2})^2$ energy becomes so large that this configuration may no longer be the sole contributor to the lowest 8+ and 10+ states. Four quasi-particle states and vibrational states may also contribute. This may explain why the long-lived isomer does not exist in this region.

5) It is interesting that for $N < 64$ the highly-populated levels terminate at 6+. This is consistent with the fact that no higher spin state is available at low energy with the d5/2 and g7/2 orbits. The half-life of the 6+ state in $^{112}$Sn was found to be 14 nsec., which is comparable to the Weisskopf estimate for an E2 transition. It is striking that no delayed state was observed for the lighter nuclei, although the level structure is unchanged. This implies that the B(E2, 6+ → 4+) value increases rapidly as N decreases. This is in disagreement with the pairing theory which predicts that the B(E2, 6+ → 4+) should decrease due to the UU-VV factor. This will be discussed in detail in the forthcoming paper[7].

In conclusion we emphasize the possible presence of $(h_{11/2})^2_{10+} \rightarrow (h_{11/2})^2_{8+}$ isomers in the region $N \geq 64$, which preserve pure quasi-particle states. In the region $N < 64$, such an isomer is not observed and the ground band terminates at a spin of 6+. The discontinuance of the ground band structure at $N \leq 64$ is also seen in the Z = 52 Te isotopes studied by Bergström et al[13]. These facts raise the question as to the real nature of the ground band of levels and the relation between the so-called vibrational sequence and the lowest seniority scheme. Further experiments are planned to elucidate these problems.

We are grateful to Drs. J.M. Hollander and B.G. Harvey for support and encouragement, to Prof. Mottelson and Drs. T. Udagawa and R.A. Broglia for stimulating discussions, and to Prof. I. Perlman and J.O. Rasmussen for the hospitality of the Lawrence Radiation Laboratory.

References

1) T. Yamazaki and G.T. Ewan, Phys. Letters 24B, 278 (1967); Nucl. Instr. and Methods, in press (1968).

2) H.H. Bolotin, A.C. Li and A. Schwarzschild, Phys. Rev. 124, 213 (1961).

3) H. Ikegami and T. Udagawa, Phys. Rev. 124, 1518 (1961).

4) H.H. Bolotin, Phys. Rev. 136, B1557, B1566 (1964).

5) E. Bodenstedt et al., Nucl. Phys. 89, 305 (1966).

6) Data taken from C.M. Lederer, J.M. Hollander and I. Perlman; "Table of Isotopes, 6th Edition" (John Wiley and Sons, New York, 1967).

7) T. Yamazaki, G.T. Ewan and S.G. Prussin, in preparation.

8) L.S. Kisslinger and R.A. Sorensen, Mat. Fys. Medd. Dan. Vid. Selsk. 32, No. 9 (1960).

9) R. Arvieu, E. Baranger, M. Veneroni, M. Baranger and V. Gillet, Phys. Letters 4, 119 (1963).

10) A. Plastino, R. Arvieu and S.A. Moszkowski, Phys. Rev. 145, 837 (1966).

11) R. Arvieu, Ann. de Phys. 8, 407 (1963).

12) C.H. Chang, G.B. Hageman and T. Yamazaki, private communication (September, 1967).

13) I. Bergström, C.J. Herrlander, A. Kerek and A. Lukko, to be published; Contribution to the International Conference on Nuclear Structure (Tokyo, September, 1967).

Fig. 1 Prompt and delayed energy spectra of $\gamma$ rays emitted in $^{110}Cd$ + 28 MeV $\alpha$ reaction. Most $\gamma$-rays are due to the $^{110}Cd$ $(\alpha,2n)$ $^{112}Sn$ reaction. Delayed $\gamma$-rays in $^{18}F$, $^{19}F$ and $^{19}Ne$ from the backing and impurity are also identified.

Fig. 2 Time distribution curves for three prominent delayed $\gamma$-rays of $^{112}Sn$.

Fig. 3 The level and isomer systematics revealed in the present study. Levels of the same spin and parity are connected by broken lines. Previously known levels are $^{112}Sn$(2+), $^{114}Sn$(2+ and 3-), $^{116}Sn$(2+, 3-, 5-, 4+, 6- and 7-), $^{118}Sn$(2+, 4+, 5- and 7-), $^{120}Sn$(2+, 4+, 5- and 7-), $^{122}Sn$(2+ and 4+), and $^{124}Sn$(2+ and 4+). References are given in the recent compilation of Lederer, Hollander and Perlman[6].


110Cd + 28 MeV α 7 cc Ge (Li)
18F 184.4
18F 197.3
19Ne 238.5
263.3
112Sn 301.9
Prompt
511.0
113Sn 661.3
112Sn 806.6
18F 939.2
112Sn 992.8
112Sn 1257.7
Delayed
≈ 50 nsec
72Ge 694
Counts
Channel numbers

Fig. 1


Time (nsec)
100
50
0
$^{110}$Cd(α, 2n) $^{112}$Sn
$E_\alpha$ = 28 MeV
1258 keV
14.6 ± 0.6 nsec
993 keV
13.7 ± 0.6 nsec
302 keV
13.6 ± 0.6 nsec
Counts (arbitrary)
10
5
2
8
12
16
Channel Number

Fig. 2



$(h_{11/2})^2$
$(d_{5/2})^2$
$(g_{7/2})^2$
>0.5 μs
3232
>0.5 μs
3090
>0.5 μs
3088
7-
2913
2777
5-
2817
6-
0.35 μs
2393
0.23 μs
2580
11 μs
2490
20 ns
2326
5.5 ns
2280
2180
6+
<7 ns
2365
<7 ns
2480
14 ns
2552
3-
2274
2368
2266
2285
2180
2100
4+
2111
2199
2201
2189
2+
1205
1211
1258
1299
1293
1231
1171
1140
1131
0+
0
$^{108}_{50}Sn_{58}$
$^{110}_{50}Sn_{60}$
$^{112}_{50}Sn_{62}$
$^{114}_{50}Sn_{64}$
$^{116}_{50}Sn_{66}$
$^{118}_{50}Sn_{68}$
$^{120}_{50}Sn_{70}$
$^{122}_{50}Sn_{72}$
$^{124}_{50}Sn_{74}$
XBL6712-5877


Fig. 3

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.