UCRL-18449

cy. 2

University of California

Ernest O. Lawrence Radiation Laboratory

RECEIVED
LAWRENCE
RADIATION LABORATORY
OCT 21 1968
LIBRARY AND
DOCUMENTS SECTIO

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

RECONSTITUTION ("REPAIR") MECHANISMS IN THE RADIOLYSIS OF AQUEOUS BIOCHEMICAL SYSTEMS: INHIBITIVE EFFECTS OF THIOLS

John Holian and Warren M. Garrison

September 6, 1968

Berkeley, California

UCRL-18449
cy. 2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to Nature

UCRL-18449
Preprint

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

RECONSTITUTION ("REPAIR") MECHANISMS IN THE RADIOLYSIS OF AQUEOUS BIOCHEMICAL SYSTEMS: INHIBITIVE EFFECTS OF THIOLS

John Holian and Warren M. Garrison

September 6, 1968

RECONSTITUTION ("REPAIR") MECHANISMS IN THE RADIOLYSIS OF AQUEOUS BIOCHEMICAL SYSTEMS: INHIBITIVE EFFECTS OF THIOLS

John Holian and Warren M. Garrison

Lawrence Radiation Laboratory
University of California
Berkeley, California

September 6, 1968

The hydrated electron, e^-_{aq}, and the OH radical are the principal products of the radiation-induced decomposition of liquid water; with γ-rays the 100 ev yields[1] of these reactive intermediates are closely approximated by the values: $G_{e^-_{aq}} \simeq G_{OH} \simeq 2.6$. While certain classes of organic compounds are known to react quantitatively with both e^-_{aq} and OH in oxygen-free solution, still, the observed yield for net destruction of such solutes may be quite low as compared to the G value for water decomposition. In interpreting the radiation chemistry of such systems the concept of a reconstitution or back-reaction must be invoked.

For example, in the γ-radiolysis of oxygen-free solutions of the pyrimidine bases, both e^-_{aq} and OH add preferentially to the 5,6 double bond[2,3]

$$e^-_{aq} + B + H_2O \rightarrow \dot{B}H + OH^- \qquad (2)$$

$$OH + B \rightarrow \dot{B}OH \qquad (3)$$

---

[1] Czapski, G., Advan. Chem. Ser., 81 106 (1968).

[2] Weiss, J., Prog. Nucleic Acid Res. and Mol. Biol., 3 103 (1964).

[3] Kamal, A. and Garrison, W.M., Nature 206 1315 (1965).

where the rate constants k_2, k_3 fall in the range 10^9 to 10^{10} M^{-1} sec^{-1} (ref. 4,5). Yet, the observed yield for base destruction in millimolar solutions is uniformly low with $G(-B) < 0.9$. The evidence is that a back-reaction[3]

$$\dot{B}H + \dot{B}OH \rightarrow 2B + H_2O \qquad (4)$$

leads to a reconstitution of the base.

We now find that certain labile compounds such as ascorbic acid and the thiol, cysteine, are effective at low concentrations in blocking reaction (4) by virtue of the H-atom transfer reaction

$$\dot{B}H + R'H \rightarrow BH_2 + R' \qquad (5)$$

where BH_2 represents the dihydro derivative of the pyrimidine nucleus. The effects of added cysteine on the yields for radiolytic reduction of cytosine in oxygen-free solution are shown in table 1A. Note that dihydrocytosine is unstable and spontaneously hydrolyses to yield ammonia and dihydrouracil[6] as the experimentally observed product. The product dihydrouracil was isolated and characterized both chromatographically and spectrophotometrically. The data of table 1A show that the yield for the radiolytic reduction of the 5,6 double bond of cytosine is essentially quantitative in the presence of either cysteine or ascorbic acid at low concentrations.

---

[4] Hart, E.J., Thomas, J.K., Gordon, S., Radiation Res., Suppl. 4 74 (1964).

[5] Scholes, P., Shaw, P., Willson, R. L., and Ebert, M., "Pulse Radiolysis" p. 151 Academic Press, New York, 1965.

[6] Green, M., and Cohen, S.C., J. Biol Chem., 228 601 (1957).

Similarly, in the radiolysis of primary amides and monosubstituted primary amides (peptides) in oxygen-free neutral solution, the reducing and oxidizing species e^-_{aq} and OH are removed through reactions of the type

$$e^-_{aq} + RCONHR + H_2O \rightarrow R\dot{C}(OH)NHR + OH^- \quad (6)$$

$$OH + RCONHR \rightarrow \dot{P} + H_2O \quad (7)$$

where k_6, k_7 fall in the range 10^7 to 10^8 M^{-1} sec^{-1} (ref. 7,8). Combination of $R\dot{C}(OH)NHR$ with like species or with the radical $\dot{P}$ would lead to formation of ketonic products. However, the combined yield of such products is low with $G(>CO) \simeq 0.2$; the evidence is that the reconstitution reaction

$$R\dot{C}(OH)NHR + \dot{P} \rightarrow 2RCONHR \quad (8)$$

represents the major stoichimetry for removal of organic radical in such systems.[8]

In accord with this, we find that cysteine effectively blocks the reconstitituion reaction 8 through the step

$$R\dot{C}(OH)NHR + R'H \rightarrow RCHO + NH_2R + R'$$

and, as shown in table 1B, the reductive deamination of the peptide bond in the presence of cysteine at low concentrations is essentially quantitative with $G(CH_3CHO) \simeq 2.5$.

---

[7] Willix, R.L.S., Garrison, W.M., Radiation Res., 32 452 (1967).

[8] Rodgers, M.A.J., Garrison, W.M., J. Phys. Chem., 72 758 (1968).

We find then that (a) the presence of cysteine (or ascorbic acid) at low concentrations leads to a very marked enhancement in the radiolytic lability of the pyrimidine and peptide moieties in oxygen-free solution, and that (b) such enhancement arises as a consequence of the blocking by the second solute of the reconstitution reactions formulated in equations 4, 8.

These results would appear to have interesting applications in the study and identification of reductive processes involving reactions of e^-_{aq} with unsaturated organic functions both in vitro and in vivo.

This work was performed under the auspices of the United States Atomic Energy Commission.

Table 1. Effect of cysteine and ascorbic acid ($R'H$) on the γ-ray induced reduction of pyrimidine (C = C) and peptide (C = O) linkages in oxygen-free solutions*.

A. Cytosine → dihydrouracil (BH_2) + ammonia†

solution	$R'H$ ‡	$G(BH_2)$
.05 M Cytosine, pH7	none	< 0.9
.05 M Cytosine, pH7	cysteine, 2.5×10^{-3} M	2.8
.05 M Cytosine, pH7	ascorbic acid, 1.5×10^{-3} M	2.9

B. N-ethylacetamide → acetaldehyde (RCHO) + ethylamine

solution	$R'H$	G(RCHO)
1M N-ethylacetamide, pH7	none	< 0.1
1M N-ethylacetamide, pH7	cysteine, 4×10^{-4} M	2.8

*At dosages below $\sim 2.5 \times 10^{18}$ ev/gm.

†Dihydrocytosine is unstable and hydrolyzes spontaneously to yield dihydrouracil and ammonia.

‡The indicated concentrations of $R'H$ give the maximum enhancement in the yield of reduced products.

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.