Submitted to Inorganic Chemistry

UCRL-18870
Preprint

RECEIVED
LAWRENCE
RADIATION LABORATORY

MAY 13 1969

LIBRARY AND
DOCUMENTS SECTION

THE CRYSTAL STRUCTURE OF $Cs_2(B_9C_2H_{11})Co(B_8C_2H_{10})Co(B_9C_2H_{11})\cdot H_2O$, A SALT OF A THREE-ICOSAHEDRAL-FRAGMENT METALLOCARBORANE

David St. Clair, Allan Zalkin, and David H. Templeton

April 1969

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

LAWRENCE RADIATION LABORATORY
UNIVERSITY of CALIFORNIA BERKELEY

UCRL-18870

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to Inorganic Chemistry

UCRL-18870
Preprint

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California 94720

AEC Contract No. W-7405-eng-48

CONTRIBUTION FROM THE LAWRENCE RADIATION LABORATORY AND DEPARTMENT OF CHEMISTRY, UNIVERSITY OF CALIFORNIA, BERKELEY, CALIFORNIA 94720

The Crystal Structure of $Cs_2(B_9C_2H_{11})Co(B_8C_2H_{10})Co(B_9C_2H_{11})\cdot H_2O$, a Salt of a Three-Icosahedral-Fragment Metallocarborane[1]

By David St. Clair, Allan Zalkin, and David H. Templeton

---

(1) Work done under the auspices of the U. S. Atomic Energy Commission.

The crystal structure of the monohydrate of the cesium salt of the bis-π-[(3)-1,2-dicarbollylcobalt]-π-(3,6)-1,2-dicarbacanastide ion, $Cs_2Co_2B_{26}C_6H_{32}\cdot H_2O$, has been determined from 3649 independent X-ray data obtained from a single-crystal by counter methods. The crystals are monoclinic, space group $P2_1/c$. Four molecules are in the unit cell with $\underline{a}$ = 7.089 Å., $\underline{b}$ = 19.240 Å., $\underline{c}$ = 20.682 Å., $\underline{\beta}$ = 98.00°. The calculated density is $\underline{\rho}$ = 1.871 g/cc. The structure was refined by least squares to a conventional R factor of 3.4 %. The carborane anion, $[(B_9C_2H_{11})Co(B_8C_2H_{10})Co(B_9C_2H_{11})]^{-2}$, consists of three linked icosahedra. Two corners of a central ten light atom icosahedron are occupied by cobalts each of which is common to another icosahedron containing eleven light atoms. Five of the positions in the icosahedra are occupied by carbons but, because of disorder, two positions contain half-boron-half-carbon atoms. The cages are staggered across the cobalts and all carbons are as close to each other as is possible in this configuration. All hydrogens except those in the water molecule were located.

Introduction

A novel anion of the carborane type, $[(B_9C_2H_{11})Co(B_8C_2H_{10})Co(B_9C_2H_{11})]^{-2}$, the [bis[undecahydro-1,2-dicarba-3-cobalta-closo-dodecaborano]-[3',3';3,6]-commo-decahydro-1,2-dicarba-3,6-dicobalta-closo-dodecaborate](2-) ion, has recently been prepared[2,3]. Francis and Hawthorne proposed that the anion

---

(2) J. M. Francis and M. F. Hawthorne, J. Am. Chem. Soc., 90, 1663(1968).

(3) This name is an extension of the ACS-approved nomenclature rules given in Inorg. Chem., 7, 1945(1968), which do not explicitly cover the naming of ions in which a single polyhedron is completed by two identical groups attached at two different positions in the cage. The name originally given by Francis and Hawthorne in reference 2 is the bis-π-[(3)-1,2-dicarbollylcobalt]-π-(3,6)-1,2-dicarbacanastide ion. Both names fail to describe the rotational isomerism of the π bonding.

---

consisted of two of the previously known eleven atom icosahedral fragments each having a cobalt atom bonded to the open face with the cobalts linked together via a new ten atom icosahedral fragment. In this paper, we report the results of an X-ray analysis of the cesium salt of this anion, $Cs_2(Co_2B_{26}C_6H_{32})\cdot H_2O$, shown in Figure 1. We confirm the structure assigned for the anion by Francis and Hawthorne. We show that the rotational configuration of the cages in the crystal are those in which the cages are staggered and that the carbons are as close to each other as possible in this staggered arrangement.

Experimental

The dark red crystals were received from Professor Hawthorne of the University of California, Riverside. One was cut to the dimensions 0.11 x 0.12 x 0.23 mm and was glued to a glass fiber with its b axis coincident with the rotation axis of the fiber.

The space group and approximate cell dimensions were determined from oscillation, Weissenberg and precession photographs. Using a manual General Electric XRD-5 diffractometer, space group absences were checked and accurate unit cell dimensions were obtained from measurements of the h00, 0k0, and 00ℓ reflections with the b axis of the crystal parallel to the φ axis of the goniostat. Intensity data were measured with this manually operated diffractometer using a quarter circle, Eulerian cradle goniostat, and a scintillation counter equipped with a pulse height discriminator for detection of the diffracted beam. Molybdenum Kα X-rays were used (Mo $K\alpha_1$; $\lambda = 0.70926$ Å.) after the Kβ radiation was filtered out by a 0.003 inch thick Zr filter placed in front of the receiving slit. One complete set of independent reflections (including space group absences) with $2\theta \leq 45^\circ$ ($\sin\theta/\lambda = 0.995$) was measured using the stationary crystal, stationary counter technique by taking ten second counts at the maximum intensity settings. Excluding space group extinctions, there were 3649 independent reflections of which 344 had intensities less than their estimated standard deviations.

The absorption coefficient is $\mu = 38\ cm^{-1}$. Absorption effects on the data were neglected. It is estimated that the scale factors which would be applied to the intensity data would very by less than 12 %.

Calculations were done on a CDC 6600 computer. Lorentz and polarization corrections were applied and the data were converted to structure factors. Refinements of parameters were done with our least squares program which

minimizes the quantity $\Sigma w(|F_o| - |F_c|)^2/\Sigma w|F_o|^2$ where F_o and F_c are the observed and calculated structure factors, respectively. The weighting factor, $w = [\sigma(F_o)]^{-2}$, is derived from the standard deviation of the intensity, $\sigma(I)$, estimated by the equation, $\sigma(I) = [I + 2I_b + (0.05I)^2]^{\frac{1}{2}}$, where I_b is the number of counts in the background and I is the number of counts on the peak minus I_b.

All atoms were considered to be in their neutral valence state. The scattering factors of Cromer and Waber[4] were used for all except the hydrogen

---

(4) D. T. Cromer and J. T. Waber, Acta Cryst., 18, 104(1965).

---

atoms whose scattering factors are those of Stewart, Davidson and Simpson[5].

---

(5) R. F. Stewart, E. R. Davidson and W. T. Simpson, Table II, J. Chem. Phys., 42, 3175(1965).

Cromer's corrections for anomalous dispersion[6] applied to the scattering

---

(6) D. T. Cromer, Acta Cryst., 18, 17(1965).

---

factors of the Cs and Co are $\Delta f' = -0.5$ and $\Delta f'' = +2.6$ electrons and $\Delta f' = +0.4$ and $\Delta f'' = +1.0$ electrons, respectively. The anisotropic temperature factors were of the form $\exp(-\beta_{11}h^2-\beta_{22}k^2-\beta_{33}\ell^2-2\beta_{12}hk-2\beta_{13}h\ell-2\beta_{23}k\ell)$.

Crystal Data

The crystals were found to be monoclinic with unit cell dimensions, measured at room temperature (~22°), $\underline{a}$ = 7.089 ± 0.004 Å., $\underline{b}$ = 19.240 ± 0.014 Å., $\underline{c}$ = 20.682 ± 0.004 Å., $\underline{\beta}$ = 98.00 ± 0.06 degrees. There are four formula units, $Cs_2Co_2B_{26}C_6OH_{34}$, per unit cell. The density is calculated to be $\underline{\rho}$ = 1.87 g/cc. No density measurement was made because the sample was lost.

The observed extinction rules, $\underline{h0\ell}$, $\underline{\ell} = \underline{2n}$ and $\underline{0k0}$, $\underline{k} = \underline{2n}$, correspond to space group $P2_1/c$. All atoms lie on the general symmetry related positions, ± ($\underline{x,y,z}$ and $\underline{x,\frac{1}{2}-y,\frac{1}{2}+z}$).

Determination of the Structure

The positions of the cesium and cobalt atoms were deduced from sections through a three-dimensional Patterson function synthesis. Full matrix least squares refinement of a scale factor and the positional and isotropic thermal parameters of the Cs and Co atoms gave R = 0.208. These atoms were used to calculate approximate phases for the reflections and a three-dimensional difference Fourier was synthesized from which the coordinates of the thirty two boron and carbon atoms were found. Since a water of crystallization was not expected, the oxygen atom which appeared in this Fourier was disregarded. When the positions and thermal parameters of these thirty two light atoms, all with the scattering power of boron, were incorporated into the refinement, the R value dropped to 0.105. Inspection of the bond distances and thermal parameters allowed location of five of the six carbon atoms and gave indications that the sixth carbon was disordered between only two atomic positions in one of the icosahedral cages. So in all further refinements, five of these atoms were given the scattering power of carbon and two were given the scattering power of half carbon and half boron. Following another series of refinements in which the Cs and Co atoms were given anisotropic thermal parameters, another difference Fourier was prepared in which again appeared the oxygen atom as well as thirty two of the thirty four hydrogen atoms. The two hydrogens in the water molecule did not appear. After some blunders in the intensity data were corrected, a final series of full matrix least squares refinement was carried out in which the scattering factors of the Cs and Co atoms were corrected for anomalous dispersion and zero weight was assigned to all reflections which had $I < \sigma(I)$. A scale factor, three positional parameters for each of the sixty nine atoms, six anisotropic temperature parameters for each Cs and Co atom, individual isotropic temperature parameters for the thirty three light

atoms excluding hydrogens, and an average temperature parameter for all thirty two hydrogens were refined. The conventional R value for all non-zero-weighted data reached R = 0.034 for 3305 data, the weighted R value was $R_2 = 0.039$ and the conventional R value calculated with all 3649 data was R = 0.041. The standard deviation of observation of unit weight was 1.2 and the r.m.s. values of $(|F_o| - |F_c|)/\sigma(F_o)$ averaged over small intensity intervals approximated this value over the entire intensity range. No parameter in the final refinement differed from that of the previous refinement by an amount greater than one tenth of its estimated standard deviation.

Results and Discussion of the Structure

The final values for positional and thermal parameters are given in Table I for the heavy atoms, Table II for the oxygen, carbon, and boron atoms and in Table III for the hydrogen atoms. The observed and calculated structure factors are given in Table IV.

A drawing of the anion skeleton is given in Figure 1. It consists of two eleven-light atom icosahedral fragments, arbitrarily labeled cages I and III, each having a cobalt ion completing the icosahedron by bonding to the apex position of each cage. The cobalts are linked via the basket-shaped canastide ion completing a third icosahedron, labeled cage II, by bonding to the open faces in this bridging fragment. The cages are staggered with respect to each other across the cobalts. That is, atoms number 7 and 22 are equidistant from atoms 12 and 13. The two carbon atoms in each of cages I and II were located at positions 7 and 8 and 12 and 13, respectively. Disorder was found, however, in cage III, one carbon being localized at position 22 and the other being equally distributed between positions 23 and 26 adjacent to position 22. This disorder indicates the presence of three rotational isomers in the crystal differing only in the directions of rotation of the end cages. Beginning from a hypothetical configuration in which the carbons are all eclipsed, each end cage is rotated 36° either to the right or to the left. The isomers consist of an enantiomorphic pair with point symmetry 2 and a third isomer with point symmetry m. Since the external shape is not sensitive to the carbon positions, one might expect disorder at both ends of the anion. We have no explanation for the fact that all the disorder seems to be at cage III. Ignoring the distinction between boron and carbon, the anion has approximately 2mm (C_{2v}) point symmetry in which one of the mirror planes contains atoms in positions 12, 13, 18, and 19 of cage II and the other mirror plane is perpendicular to

TABLE I

Coordinates($\times 10^5$) and Thermal Parameters($\times 10^2$)[a] of Heavy Atoms[b]

Atom	x	y	z	B_{11}	B_{22}	B_{33}	B_{12}	B_{13}	B_{23}
Cs(1)	21468(6)	27505(2)	17044(2)	434(2)	336(2)	318(2)	017(1)	054(1)	-012(1)
Cs(2)	05321(6)	02870(2)	34230(2)	503(2)	280(2)	449(2)	031(2)	106(2)	-044(1)
Co(1)	23982(10)	76225(4)	58130(3)	250(3)	195(3)	229(3)	020(2)	-002(2)	010(2)
Co(2)	25769(10)	57752(4)	64795(3)	231(3)	214(3)	239(3)	-002(2)	002(2)	019(2)

(a) Anisotropic thermal parameters, $\underline{B}$, in units of $\mathring{A}^2$, are given by $\underline{B} = 4\beta_{ij}/a_i^* a_j^*$ where a_i^* is the ith reciprocal cell length.

(b) Estimated standard deviations are given in parentheses following the parameter.

TABLE II

Coordinates($x10^4$) and Thermal Parameters($x10^2$) of the Light Atoms[a,b]

Atom	x	y	z	B(Å^2)
O(1)	7075(8)	5640(3)	8419(3)	643(12)
		Atoms in Cage I		
B(1)	3781(10)	8967(4)	4548(3)	329(13)
B(2)	3175(10)	8098(4)	4321(3)	321(13)
B(3)	5330(10)	8283(4)	4825(3)	326(13)
B(4)	4890(9)	8951(4)	5360(3)	294(12)
B(5)	2461(9)	9202(4)	5183(3)	292(12)
B(6)	1410(10)	8660(4)	4538(3)	309(13)
C(7)	3825(8)	7631(3)	5016(3)	293(11)
C(8)	4806(8)	8116(3)	5604(3)	295(11)
B(9)	3196(9)	8664(3)	5870(3)	278(12)
B(10)	0994(9)	8496(3)	5351(3)	266(12)
B(11)	1482(9)	7788(4)	4818(3)	294(12)
		Atoms in Cage II		
C(12)	1639(7)	6581(3)	5850(3)	232(9)
C(13)	3550(8)	6763(3)	6316(2)	236(10)
B(14)	3294(9)	7459(3)	6797(3)	264(12)
B(15)	0822(9)	7708(4)	6597(3)	299(12)
B(16)	-0137(9)	7143(3)	5945(3)	251(11)
B(17)	3360(9)	6575(3)	7117(3)	264(12)
B(18)	1585(10)	7179(4)	7298(3)	318(13)
B(19)	-0546(9)	6978(3)	6762(3)	275(12)
B(20)	-0047(9)	6259(3)	6267(3)	260(11)
B(21)	0951(9)	6289(3)	7108(3)	284(12)

Atoms in Cage III

C(22)	4005(8)	5193(3)	5879(3)	297(11)
BC(23)	5182(9)	5261(3)	6618(3)	292(11)
B(24)	3661(9)	5032(3)	7168(3)	256(12)
B(25)	1423(9)	4807(3)	6691(3)	265(12)
BC(26)	1816(9)	4931(3)	5893(3)	306(12)
B(27)	5659(11)	4569(4)	6115(4)	363(14)
B(28)	5409(10)	4434(4)	6947(3)	337(13)
B(29)	3074(9)	4136(3)	6982(3)	284(12)
B(30)	1870(10)	4088(4)	6165(3)	312(13)
B(31)	3488(10)	4355(4)	5649(4)	339(14)
B(32)	4321(10)	3856(4)	6337(3)	325(13)

(a) Numbering system used is shown in Figure 1.

(b) BC refers to disordered atoms, half boron and half carbon.

TABLE III

Coordinates($\times 10^3$) of the Hydrogen Atoms[a,b,c]

Atom	x	y	z	Atom	x	y	z
H(1)	410	931	415	H(17)	473	650	745
H(2)	326	787	385	H(18)	142	741	786
H(3)	677	687	973	H(19)	-200	703	687
H(4)	615	927	556	H(20)	-123	592	601
H(5)	200	975	526	H(21)	038	601	753
H(6)	038	885	415	H(22)	420	551	552
H(7)	430	725	498	H(23)	623	559	667
H(8)	603	793	587	H(24)	379	516	767
H(9)	352	892	635	H(25)	000	484	684
H(10)	956	864	548	H(26)	105	507	556
H(11)	049	738	458	H(27)	701	455	589
H(12)	189	642	544	H(28)	662	430	733
H(13)	457	671	617	H(29)	269	376	738
H(14)	439	777	697	H(30)	067	372	601
H(15)	027	821	675	H(31)	331	424	518
H(16)	860	722	555	H(32)	466	332	628

(a) Number refers to the position in the cage occupied by the atom to which the hydrogen atom is attached. See Figure 1.

(b) Refined average temperature factor is $\underline{B} = 2.6 \pm 0.2\ \text{Å}^2$.

(c) Standard deviations for $\underline{x}$, $\underline{y}$, and $\underline{z}$ are ±0.007, ±0.003, and ±0.003, respectively.

TABLE IV

Observed and Calculated Structure Factors

Reflections given zero weight in the final refinement are marked with an asterisk.

OBSERVED AND CALCULATED STRUCTURE FACTORS

FCA(0,0,0) = 2960

L	FOB	FCA
H,K= -1, -6		
21	0	8*
H,K= -1, -5		
21	62	65
H,K= -1, -4		
21	106	110
H,K= -1, -3		
21	0	1*
22	0	13*
H,K= -1, -2		
21	33	37
22	93	100
H,K= -1, -1		
21	80	79
22	17	19*
H,K= 0, 0		
2	199	181
4	737	761
6	637	648
8	328	333
10	341	353
12	531	539
14	42	61
16	264	273
18	68	74
20	42	53
22	38	37
H,K= 0, 1		
1	78	69
3	84	76
4	365	356
5	201	196
6	88	85
7	210	208
8	295	291
9	325	320
10	73	77
11	46	42
12	168	168
13	105	103
14	3	31
15	104	108
16	256	256
17	41	30
18	93	95
19	71	70
20	122	121
21	64	63
22	0	34*
H,K= 0, 2		
0	117	112
1	216	199
2	167	172
3	40	37
4	219	212
5	147	150
6	176	174
7	56	56
8	264	262
9	67	68
10	269	270
11	64	69
12	174	173
13	0	17*
14	155	153
15	30	34
16	142	144
17	0	22*
18	21	20
19	0	2*
20	219	215
21	0	12*
H,K= 0, 3		
1	192	178
2	296	291
3	307	296
4	572	575
5	8	6*
6	120	116
7	247	243
8	676	623
9	46	36
10	76	77
11	152	152
12	85	85
13	59	82
14	194	193
15	0	23*
16	126	125
17	70	70
18	78	77
19	60	63
20	114	117
21	0	21*
H,K= 0, 4		
0	69	76
1	418	402
2	260	247
3	44	44
4	328	321
5	183	180
6	363	354
7	259	255
8	295	299
9	45	46
10	65	67
11	35	31
12	514	521
13	197	199
14	47	48
15	73	73
16	136	131
17	80	77
18	119	124
19	163	153
20	101	99
21	C	4*
H,K= 0, 5		
1	331	324
2	10C	104
3	538	529
4	111	106
5	146	142
6	176	173
7	328	318
8	105	100
9	340	335
10	39	39
11	49	46
12	217	223
13	216	223
14	166	164
15	75	60
16	176	173
17	0	4*
18	108	113
19	65	63
20	36	34
21	69	65
H,K= 0, 6		
0	369	353
1	19	16
2	90	93
3	15	14
4	101	99
5	173	122
6	139	137
7	84	86
8	82	85
9	16	25
10	127	11
11	48	44
12	152	156
13	124	121
14	0	9*
15	116	116
16	0	11*
17	0	7*
18	38	34
19	73	78
20	28	35
21	90	85
H,K= 0, 7		
1	137	134

L	FOB	FCA
	2	277 275
	8	84 84
	4	401 403
	5	94 95
	6	67 63
	7	137 137
	8	367 363
	9	84 84
	10	91 92
	11	270 272
	12	67 70
	13	0 10*
	14	82 84
	15	101 103
	16	150 156
	17	116 124
	18	66 63
	19	54 50
	20	104 99
	21	80 79
	22	0 2*
H,K= 0, 0		
2	199	181
4	737	761
6	637	648
8	328	333
10	341	353
12	531	539
14	42	61
16	264	273
18	68	74
20	42	53
22	38	37
H,K= 0, 1		
1	78	69
3	84	76
4	365	356
5	201	196
6	88	85
7	210	208
8	295	291
9	325	320
10	73	77
11	46	42
12	168	168
13	105	103
14	3	31
15	104	108
16	256	256
17	41	30
18	93	95
19	71	70
20	122	121
21	64	63
22	0	34*
H,K= 0, 2		
0	117	112
1	216	199
2	167	172
3	40	37
4	219	212
5	147	150
6	176	174
7	56	56
8	264	262
9	67	68
10	269	270
11	64	69
12	174	173
13	0	17*
14	155	153
15	30	34
16	142	144
17	0	22*
18	21	20
19	0	2*
20	219	215
21	0	12*
H,K= 0, 3		
1	192	178
2	296	291
3	307	296
4	572	575
5	8	6*
6	120	116
7	247	243
8	676	623
9	46	36
10	76	77
11	152	152
12	85	85
13	59	82
14	194	193
15	0	23*
16	126	125
17	70	70
18	78	77
19	60	63
20	114	117
21	0	21*
H,K= 0, 4		
0	69	76
1	418	402
2	260	247
3	44	44
4	328	321
5	183	180
6	363	354
7	259	255
8	295	299
9	45	46
10	65	67
11	35	31
12	514	521
13	197	199
14	47	48
15	73	73
16	136	131
17	80	77
18	119	124
19	163	153
20	101	99
21	C	4*
H,K= 0, 5		
1	331	324
2	10C	104
3	538	529
4	111	106
5	146	142
6	176	173
7	328	318
8	105	100
9	340	335
10	39	39
11	49	46
12	217	223
13	216	223
14	166	164
15	75	60
16	176	173
17	0	4*
18	108	113
19	65	63
20	36	34
21	69	65
H,K= 0, 6		
0	369	353
1	19	16
2	90	93
3	15	14
4	101	99
5	173	122
6	139	137
7	84	86
8	82	85
9	16	25
10	127	11
11	48	44
12	152	156
13	124	121
14	0	9*
15	116	116
16	0	11*
17	0	7*
18	38	34
19	73	78
20	28	35
21	90	85
H,K= 0, 7		
1	137	134

L	FOB	FCA
	8	53 46
	9	213 205
	10	108 107
	11	78 77
	12	116 108
	13	178 170
	14	33 26
	15	91 92
H,K= 0, 16		
	20	73 72
	21	17 27*
H,K= 1, 3		
	16	0 8*
	17	137 131
	18	0 2*
H,K= 1, 11		
	16	91 90
	17	0 2*
H,K= 1, 7		
	12	63 65
	13	186 193
	14	117 117
	15	52 55
	16	142 149
	17	0 1*
H,K= 1, 8		
	13	143 149
	14	130 134
	15	38 41
	16	135 135
	17	0 4*
H,K= 1, 9		
	13	233 238
	14	350 357
	15	128 124
	16	139 143
	17	9 44
H,K= 1, 10		
	13	40 40
	14	160 162
	15	0 1*
H,K= 1, 11		
	13	138 143
	14	0 3*
H,K= 1, 12		
	13	0 8*
H,K= 1, 13		
	12	0 8*
H,K= 1, 14		
	11	0 4*
H,K= 1, 15		
	10	0 4*
H,K= 1, 16		
	9	0 1*
H,K= 1, 17		
	8	0 3*
H,K= 1, 18		
	7	0 3*
H,K= 1, 19		
	6	0 3*
H,K= 1, 20		
	5	0 2*
H,K= 1, 21		
	4	0 4*
H,K= 1, 22		
	3	0 3*
H,K= 1, 23		
	2	0 3*
H,K= 1, 24		
	1	0 3*
H,K= 1, 25		
	0	0 3*
H,K= 2, 0		
	2	0 1*
H,K= 2, 1		
	1	0 1*
H,K= 2, 2		
	0	0 1*
H,K= 2, 3		
	1	0 1*
H,K= 2, 4		
	2	0 1*
H,K= 2, 5		
	3	0 1*
H,K= 2, 6		
	4	0 1*
H,K= 2, 7		
	5	0 1*
H,K= 2, 8		
	6	0 1*
H,K= 2, 9		
	7	0 1*
H,K= 2, 10		
	8	0 1*
H,K= 2, 11		
	9	0 1*
H,K= 2, 12		
	10	0 1*
H,K= 2, 13		
	11	0 1*
H,K= 2, 14		
	12	0 1*
H,K= 2, 15		
	13	0 1*
H,K= 2, 16		
	14	0 1*
H,K= 2, 17		
	15	0 1*
H,K= 2, 18		
	16	0 1*
H,K= 2, 19		
	17	0 1*
H,K= 2, 20		
	18	0 1*
H,K= 2, 21		
	19	0 1*
H,K= 2, 22		
	20	0 1*
H,K= 2, 23		
	21	0 1*
H,K= 2, 24		
	22	0 1*
H,K= 2, 25		
	23	0 1*
H,K= 3, 0		
	3	0 1*
H,K= 3, 1		
	2	0 1*
H,K= 3, 2		
	1	0 1*
H,K= 3, 3		
	0	0 1*
H,K= 3, 4		
	1	0 1*
H,K= 3, 5		
	2	0 1*
H,K= 3, 6		
	3	0 1*
H,K= 3, 7		
	4	0 1*
H,K= 3, 8		
	5	0 1*
H,K= 3, 9		
	6	0 1*
H,K= 3, 10		
	7	0 1*
H,K= 3, 11		
	8	0 1*
H,K= 3, 12		
	9	0 1*
H,K= 3, 13		
	10	0 1*
H,K= 3, 14		
	11	0 1*
H,K= 3, 15		
	12	0 1*
H,K= 3, 16		
	13	0 1*
H,K= 3, 17		
	14	0 1*
H,K= 3, 18		
	15	0 1*
H,K= 3, 19		
	16	0 1*
H,K= 3, 20		
	17	0 1*
H,K= 3, 21		
	18	0 1*
H,K= 3, 22		
	19	0 1*
H,K= 3, 23		
	20	0 1*
H,K= 3, 24		
	21	0 1*
H,K= 3, 25		
	22	0 1*
H,K= 4, 0		
	4	0 1*
H,K= 4, 1		
	3	0 1*
H,K= 4, 2		
	2	0 1*
H,K= 4, 3		
	1	0 1*
H,K= 4, 4		
	0	0 1*
H,K= 4, 5		
	1	0 1*
H,K= 4, 6		
	2	0 1*
H,K= 4, 7		
	3	0 1*
H,K= 4, 8		
	4	0 1*
H,K= 4, 9		
	5	0 1*
H,K= 4, 10		
	6	0 1*
H,K= 4, 11		
	7	0 1*
H,K= 4, 12		
	8	0 1*
H,K= 4, 13		
	9	0 1*
H,K= 4, 14		
	10	0 1*
H,K= 4, 15		
	11	0 1*
H,K= 4, 16		
	12	0 1*
H,K= 4, 17		
	13	0 1*
H,K= 4, 18		
	14	0 1*
H,K= 4, 19		
	15	0 1*
H,K= 4, 20		
	16	0 1*
H,K= 4, 21		
	17	0 1*
H,K= 4, 22		
	18	0 1*
H,K= 4, 23		
	19	0 1*
H,K= 4, 24		
	20	0 1*
H,K= 4, 25		
	21	0 1*
H,K= 5, 0		
	5	0 1*
H,K= 5, 1		
	4	0 1*
H,K= 5, 2		
	3	0 1*
H,K= 5, 3		
	2	0 1*
H,K= 5, 4		
	1	0 1*
H,K= 5, 5		
	0	0 1*
H,K= 5, 6		
	1	0 1*
H,K= 5, 7		
	2	0 1*
H,K= 5, 8		
	3	0 1*
H,K= 5, 9		
	4	0 1*
H,K= 5, 10		
	5	0 1*
H,K= 5, 11		
	6	0 1*
H,K= 5, 12		
	7	0 1*
H,K= 5, 13		
	8	0 1*
H,K= 5, 14		
	9	0 1*
H,K= 5, 15		
	10	0 1*
H,K= 5, 16		
	11	0 1*
H,K= 5, 17		
	12	0 1*
H,K= 5, 18		
	13	0 1*
H,K= 5, 19		
	14	0 1*
H,K= 5, 20		
	15	0 1*
H,K= 5, 21		
	16	0 1*
H,K= 5, 22		
	17	0 1*
H,K= 5, 23		
	18	0 1*
H,K= 5, 24		
	19	0 1*
H,K= 5, 25		
	20	0 1*
H,K= 6, 0		
	6	0 1*
H,K= 6, 1		
	5	0 1*
H,K= 6, 2		
	4	0 1*
H,K= 6, 3		
	3	0 1*
H,K= 6, 4		
	2	0 1*
H,K= 6, 5		
	1	0 1*
H,K= 6, 6		
	0	0 1*
H,K= 6, 7		
	1	0 1*
H,K= 6, 8		
	2	0 1*
H,K= 6, 9		
	3	0 1*
H,K= 6, 10		
	4	0 1*
H,K= 6, 11		
	5	0 1*
H,K= 6, 12		
	6	0 1*
H,K= 6, 13		
	7	0 1*
H,K= 6, 14		
	8	0 1*
H,K= 6, 15		
	9	0 1*
H,K= 6, 16		
	10	0 1*
H,K= 6, 17		
	11	0 1*
H,K= 6, 18		
	12	0 1*
H,K= 6, 19		
	13	0 1*
H,K= 6, 20		
	14	0 1*
H,K= 6, 21		
	15	0 1*
H,K= 6, 22		
	16	0 1*
H,K= 6, 23		
	17	0 1*
H,K= 6, 24		
	18	0 1*
H,K= 6, 25		
	19	0 1*
H,K= 7, 0		
	7	0 1*
H,K= 7, 1		
	6	0 1*
H,K= 7, 2		
	5	0 1*
H,K= 7, 3		
	4	0 1*
H,K= 7, 4		
	3	0 1*
H,K= 7, 5		
	2	0 1*
H,K= 7, 6		
	1	0 1*
H,K= 7, 7		
	0	0 1*
H,K= 7, 8		
	1	0 1*
H,K= 7, 9		
	2	0 1*
H,K= 7, 10		
	3	0 1*
H,K= 7, 11		
	4	0 1*
H,K= 7, 12		
	5	0 1*
H,K= 7, 13		
	6	0 1*
H,K= 7, 14		
	7	0 1*
H,K= 7, 15		
	8	0 1*
H,K= 7, 16		
	9	0 1*
H,K= 7, 17		
	10	0 1*
H,K= 7, 18		
	11	0 1*
H,K= 7, 19		
	12	0 1*
H,K= 7, 20		
	13	0 1*
H,K= 7, 21		
	14	0 1*
H,K= 7, 22		
	15	0 1*
H,K= 7, 23		
	16	0 1*
H,K= 7, 24		
	17	0 1*
H,K= 7, 25		
	18	0 1*
H,K= 8, 0		
	8	0 1*
H,K= 8, 1		
	7	0 1*
H,K= 8, 2		
	6	0 1*
H,K= 8, 3		
	5	0 1*
H,K= 8, 4		
	4	0 1*
H,K= 8, 5		
	3	0 1*
H,K= 8, 6		
	2	0 1*
H,K= 8, 7		
	1	0 1*
H,K= 8, 8		
	0	0 1*
H,K= 8, 9		
	1	0 1*
H,K= 8, 10		
	2	0 1*
H,K= 8, 11		
	3	0 1*
H,K= 8, 12		
	4	0 1*
H,K= 8, 13		
	5	0 1*
H,K= 8, 14		
	6	0 1*
H,K= 8, 15		
	7	0 1*
H,K= 8, 16		
	8	0 1*
H,K= 8, 17		
	9	0 1*
H,K= 8, 18		
	10	0 1*
H,K= 8, 19		
	11	0 1*
H,K= 8, 20		
	12	0 1*
H,K= 8, 21		
	13	0 1*
H,K= 8, 22		
	14	0 1*
H,K= 8, 23		
	15	0 1*
H,K= 8, 24		
	16	0 1*
H,K= 8, 25		
	17	0 1*
H,K= 9, 0		
	9	0 1*
H,K= 9, 1		
	8	0 1*
H,K= 9, 2		
	7	0 1*
H,K= 9, 3		
	6	0 1*
H,K= 9, 4		
	5	0 1*
H,K= 9, 5		
	4	0 1*
H,K= 9, 6		
	3	0 1*
H,K= 9, 7		
	2	0 1*
H,K= 9, 8		
	1	0 1*
H,K= 9, 9		
	0	0 1*
H,K= 9, 10		
	1	0 1*
H,K= 9, 11		
	2	0 1*
H,K= 9, 12		
	3	0 1*
H,K= 9, 13		
	4	0 1*
H,K= 9, 14		
	5	0 1*
H,K= 9, 15		
	6	0 1*
H,K= 9, 16		
	7	0 1*
H,K= 9, 17		
	8	0 1*
H,K= 9, 18		
	9	0 1*
H,K= 9, 19		
	10	0 1*
H,K= 9, 20		
	11	0 1*
H,K= 9, 21		
	12	0 1*
H,K= 9, 22		
	13	0 1*
H,K= 9, 23		
	14	0 1*
H,K= 9, 24		
	15	0 1*
H,K= 9, 25		
	16	0 1*
H,K= 10, 0		
	10	0 1*
H,K= 10, 1		
	9	0 1*
H,K= 10, 2		
	8	0 1*
H,K= 10, 3		
	7	0 1*
H,K= 10, 4		
	6	0 1*
H,K= 10, 5		
	5	0 1*
H,K= 10, 6		
	4	0 1*
H,K= 10, 7		
	3	0 1*
H,K= 10, 8		
	2	0 1*
H,K= 10, 9		
	1	0 1*
H,K= 10, 10		
	0	0 1*
H,K= 10, 11		
	1	0 1*
H,K= 10, 12		
	2	0 1*
H,K= 10, 13		
	3	0 1*
H,K= 10, 14		
	4	0 1*
H,K= 10, 15		
	5	0 1*
H,K= 10, 16		
	6	0 1*
H,K= 10, 17		
	7	0 1*
H,K= 10, 18		
	8	0 1*
H,K= 10, 19		
	9	0 1*
H,K= 10, 20		
	10	0 1*
H,K= 10, 21		
	11	0 1*
H,K= 10, 22		
	12	0 1*
H,K= 10, 23		
	13	0 1*
H,K= 10, 24		
	14	0 1*
H,K= 10, 25		
	15	0 1*
H,K= 11, 0		
	11	0 1*
H,K= 11, 1		
	10	0 1*
H,K= 11, 2		
	9	0 1*
H,K= 11, 3		
	8	0 1*
H,K= 11, 4		
	7	0 1*
H,K= 11, 5		
	6	0 1*
H,K= 11, 6		
	5	0 1*
H,K= 11, 7		
	4	0 1*
H,K= 11, 8		
	3	0 1*
H,K= 11, 9		
	2	0 1*
H,K= 11, 10		
	1	0 1*
H,K= 11, 11		
	0	0 1*
H,K= 11, 12		
	1	0 1*
H,K= 11, 13		
	2	0 1*
H,K= 11, 14		
	3	0 1*
H,K= 11, 15		
	4	0 1*
H,K= 11, 16		
	5	0 1*
H,K= 11, 17		
	6	0 1*
H,K= 11, 18		
	7	0 1*
H,K= 11, 19		
	8	0 1*
H,K= 11, 20		
	9	0 1*
H,K= 11, 21		
	10	0 1*
H,K= 11, 22		
	11	0 1*
H,K= 11, 23		
	12	0 1*
H,K= 11, 24		
	13	0 1*
H,K= 11, 25		
	14	0 1*
H,K= 12, 0		
	12	0 1*
H,K= 12, 1		
	11	0 1*
H,K= 12, 2		
	10	0 1*
H,K= 12, 3		
	9	0 1*
H,K= 12, 4		
	8	0 1*
H,K= 12, 5		
	7	0 1*
H,K= 12, 6		
	6	0 1*
H,K= 12, 7		
	5	0 1*
H,K= 12, 8		
	4	0 1*
H,K= 12, 9		
	3	0 1*
H,K= 12, 10		
	2	0 1*
H,K= 12, 11		
	1	0 1*
H,K= 12, 12		
	0	0 1*
H,K= 12, 13		
	1	0 1*
H,K= 12, 14		
	2	0 1*
H,K= 12, 15		
	3	0 1*
H,K= 12, 16		
	4	0 1*
H,K= 12, 17		
	5	0 1*
H,K= 12, 18		
	6	0 1*
H,K= 12, 19		
	7	0 1*
H,K= 12, 20		
	8	0 1*
H,K= 12, 21		
	9	0 1*
H,K= 12, 22		
	10	0 1*
H,K= 12, 23		
	11	0 1*
H,K= 12, 24		
	12	0 1*
H,K= 12, 25		
	13	0 1*
H,K= 13, 0		
	13	0 1*
H,K= 13, 1		
	12	0 1*
H,K= 13, 2		
	11	0 1*
H,K= 13, 3		
	10	0 1*
H,K= 13, 4		
	9	0 1*
H,K= 13, 5		
	8	0 1*
H,K= 13, 6		
	7	0 1*
H,K= 13, 7		
	6	0 1*
H,K= 13, 8		
	5	0 1*
H,K= 13, 9		
	4	0 1*
H,K= 13, 10		
	3	0 1*
H,K= 13, 11		
	2	0 1*
H,K= 13, 12		
	1	0 1*
H,K= 13, 13		
	0	0 1*
H,K= 13, 14		
	1	0 1*
H,K= 13, 15		
	2	0 1*
H,K= 13, 16		
	3	0 1*
H,K= 13, 17		
	4	0 1*
H,K= 13, 18		
	5	0 1*
H,K= 13, 19		
	6	0 1*
H,K= 13, 20		
	7	0 1*
H,K= 13, 21		
	8	0 1*
H,K= 13, 22		
	9	0 1*
H,K= 13, 23		
	10	0 1*
H,K= 13, 24		
	11	0 1*
H,K= 13, 25		
	12	0 1*
H,K= 14, 0		
	14	0 1*
H,K= 14, 1		
	13	0 1*
H,K= 14, 2		
	12	0 1*
H,K= 14, 3		
	11	0 1*
H,K= 14, 4		
	10	0 1*
H,K= 14, 5		
	9	0 1*
H,K= 14, 6		
	8	0 1*
H,K= 14, 7		
	7	0 1*
H,K= 14, 8		
	6	0 1*
H,K= 14, 9		
	5	0 1*
H,K= 14, 10		
	4	0 1*
H,K= 14, 11		
	3	0 1*
H,K= 14, 12		
	2	0 1*
H,K= 14, 13		
	1	0 1*
H,K= 14, 14		
	0	0 1*
H,K= 14, 15		
	1	0 1*
H,K= 14, 16		
	2	0 1*
H,K= 14, 17		
	3	0 1*
H,K= 14, 18		
	4	0 1*
H,K= 14, 19		
	5	0 1*
H,K= 14, 20		
	6	0 1*
H,K= 14, 21		
	7	0 1*
H,K= 14, 22		
	8	0 1*
H,K= 14, 23		
	9	0 1*
H,K= 14, 24		
	10	0 1*
H,K= 14, 25		
	11	0 1*
H,K= 15, 0		
	15	0 1*
H,K= 15, 1		
	14	0 1*
H,K= 15, 2		
	13	0 1*
H,K= 15, 3		
	12	0 1*
H,K= 15, 4		
	11	0 1*
H,K= 15, 5		
	10	0 1*
H,K= 15, 6		
	9	0 1*
H,K= 15, 7		
	8	0 1*
H,K= 15, 8		
	7	0 1*
H,K= 15, 9		
	6	0 1*
H,K= 15, 10		
	5	0 1*
H,K= 15, 11		
	4	0 1*
H,K= 15, 12		
	3	0 1*
H,K= 15, 13		
	2	0 1*
H,K= 15, 14		
	1	0 1*
H,K= 15, 15		
	0	0 1*
H,K= 15, 16		
	1	0 1*
H,K= 15, 17		
	2	0 1*
H,K= 15, 18		
	3	0 1*
H,K= 15, 19		
	4	0 1*
H,K= 15, 20		
	5	0 1*
H,K= 15, 21		
	6	0 1*
H,K= 15, 22		
	7	0 1*
H,K= 15, 23		
	8	0 1*
H,K= 15, 24		
	9	0 1*
H,K= 15, 25		
	10	0 1*
H,K=

| | | |

the first one and bisects the bonds between atoms 12 to 13 and 18 to 19. The 2-fold rotation axis is along the intersection of these two mirror planes. If the carbon atom in position 8 of cage I were disordered equally between positions 8 and 11, the anion would have 2mm point symmetry with respect to all atoms including borons and carbons. The axis of cage I, Co(1) to B(1), makes an angle of 117° with the axis of cage III, Co(2) to B(32). The cages are arranged such that all the carbon atoms lie on the interior side of this angle and are as close to each other as is possible in this staggered configuration.

The bond distances between all the atoms in the icosahedral framework except hydrogens are listed in Table V and the distances between the hydrogen atoms and the atoms to which they are bonded are given in Table VI. The standard deviations quoted on the bond distances in these tables are calculated from the least squares estimates of the accuracy of the final positional parameters. Probably a better estimate of their accuracy is given by the larger root-mean-square deviations of equivalent bond distances from their average values, given in Table VII. A complete list of bond angles would be a rather long list indeed. Therefore, since both the cages I and III are analogous to the previously investigated eleven-atom icosahedral fragments[7], only the angles involving the

(7) A. Zalkin, T. E. Hopkins, and D. H. Templeton, Inorg. Chem., 5, 1189(1966).

ten-atom fragment, cage II, are reported in Table VIII.

The molecular packing in the crystal is shown in Figure 2 by a framework drawing of the projection of the contents of the unit cell seen from a view along the a axis. This drawing together with the distances given in Table IX reveals a series of interactions between the cesium ions, the oxygens of the water molecules and the hydrogen atoms attached to the anion. The cesium ions

TABLE V

Bond Distances

Atom 1	Atom 2	Distance (Å)	Atom 1	Atom 2	Distance (Å)
Co(1)	C(8)	2.051 ± .006	C(7)	B(2)	1.702 ± .009
Co(1)	C(7)	2.051 ± .006	B(11)	B(10)	1.815 ± .009
Co(1)	B(11)	2.094 ± .007	B(11)	B(2)	1.787 ± .009
Co(1)	B(10)	2.113 ± .007	B(11)	B(6)	1.772 ± .009
Co(1)	B(9)	2.081 ± .007	B(10)	B(9)	1.797 ± .009
Co(1)	C(12)	2.070 ± .005	B(10)	B(6)	1.775 ± .009
Co(1)	C(13)	2.061 ± .005	B(10)	B(5)	1.774 ± .009
Co(1)	B(14)	2.070 ± .006	B(9)	B(5)	1.776 ± .009
Co(1)	B(15)	2.100 ± .007	B(9)	B(4)	1.792 ± .009
Co(1)	B(16)	2.072 ± .006	B(3)	B(2)	1.762 ± .010
Co(2)	C(12)	2.065 ± .005	B(3)	B(4)	1.754 ± .009
Co(2)	C(13)	2.065 ± .005	B(2)	B(6)	1.759 ± .009
Co(2)	B(17)	2.053 ± .007	B(6)	B(5)	1.772 ± .009
Co(2)	B(20)	2.072 ± .006	B(5)	B(4)	1.776 ± .009
Co(2)	B(21)	2.101 ± .007	B(1)	B(3)	1.759 ± .010
Co(2)	C(22)	2.043 ± .006	B(1)	B(2)	1.775 ± .010
Co(2)	BC(23)	2.080 ± .006	B(1)	B(6)	1.779 ± .009
Co(2)	B(24)	2.087 ± .007	B(1)	B(5)	1.774 ± .009
Co(2)	B(25)	2.105 ± .007	B(1)	B(4)	1.754 ± .010
Co(2)	BC(26)	2.055 ± .006	C(12)	C(13)	1.561 ± .007
C(8)	C(7)	1.613 ± .008	C(12)	B(16)	1.719 ± .008
C(8)	B(9)	1.700 ± .008	C(12)	B(20)	1.713 ± .008
C(8)	B(3)	1.734 ± .009	C(13)	B(14)	1.695 ± .008
C(8)	B(4)	1.688 ± .009	C(13)	B(17)	1.720 ± .008
C(7)	B(11)	1.682 ± .008	B(14)	B(15)	1.808 ± .009

C(7)	B(3)	1.727 ± .009	B(14)	B(17)	1.823 ± .009
B(14)	B(18)	1.783 ± .009	BC(23)	B(28)	1.728 ± .009
B(15)	B(16)	1.791 ± .009	B(24)	B(25)	1.801 ± .009
B(15)	B(18)	1.791 ± .009	B(24)	B(28)	1.797 ± .009
B(15)	B(19)	1.768 ± .009	B(24)	B(29)	1.804 ± .009
B(16)	B(19)	1.783 ± .009	B(25)	BC(26)	1.729 ± .009
B(16)	B(20)	1.824 ± .009	B(25)	B(29)	1.790 ± .009
B(17)	B(18)	1.790 ± .009	B(25)	B(30)	1.814 ± .009
B(18)	B(19)	1.786 ± .009	BC(26)	B(30)	1.715 ± .009
B(19)	B(20)	1.786 ± .009	BC(26)	B(31)	1.747 ± .009
B(21)	B(17)	1.792 ± .009	B(27)	B(28)	1.772 ± .010
B(21)	B(18)	1.800 ± .009	B(27)	B(31)	1.748 ± .010
B(21)	B(19)	1.783 ± .009	B(28)	B(29)	1.764 ± .009
B(21)	B(20)	1.786 ± .009	B(29)	B(30)	1.787 ± .009
C(22)	BC(23)	1.643 ± .008	B(30)	B(31)	1.749 ± .010
C(22)	BC(26)	1.636 ± .008	B(32)	B(27)	1.764 ± .010
C(22)	B(27)	1.702 ± .009	B(32)	B(28)	1.775 ± .010
C(22)	B(31)	1.706 ± .009	B(32)	B(29)	1.784 ± .009
BC(23)	B(24)	1.729 ± .009	B(32)	B(30)	1.782 ± .009
BC(23)	B(27)	1.752 ± .009	B(32)	B(31)	1.750 ± .010

TABLE VI

Hydrogen to Carbon or Boron Distances(Å)[a]

Atoms in Cage I		Atoms in Cage II		Atoms in Cage III	
Atom	Distance	Atom	Distance	Atom	Distance
C(8)	1.02	C(12)	0.94	C(22)	0.99
C(7)	0.81	C(13)	0.82	BC(23)	0.97
B(11)	1.12	B(14)	1.01	B(24)	1.05
B(10)	1.12	B(15)	1.11	B(25)	1.10
B(9)	1.10	B(16)	1.14	BC(26)	0.86
B(3)	1.11	B(17)	1.12	B(27)	1.12
B(2)	1.08	B(18)	1.26	B(28)	1.11
B(6)	1.07	B(19)	1.09	B(29)	1.15
B(5)	1.13	B(20)	1.14	B(30)	1.12
B(4)	1.11	B(21)	1.14	B(31)	0.98
B(1)	1.09			B(32)	1.07

(a) Estimated standard deviations on all distances are ±0.05 Å.

TABLE VII

Averages of Equivalent Bond Distances(Å)[a,b]

Type	Number	Average	σ
Co - C	7	2.06	0.01
Co - B	11	2.09	0.02
C - B	12	1.71	0.02
B - B	48	1.78	0.02
C - H	5	0.92	0.10
B - H	25	1.11	0.05

(a) An average is given only is there are 5 or more values to include in the calculation of that average.

(b) The root-mean-square deviation from the average is given by $\sigma = [\Sigma\Delta^2/(n - 1)]^{\frac{1}{2}}$, where Δ_i is the difference between the ith bond distance and the average of n distances of the ith type.

TABLE VIII

Angles Involving the Cobalts and the Canastide Ion

Atom 1	Atom 2	Atom 3	Angle(deg)	Atom 1	Atom 2	Atom 3	Angle(deg)
C(12)	Co(1)	C(13)	44.4 ± .2	Co(1)	B(16)	B(15)	65.4 ± .3
C(12)	Co(2)	C(13)	44.4 ± .2	Co(2)	B(17)	B(21)	65.9 ± .3
C(12)	Co(1)	B(16)	49.0 ± .2	Co(2)	B(21)	B(17)	63.1 ± .3
C(13)	Co(1)	B(14)	48.5 ± .2	Co(2)	B(21)	B(20)	63.8 ± .3
C(12)	Co(2)	B(20)	48.9 ± .2	Co(2)	B(20)	B(21)	65.5 ± .3
C(13)	Co(2)	B(17)	49.4 ± .2	C(12)	C(13)	B(14)	112.9 ± .4
B(14)	Co(1)	B(15)	51.4 ± .2	C(12)	C(13)	B(17)	112.0 ± .4
B(15)	Co(1)	B(16)	50.8 ± .2	C(13)	C(12)	B(16)	112.0 ± .4
B(17)	Co(2)	B(21)	51.1 ± .2	C(13)	C(12)	B(20)	112.1 ± .4
B(20)	Co(2)	B(21)	50.7 ± .2	B(14)	C(13)	B(17)	64.5 ± .4
Co(1)	C(12)	C(13)	67.5 ± .2	B(16)	C(12)	B(20)	64.2 ± .4
Co(1)	C(13)	C(12)	68.1 ± .2	C(13)	B(14)	B(17)	58.4 ±..3
Co(2)	C(12)	C(13)	67.8 ± .2	C(13)	B(17)	B(14)	57.1 ± .3
Co(2)	C(13)	C(12)	67.8 ± .2	C(12)	B(16)	B(20)	57.7 ± .3
Co(1)	C(12)	B(16)	65.5 ± .3	C(12)	B(20)	B(16)	58.1 ± .3
Co(1)	C(13)	B(14)	66.1 ± .3	C(13)	B(14)	B(15)	104.7 ± .4
Co(2)	C(12)	B(20)	65.8 ± .3	C(12)	B(16)	B(15)	104.7 ± .4
Co(2)	C(13)	B(17)	64.9 ± .3	C(13)	B(17)	B(21)	104.9 ± .4
Co(1)	B(14)	C(13)	65.5 ± .3	C(12)	B(20)	B(21)	105.3 ± .4
Co(1)	B(16)	C(12)	65.4 ± .3	B(14)	B(15)	B(16)	105.5 ± .4
Co(2)	B(17)	C(13)	65.7 ± .3	B(17)	B(21)	B(20)	105.5 ± .4
Co(2)	B(20)	C(12)	65.3 ± .2	C(13)	B(14)	B(18)	103.9 ± .4
Co(1)	B(14)	B(15)	65.2 ± .3	C(13)	B(17)	B(18)	102.5 ± .4
Co(1)	B(15)	B(14)	63.5 ± .3	C(12)	B(16)	B(19)	102.7 ± .4
Co(1)	B(15)	B(16)	63.8 ± .3	C(12)	B(20)	B(19)	102.8 ± .4

B(15)	B(14)	B(18)	59.8 ± .4	B(16)	B(19)	B(21)	110.3 ± .4
B(21)	B(17)	B(18)	60.3 ± .4	B(18)	B(19)	B(20)	107.9 ± .5
B(15)	B(16)	B(19)	59.3 ± .4	Co(1)	C(13)	B(17)	125.1 ± .3
B(21)	B(20)	B(19)	59.9 ± .4	Co(1)	B(14)	B(17)	119.1 ± .3
B(14)	B(15)	B(18)	59.4 ± .4	Co(1)	B(14)	B(18)	118.5 ± .4
B(17)	B(21)	B(18)	59.8 ± .4	Co(1)	B(15)	B(18)	116.6 ± .3
B(16)	B(15)	B(19)	60.1 ± .4	Co(1)	B(15)	B(19)	116.9 ± .3
B(20)	B(21)	B(19)	60.0 ± .4	Co(1)	B(16)	B(19)	117.6 ± .3
B(14)	B(17)	B(18)	59.1 ± .4	Co(1)	B(16)	B(20)	118.4 ± .3
B(17)	B(14)	B(18)	59.5 ± .4	Co(1)	C(12)	B(20)	124.3 ± .3
B(16)	B(20)	B(19)	59.2 ± .4	Co(2)	C(12)	B(16)	124.7 ± .3
B(20)	B(16)	B(19)	59.3 ± .4	Co(2)	B(20)	B(16)	118.8 ± .3
B(17)	B(18)	B(15)	110.0 ± .5	Co(2)	B(20)	B(19)	118.0 ± .3
B(17)	B(18)	B(19)	106.7 ± .5	Co(2)	B(21)	B(19)	116.7 ± .3
B(14)	B(18)	B(19)	106.9 ± .5	Co(2)	B(21)	B(18)	116.2 ± .3
B(14)	B(18)	B(21)	109.3 ± .5	Co(2)	B(17)	B(18)	119.1 ± .3
B(15)	B(18)	B(21)	109.0 ± .5	Co(2)	B(17)	B(14)	118.4 ± .3
B(15)	B(19)	B(21)	110.9 ± .5	Co(2)	C(13)	B(14)	124.4 ± .3
B(15)	B(19)	B(20)	110.3 ± .5	Co(1)	C(12)	Co(2)	134.1 ± .2
B(16)	B(19)	B(18)	108.3 ± .5	Co(1)	C(13)	Co(2)	134.7 ± .2

and the anion cages are held in their relative orientations by the balance of the attraction of their unlike charges with the repulsive forces of the electrons in the cesium ions and the hydrogen atoms attached to the cages. All of the shortest distances to the oxygen, given in Table IX, correspond to reasonable van der Waals contacts expected for a normal water of hydration.

TABLE IX

Shortest Distances(Å) to Cesiums and Oxygen[a]

Atom	To O(1)	Atom	To Cs(1)	Atom	To Cs(2)
H(1)	2.76	H(18)	2.82	H(15)	2.95
H(24)	2.78	H(32)	2.95	H(4)	3.05
H(6)	2.78	H(17)	2.99	H(10)	3.08
H(17)	2.93	H(19)	3.00	H(24)	3.08
Cs(2)	3.029	H(11)	3.12	H(21)	3.10
Cs(1)	3.163	H(6)	3.15	H(6)	3.16
H(5)	3.21	H(29)	3.23	H(25)	3.25
H(21)	3.26	H(3)	3.24		
		H(30)	3.29		

(a) Estimated standard deviations are all ±0.05 Å. except O(1) to Cs(1) and to Cs(2) which are ±0.006 Å.

Acknowledgment

We wish to thank J. N. Francis and M. F. Hawthorne of the University of California, Riverside, for providing us with the crystals used in this work.

Figure Captions

Figure 1: Drawing of the $(Co_2B_{26}C_6H_{32})^{-2}$ anion (hydrogen atoms not shown).

Figure 2: Framework drawing of the projection of the unit cell contents down the a axis.



Figure I

XBL6810-6853


Co
Cs
H_2O
95
71
78
26
24
78
71
05
26
95
29
74
22
76
22
76
74
29
05
b
c sin β
0 1 2 3 A
1
0
1
0

XBL 681-83

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

TECHNICAL INFORMATION DIVISION
LAWRENCE RADIATION LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720