

eyo

OBSERVATION OF NMR IN AN ISOMERIC STATE
FOLLOWING A NUCLEAR REACTION

D. Quitmann, J. M. Jaklevic, and D. A. Shirley

April 1969

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

LAWRENCE RADIATION LABORATORY
UNIVERSITY of CALIFORNIA BERKELEY

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

OBSERVATION OF NMR IN AN ISOMERIC STATE
FOLLOWING A NUCLEAR REACTION*

D. Quitmann,[†] J. M. Jaklevic and D. A. Shirley

Lawrence Radiation Laboratory
University of California
Berkeley, California 94720

April 1969

The isomeric state in As^{73} with $I^\pi = 9/2^+$ at 426 keV was produced and aligned by the $\text{Ga}^{71}(\alpha,2n)$ reaction in a liquid Ga metal target. The resulting γ -ray anisotropy was attenuated by inducing NMR transitions with a 1 MHz rf field. We obtained $g = +1.146 \pm 0.007$ for the 426-keV state.

Nuclear magnetic resonance transitions in the 5.8- μ sec, 426 keV state of As^{73} have been induced by an external rf magnetic field and observed through their effect on the angular distribution of the gamma rays emitted. The isomeric state was produced and aligned by the $\text{Ga}^{71}(\alpha,2n)$ reaction on a liquid metal target. This result extends the classical NMR experiments to isomeric states with lifetimes in the 10^{-6} - 10^{-3} sec range. Using this technique, NMR studies of a large number of nuclear states (and of elements) should be possible, providing accurate values of the magnetic moments. It could also help elucidate the dynamics of environmental effects with characteristic times in the 10^{-6} - 10^{-3} sec range, and serve to determine hyperfine constants and Knight shifts for solute atoms.

The radiative detection of magnetic resonance in nuclei (NMR/RD) began with the positronium experiments of Deutsch and Brown.¹ Bloembergen and Temmer suggested combining magnetic resonance with nuclear orientation (NMR/ON),²

and Abragam and Pound proposed the use of perturbed angular correlations (NMR/PAC).³ These two methods as well as the alignment by nuclear reactions, have attracted increasing interest.⁴ From the standpoint of the applicability of NMR to metastable nuclear states in condensed matter, the present experimental situation can be summarized as follows. States with lifetimes longer than 10^3 sec are generally accessible to NMR/ON, usually employing lattices of ferromagnetic metals. For states with lifetimes of the order of 10^{-6} sec, NMR/PAC may be applicable, again using ferromagnetic hosts. However, since neither method is generally feasible for states in the $10^{-6} - 10^{+3}$ sec range, nuclear reactions must be used to provide the oriented nuclei. Several experiments have been reported on light beta - emitting nuclei with lifetimes in the upper half of this range (i.e. 10^{-2} sec to 10^2 sec).⁵ The purpose of the present work was to apply the NMR/RD method to one of the many γ -decaying isomeric states in the $10^{-6} - 10^{-3}$ sec range.

Preliminary measurements were made on several targets using (α, xn) reactions to produce the aligned isomers. In order to decide whether a NMR/RD experiment would be feasible, two criteria were employed: (1) with no external fields a time-integral γ -ray anisotropy of magnitude comparable to the one predicted for the expected alignment must be exhibited, and (2) this anisotropy must disappear when a (small) dc magnetic field is applied perpendicular to the beam-detector plane.

Metallic targets are attractive for such experiments, since their spin-lattice relaxation times (T_1) are sufficiently long at room temperature. When employing nuclear reactions, the alignment may still be lost during the stopping of the recoil nucleus or through quadrupole interactions. Liquid metals, however,

reduce these effects while still having⁶ relaxation times of the order 10^{-4} to 10^{-3} sec. Liquid gallium metal was recently used by Christiansen et al. in their stroboscopic measurement of the magnetic moment of the 5- μ sec isomer of Ge⁶⁹.⁷ Similarly our experiments showed that As⁷³ in liquid gallium would be a suitable system to attempt NMR/RD.

Irradiation with 30-MeV α particles produces the 426-keV, $I^\pi = 9/2^+$ isomer of As⁷³ from Ga⁷¹ ($I^\pi = 3/2^-$) via the $(\alpha, 2n)$ reaction. A 360-keV (M2) and a 66-keV (M1) γ ray are emitted in the $(9/2^+) \rightarrow (5/2^-) \rightarrow (3/2^-)$ cascade depopulating this state. The angular distribution will be described by the expansion

$$W(\theta) = 1 + \sum A_\nu P_\nu(\cos\theta), \quad \nu = 2, 4 \quad (1)$$

If one assumes the highest alignment possible in this experiment for the $9/2^+$ state (population of the $m = \pm 1/2, \pm 3/2$ substates only, and equal distribution among these), and furthermore transitions of pure multipolarity (see⁸), then: $A_2(360) = +0.417$; $A_4(360) = -0.167$ $A_2(66) = -0.292$; $A_4(66) = 0$. Earlier measurements⁹ had given $A_2(360) = +0.34 \pm 0.04$; $A_4(360) = -0.05 \pm 0.05$; $A_2(66) = -0.20 \pm 0.05$, while $A_4(66)$ can be assumed to be small. Therefore, only the P_2 terms will be considered in the following analysis.

The experimental geometry can be described by the relative orientations of four vectors; i.e. the beam direction \vec{k}_1 , the propagation direction of the observed γ ray \vec{k}_2 , and the directions of the two magnetic fields acting at the nucleus: the dc field \vec{H}_0 and the rf field $\vec{H}_1'(t) = 2\vec{H}_1 \cos\omega t$. The geometry employed is illustrated in the insert in Fig. 1. Taking the z-direction along the beam, \vec{H}_0 was also along z, \vec{H}_1 along $\pm x$, and the two detectors were in the $z(\theta = 0^\circ)$ and $y(\theta = 90^\circ)$ directions. The measurements were

essentially observations of the time-integral correlations, where the normalized γ -ray intensities are¹⁰

$$W(\theta) = 1 + A_2 \overline{G_2} P_2(\cos\theta) \quad (2)$$

with

$$\overline{G_2} = (1/4) (3 \cos^2\beta - 1)^2 + 3 \sin^2\beta \cos^2\beta / (1 + \omega_e \tau)^2 + (3/4) \sin^4\beta / (1 + (2 \omega_e \tau)^2) \quad (3)$$

$$\tan \beta = H_1 / (H_0 - \hbar\omega/g\mu_n) \quad (4)$$

$$\hbar\omega_e = \mu_n g \sqrt{(H_0 - \hbar\omega/g\mu_n)^2 + H_1^2} \quad (5)$$

The g -factor has been measured earlier⁹ as $g = +1.1 \pm 0.2$, based on⁸

$T_{1/2} = \tau \cdot \ln 2 + (5.8 \pm 0.5) \cdot 10^{-6}$ s. To produce a sizeable decrease of $\overline{G_2}$ the quantity $\omega_1 \tau = g \mu_n H_1 \tau / \hbar$ must be of order unity or greater. In this experiment we reached $\omega_1 \tau = 0.7$ with the available rf field strength, which was measured in situ with a test coil as $2H_1 = (31 \pm 3)$ gauss. A pair of coils provided a dc magnetic field up to 1500 gauss, with homogeneity and stability of about one part in 10^3 .

The target was liquid Ga^{71} metal, 0.15 mm thick and 3 mm wide, confined between two thin quartz plates. The skin depth at 1 MHz is 0.16 mm. The experiments were performed by stepping H_0 , with $H_1(t)$ held at constant amplitude and frequency (stable to ± 0.3 kHz). Of the pulses from two Ge(Li) detectors (1.5 keV FWHM at 122 keV) only those "delayed" events were accepted which occurred in a 0.14 μ sec interval between beam bursts (0.16 μ sec separation). The timing circuitry was similar to that described elsewhere.^{11,12} Four single-channel analyzers selected the 66-keV and 360-keV lines. The ratio of peak to background was about 10:1. The output pulses were registered in a 400-channel analyzer modified to operate as four 100-channel multiscalers which were

stepped synchronously with H_0 . Data were taken at two frequencies and four values of H_1 , sweeping H_0 by between $\pm 30\%$ and $\pm 10\%$. All runs were consistent with the conclusions discussed below. The data are presented as ratios of counts, e.g. $N(66\text{keV}, 0^\circ) / N(360\text{keV}, 0^\circ)$, to correct for beam fluctuations. Figure 1 shows representative curves.

Results from four measurements are listed in Table 1. Each measurement yielded three independent ratios at each of the values of H_0 , with H_1 and ω constant. A single seven-parameter least-squares fit was made to the combined three sets of ratios from each measurement, using the appropriate functions derived from Eqs. (2)-(5). The fits were satisfactory in each case. We note the following:

1) The results for the g -factor agree within $\pm 0.3\%$. This spread is probably due mainly to the error in the field determinations which was independently estimated as $\pm 0.5\%$. Including the Knight shift constant K explicitly, the weighted average of our results is $g \cdot (1 + K) = 1.1495 \pm 0.0057$. For As in liquid As, $K = +0.00318$.¹³ Since we estimated that As in Ga has only a slightly different Knight shift, we use as a correction $K = (3.2 \pm 1.0) \cdot 10^{-3}$. Thus

$$g = 1.146 \pm 0.007 \quad (6)$$

The large g -factor of $+1.146$ identifies the 426-keV level of As⁷³ quite unambiguously as a $g_{9/2}$ proton state. The ground states in seven odd proton nuclei in the $40 < Z < 50$ region are also classified as $g_{9/2}$ proton levels, with g -factors ranging somewhat higher, between $+1.23$ and $+1.37$. This is the first nucleus with $Z < 40$ in which this state has been identified by a magnetic moment measurement.

2) The size of the derived values of A_2 is approximately one-half that observed earlier. The neglect of $A_4(360)$ is not expected to be the reason for this.

However, the possibility of insufficient heating of the target cannot be completely ruled out. Quadrupole interactions in solid gallium (orthorhombic, melting point 29.8°C) or possibly in the liquid when near freezing would destroy the coherent motion of the nuclear spins. If one tries to describe the reduction of A_2 phenomenologically by introducing an effective lifetime τ_{eff} which is shorter than the nuclear mean lifetime $\tau = 8.37 \cdot 10^{-6}$ sec, then $\tau_{\text{eff}} \approx (6.5 \pm 0.5) \cdot 10^{-6}$ sec can explain the observed data.

3) The observed lines are distinctly non-Lorentzian. In fact the theoretical lineshapes described by Eqs. (2)-(5) have two maxima and resemble two unresolved Lorentzians split by $\sqrt{2} H_1$ when $\omega_1 \tau \geq 1$ (saturation).¹⁰ This shape is expected for experiments in which the observed quantity transforms as a second-rank tensor,¹⁴ but to our knowledge it has not been previously demonstrated in a nuclear magnetic resonance experiment.

We are indebted to E. Matthias and C. M. Lederer for their help in various stages of this experiment. Much of the experimental setup was developed by G. Gabor, W. Gagnon, E. Lampo, and D. Voronin. The kind cooperation of the 88-inch cyclotron group is gratefully acknowledged.

FOOTNOTES AND REFERENCES

* Work performed under the auspices of the U. S. Atomic Energy Commission.

† On leave of absence from Institut für Technische Kernphysik, Technische Hochschule, Darmstadt.

1. Martin Deutsch and Sanborn C. Brown, Phys. Rev. 85, 1047 (1952).
2. N. Bloembergen and G. R. Temmer, Phys. Rev. 89, 883 (1953).

3. A. Abragam and R. V. Pound, Phys. Rev. 92, 943 (1953).
4. See, e.g., "Hyperfine Interactions and Nuclear Radiations," Ed. by E. Matthias and D. A. Shirley, (North-Holland Publishing Co., Amsterdam, 1968) Chapter X.
5. Donald Connor, Phys. Rev. Letters 3, 429 (1959); Donald Connor and Tung Tsang, Phys. Rev. 126, 1506 (1962); Tung Tsang and Donald Connor, Phys. Rev. 132, 1141 (1963); K. Sugimoto, A. Mizobuchi, K. Nakai, and K. Matuda, Phys. Letters 18, 38 (1965); J. Phys. Soc. Japan 21, 213 (1966); K. Sugimoto, Ref. 4, page 859; K. Sugimoto, K. Nakai, K. Matuda, and T. Minamisono, Phys. Letters 25B, 130 (1967).
6. D. A. Cornell, Phys. Rev. 153, 208 (1967); Note that the spin-lattice relaxation time applicable in the present experiment is about 1/3 of the T_1 observed in conventional NMR.
7. J. Christiansen, H.-E. Mahnke, E. Recknagel, D. Riegel, G. Weyer, and W. Witthun, Phys. Rev. Letters 21, 554 (1968).
8. R. W. Hayward and D. D. Hoppes, Phys. Rev. 101, 93 (1956); H. H. Bolotin, Phys. Rev. 131, 774 (1963); E. Bodenstedt, G. Strube, W. Engels, H. Blumberg, R.-M. Lieder, and E. Gerdau, Phys. Letters 6, 290 (1963).
9. D. Quitmann and J. M. Jaklevic, Bull. Am. Phys. Soc. 14, 624 (1969).
10. B. Olsen, E. Matthias, and R. M. Steffen, UUIP-583, Institute of Physics University of Uppsala (April 1968). See also Ref. 4. We write here $\frac{G}{G_2}$ for G_{22}^{00} .
11. T. Yamazaki and G. T. Ewan, Nucl. Instr. Methods 62, 101 (1968).
12. J. M. Jaklevic, C. M. Lederer, and J. M. Hollander (to be published).
13. D. A. Rigney and J. A. Blodgett, to appear in J. Phys. Chem. Sol. (1969).

14. M. Guichon, J. E. Blamont, and J. Brosset, *Compt. Rend.* 243, 1859 (1956); *J. Phys. Radium* 18, 99 (1957). See also A. Kastler, *Physics Today*, September 1967, p. 34.

FIGURE CAPTION

Fig. 1. Effect of NMR on the intensity ratios for the two γ -rays which depopulate the As^{73} isomer ($9/2^+$, 426 keV) produced by $\text{Ga}^{71}(\alpha, 2n)$. The ordinates at the left give the observed ratios. The scales at the right show the change in each ratio ($\epsilon = 1$ corresponds to $\overline{G_2} = 0$); to within a few percent, $\epsilon = 1 - \overline{G_2}$ holds. Given with each curve are the rf frequency $\omega/2\pi$, the field strength in the rotating frame H_1 , and the identification of the ratio. H_0 is the applied dc field. Statistical errors are shown. The curves drawn are from the least-squares fits. The insert shows the relevant vectors.

Table 1. Summary of results. The errors given parenthetically are statistical errors only

Experimental conditions		Relative effect ^{a)}			Results from fits				
$\omega/2\pi$ MHz	H_1 gauss	$\frac{N(66,0)}{N(360,0)}$	$\frac{N(66,90)}{N(360,90)}$	$\frac{N(66,0)}{N(66,90)}$	$-A_2(66)$ b)	$+A_2(360)$ b)	χ^2	g-factor	
0.9252	14.6(7)	1.123(9) ^{e)}	0.933(7)	1.084(5)	0.136(7)	0.197(14)	0.95	1.1520(10)	c)
0.8535	17.0(7)	1.139(12)	0.941(12)	1.094(6) ^{e)}	0.146(5)	0.193(13)	1.23	1.1467(10)	c)
0.8562	12.4	1.074(15)	0.932(15)	1.050(10) ^{e)}	0.114(8)	0.174(17)	1.19	1.1496(20)	a)
0.8581	8.5	1.056(20)	0.968(20)	1.032(15) ^{e)}	0.126(12)	0.184(22)	0.90	1.1531(30)	a)

a) Ratio (on resonance, i.e., averaged over about ± 25 gauss)/ratio (off resonance).

b) The four corrections for background in the energy windows were applied in the fits.

c) The value of H_1 is from the least-squares fit. The H_1 field had been increased for the second measurement by about 10%.

d) The relative values of H_1 had been directly measured for the last three entries. Therefore, H_1 was fixed correspondingly in the last two fits. Also, the ratio $A_2(66)/A_2(360)$ was restricted to be -0.7 ± 0.1 .

e) Shown in Fig. 1.




Fig. 1.

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or*
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.*

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

TECHNICAL INFORMATION DIVISION
LAWRENCE RADIATION LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720