Submitted to Physical Review Letters

UCRL-18930
Preprint

cy. 2

EVIDENCE FOR Y_4 DEFORMATION IN ^{20}Ne AND OTHER s-d SHELL NUCLEI

RECEIVED
LAWRENCE
RADIATION LABORATORY
JUL 14 1969
LIBRARY AND
DOCUMENTS SECTION

R. de Swiniarski, C. Glashausser, D. L. Hendrie,
J. Sherman, A. D. Bacher, and E. A. McClatchie

June 1969

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

LAWRENCE RADIATION LABORATORY
UNIVERSITY of CALIFORNIA BERKELEY

UCRL-18930
cy. 2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

EVIDENCE FOR Y_4 DEFORMATION IN ^{20}Ne AND OTHER s-d SHELL NUCLEI*

R. de Swiniarski,† C. Glashausser, D. L. Hendrie, J. Sherman, A. D. Bacher, and E. A. McClatchie

Lawrence Radiation Laboratory
University of California
Berkeley, California 94720

June 1969

The sign and magnitude of the hexadecapole deformation in ^{20}Ne is determined from measurements of the inelastic scattering of 24.5-MeV protons. A coupled-channels analysis yields a value for β_4 of +0.28. A similar analysis of other available data in the s-d shell suggests a large hexadecapole deformation in ^{28}Si(+0.25) and ^{32}S(+0.25); ^{24}Mg is found to have a Y_4 moment close to zero.

Accurate measurements of the large intrinsic quadrupole deformation of the first excited 2^+ states in ^{20}Ne, ^{24}Mg, and ^{28}Si have recently been performed.[1] Such data are a critical test of the detailed microscopic calculations of nuclear properties which are now being carried out by methods such as deformed Hartree-Fock.[2] Some of these calculations suggest that nuclei of the 2s-1d shell should also have a ground-state hexadecapole deformation which changes both in size and sign through the shell.[3,4,5] The size and sign of Y_4 moments of rare-earth nuclei have previously been determined by a coupled-channels analysis of the cross sections for excitation of the ground-state rotational band by 50 MeV alpha particles.[6] The advantage of this method was that all multiple-excitation paths between these states were treated

consistently. A similar analysis of scattering data in the 2s-1d shell has been performed only for ^{24}Mg, but no Y_4 deformation was observed in the ground state band.[7] On the other hand, previous inelastic scattering results analyzed with DWBA, Austern-Blair, and other less sophisticated models indicate that large direct transition strengths are needed in order to explain the magnitude of the cross sections for the first 4^+ states in ^{20}Ne and ^{28}Si.[8,9,10]

We have measured the inelastic scattering of 24.5 MeV protons from ^{20}Ne. The cross sections for the lowest 0^+, 2^+, 4^+ and 6^+ states are analyzed with the same coupled-channels method used for the rare-earth nuclei. The same model is then used to analyze the data of Crawley and Garvey[11] for inelastic scattering of 17.5 MeV protons from ^{24}Mg, ^{28}Si, and ^{32}S. Evidence is found for large positive hexadecapole deformation in ^{20}Ne, ^{28}Si, and ^{32}S while ^{24}Mg is determined to have a small, possibly negative, hexadecapole moment. These results are relatively independent of the size and sign of β_2.

The ^{20}Ne experiment was performed at the Berkeley 88" cyclotron. A 24.5 MeV proton beam was scattered from a gas target filled to 10 or 20 cm Hg pressure with enriched ^{20}Ne. A set of four Si(Li) detectors were used to count the scattered particles; the detectors were 4 mm thick and cooled to -25°C. The total energy resolution achieved was about 50 keV. The angular distributions are shown in Fig. 1, along with theoretical curves described below.

In the coupled-channels calculations the states explicitly coupled are assumed to be the lowest members of a pure K=0 rotational band. The intrinsic deformation of these states is parameterized according to the following definition of the nuclear radius:

$$R(\theta) = R_0 \left[1 + \beta_2 Y_{20}(\theta) + \beta_4 Y_{40}(\theta) \right] \quad .$$

The interaction potential arises from the deformation of both the real and imaginary central terms of the optical potential and is calculated correctly to all orders. Thus, all possible multiple excitation paths between the coupled states are explicitly included. Coulomb excitation and deformed spin-orbit terms are included in many of the calculations but are found to have no significant effect. The coupled-channels code of A. D. Hill, which includes a spin-orbit term in the optical potential, is used for most of the calculations. The predictions for the 6^+ state in ^{20}Ne are made with the program of N. K. Glendenning, which, however, does not include a spin-orbit term. The 2^+ and 4^+ curves are insensitive to the spin-orbit potential.

Starting optical-model parameters were obtained by fitting only the elastic cross sections using the search code MERCY. Very good fits to the elastic cross sections were obtained with several distinct sets of optical model parameters. These parameters were obtained either by searching only on the well depths using the geometrical parameters recently found for nuclei in the $1f_{7/2}$ shell[12] or by searching on all parameters. These parameters were then adjusted to preserve the fits to the elastic scattering in the coupled-channel calculations. Usually it was sufficient to adjust only W_D, a_I, and V_o. The parameters used for the curves shown in Figs. 1-2 are given in Table I.

The experimental cross sections and theoretical curves for the 0^+, 2^+, 4^+ and 6^+ states in ^{20}Ne are shown in Fig. 1. With values of +0.47 for β_2 and +0.28 for β_4, good fits are obtained to the shape and magnitude of the 0^+, 2^+,

and 4^+ cross sections; the 6^+ cross section is underestimated by a factor of about 2. The sensitivity of the predictions for the 4^+ and 6^+ states to the value of β_4 is also illustrated in this figure. When β_4 is omitted, the predicted 4^+ and 6^+ cross sections are too small by one or two orders of magnitude. If β_4 is negative, the predicted shape of the 4^+ angular distribution does not match the experimental curve. In fact, changing β_4 by +0.05 or -0.05 is sufficient to destroy the good agreement with the 4^+ cross section. Only a large positive Y_4 deformation could reproduce the shape as well as the magnitude of the 4^+ cross section. Changing β_2 has a smaller effect on the predicted cross sections. If it is omitted (as it is, e.g., in a DWBA calculation), the value of β_4 must be increased to about 0.36 and the fit deteriorates somewhat, especially at back angles.

The coupled-channels predictions for the 0^+, 2^+, and 4^+ states in ^{24}Mg, ^{28}Si, and ^{32}S are shown in Fig. 2; no 6^+ data were available. The values of β_2 and β_4 are given in Table I. The signs of the β_2 deformations in ^{24}Mg and ^{28}Si were chosen to agree with the results of Ref. 1. Hartree-Fock calculations[5] predict an oblate deformation for ^{32}S, but this has not yet been verified experimentally and is not determined by the present analysis. The sign and magnitude of β_4 for ^{32}S are not very sensitive to the sign of β_2. The fits to the elastic scattering are good for all three nuclei; the striking difference in the shapes of the 4^+ angular distributions for ^{24}Mg and ^{28}Si - ^{32}S is also qualitatively explained. However, the general quality of the fits shown in Fig. 2 is inferior to that obtained for the 0^+, 2^+, and 4^+ states in ^{20}Ne. A conservative error of ±0.08 is thus assigned to the value of β_4 determined for these nuclei, due to ambiguities in the optical parameters and imperfections of the fit. The corresponding error for ^{20}Ne is ±0.05.

Non-direct processes may be responsible for some of the discrepancies, especially for the 6^+ state in ^{20}Ne and for the 4^+ state in ^{24}Mg. However, there is evidence that such processes are not important at forward angles for the 4^+ states with larger cross sections. Prior to this experiment, an excitation function for the 0^+, 2^+, and 4^+ states in ^{20}Ne was measured at proton energies between 23 and 26 MeV in 500 keV steps. The excitation functions of these states are smooth between 23 and 26 MeV. In addition the 4^+ cross section decreases by more than an order of magnitude between forward and backward angles. The cross section for the 4^+ state in ^{28}Si measured at 21.2 MeV[10] is similar to the 17.5 MeV cross section.[11]

An additional source of ambiguity is the known imperfection in the rotational description of these nuclei. The energy levels, particularly in ^{32}S, already show considerable deviation from the strict rotational model pattern. The values of B(E2) for the intraband transitions measured in various laboratories are not always consistent.[13] However, the ratio $[B(E2)4^+ \longrightarrow 2^+]/[B(E2)2^+ \longrightarrow 0^+]$ seems to be generally slightly less than 1.0, whereas the rotational model prediction is 1.43. For ^{32}S, however, a recent measurement[14] gives 2.6 ± 0.7. This means that the multiple excitation contributions to the 4^+ cross sections are somewhat overestimated except in ^{32}S. The $[B(E2)6^+ \longrightarrow 4^+]$ in ^{20}Ne seems to be considerably larger than the measured $[B(E2)2^+ \longrightarrow 0^+]$ in this nucleus, and is larger than expected from the rotational model. This may be another reason why the predicted 6^+ cross section in ^{20}Ne is too small.

In terms of the rotational model, non-zero values of β_4 imply a hexadecapole moment in the ground state and in all the states of the rotational

band built upon it. However, the inelastic scattering data alone might be equally well described by a vibrational model, with some modifications of the values of β_2 and β_4. Thus, the interpretation of β_4 as describing the static Y_4 deformation of the ground state band relies upon measurements of a non-zero quadrupole moment; such a measurement has not yet been made for ^{32}S. The quadrupole moments of the 2^+ states of ^{20}Ne and ^{24}Mg are about 30% larger than expected[1] on the basis of the rotational model from the electromagnetic values of $[B(E2)2^+ \longrightarrow 0^+]$. Since the analysis of the inelastic scattering depends upon the evaluation of matrix elements between the ground state and excited states, the present results should be interpreted in terms of a transition probability instead of a static moment when the two are not consistent.

Benson and Flowers[4] have predicted a value of about 0.17 for β_4 in ^{20}Ne; Hartree-Fock calculations[5] predict a large Y_4 moment for ^{28}Si but a small Y_4 moment for ^{32}S. We have made an estimate of the value of β_4 expected in the Nilsson model, according to the simple method of Harada[15] which accurately predicted the relative values for rare-earth nuclei. The predictions follow the general trend of the experimental values except for ^{32}S which is again underestimated.

To summarize, the coupled-channels analysis of the present ^{20}Ne data show clearly the existence of a large hexadecapole deformation. A similar analysis of available data suggests a large hexadecapole deformation also in ^{28}Si and ^{32}S while ^{24}Mg is found to have a very small Y_4 deformation.

We are very grateful to Drs. N. K. Glendenning and A. D. Hill for the use of their coupled-channels programs and to N. Brown for performing some of the calculations presented here.

Footnotes and References

*Work supported in part by the U.S. Atomic Energy Commission.

†NATO-Fulbright fellow; permanent address: Institut des Sciences Nucleaires de Grenoble, France.

1. A. Bamberger, P. G. Bizzeti, and B. Povh, Phys. Rev. Letters 21, 1599 (1968); O. Häusser, B. W. Hooton, D. Pelte, T. K. Alexander, and H. C. Evans, Phys. Rev. Letters 22, 359 (1969); D. Schwalm and B. Povh, Phys. Letters 29B, 103 (1969); T. K. Alexander, D. Pelte, O. Häusser, B. Hooton, and H. C. Evans, Bull. Am. Phys. Soc. 14, 555 (1969); K. Nakai, F. Stephens, and R. Diamond, private communication.
2. S. Das Gupta and M. Harvey, Nucl. Phys. A94, 602 (1967); G. Ripka, Advances in Nuclear Physics, M. Baranger and E. Vogt, editors, (Plenum Press, New York, 1968).
3. C. Brihaye and G. Reidemeister, Nucl. Phys. A100, 65 (1967).
4. H. G. Benson and B. H. Flowers, Nucl. Phys. A126, 305 (1969).
5. B. Castel and J. C. Parikh, to be published.
6. D. L. Hendrie, N. K. Glendenning, B. G. Harvey, O. N. Jarvis, H. H. Duhm, J. Saudinos, and J. Mahoney, Phys. Letters 26B, 127 (1968).
7. M. Reed, thesis, Univ. of Calif., (1968), University of California Lawrence Radiation Laboratory Report UCRL-18414, (unpublished); A. A. Rush and N. K. Ganguly, Nucl. Phys. A117, 101 (1968).
8. J. S. Blair, N. Cue, G. W. Farwell, and D. Shreve, Univ. of Washington Annual Progress Report, (1964) (unpublished); A. Springer, thesis, Univ. of Calif., (1965), University of California Lawrence Radiation Laboratory Report UCRL-18681 (unpublished).

9. F. Hinterberger, G. Mairle, V. Schmidt-Rohr, G. J. Wagner, and P. Turek, Nucl. Phys. A115, 570 (1968).

10. R. W. Barnard and G. D. Jones, Nucl. Phys. A108, 655 (1968).

11. G. M. Crawley and G. T. Garvey, Phys. Rev. 160, 981 (1967); G. M. Crawley, thesis, Princeton Univ., (1965), (unpublished).

12. P. Kossanyi-Demay and R. de Swiniarski, Nucl. Phys. A108, 577 (1968).

13. S. J. Skorka, J. Hertel, and T. W. Retz-Schmidt, Nuclear Data 2, 347 (1966); J. H. Anderson and R. C. Ritter, Nucl. Phys. A128, 305 (1969).

14. G. T. Garvey, K. W. Jones, L. E. Carlson, A. G. Robertson, and D. F. H. Start, Phys. Letters 29B, 108 (1969).

15. K. Harada, Phys. Letters 10, 80 (1964).

Figure Captions

Fig. 1. Measured cross-sections and theoretical predictions for the inelastic scattering of 24.5-MeV protons from ^{20}Ne. The curves were calculated with a coupled-channels program; a rotational model was used with the values of β_2 and β_4 indicated.

Fig. 2. Experimental cross sections of Crawley and Garvey[11] for the inelastic scattering of 17.5-MeV protons from ^{24}Mg, ^{28}Si, and ^{32}S. The curves are coupled-channels predictions with the values of β_2 and β_4 indicated in Table I.

Table I.

	V_o (MeV)	r_o (F)	a_o (F)	W_D (MeV)	r_I (F)	a_I (F)	V_s (MeV)	r_s (F)	a_s (F)	β_2 (exp)	β_4 (exp)
^{20}Ne	54.4	1.05	0.73	6.30	1.26	0.55	3.57	0.95	0.33	+0.47	+0.28
^{24}Mg	46.0	1.22	0.60	3.60	1.27	0.64	7.26	1.22	0.60	+0.47	-0.05
^{28}Si	46.0	1.24	0.62	8.0	1.19	0.40	6.0	1.24	0.62	-0.34	+0.25
^{32}S	47.0	1.21	0.62	9.5	1.26	0.28	6.0	1.21	0.62	-0.30	+0.25


^{20}Ne (p,p′) ^{20}Ne*
E_p = 24.5 MeV
• Experiment
β_2=0.47 β_4=0
β_2=0.47 β_4=-0.10
β_2=0.47 β_4=0.28
0.00 MeV (0+)
1.632 MeV (2+)
4.250 MeV (4+)
8.790 MeV (6+)
Cross section (mb/sr)
$\theta_{c.m.}$ (deg)
XBL696-2984

Fig. 1.


^{24}Mg (p,p') ^{24}Mg*
E_p = 17.5 MeV
0.0 MeV (0+)
1.37 MeV (2+)
4.12 MeV (4+)
^{28}Si (p,p')^{28}Si*
E_p = 17.5 MeV
0.0 MeV (0+)
1.78 MeV (2+)
4.61 MeV (4+)
^{32}S (p,p')^{32}S*
E_p = 17.5 MeV
0.0 MeV (0+)
2.23 MeV (2+)
4.46 MeV (4+)
Cross section (mb/sr)
$\theta_{c.m.}$ (deg)
XBL696-2985

Fig. 2.

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

TECHNICAL INFORMATION DIVISION
LAWRENCE RADIATION LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720