Submitted to Inorganic Chemistry

UCRL-19094
Preprint

cy. I

REACTIONS OF DISULFUR DINITRIDE WITH ANTIMONY PENTACHLORIDE

RECEIVED
LAWRENCE
RADIATION LABORATORY
NOV 6 1969
LIBRARY AND
DOCUMENTS SECTION

R. Lyle Patton and William L. Jolly

October 1969

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545

LAWRENCE RADIATION LABORATORY
UNIVERSITY of CALIFORNIA BERKELEY

UCRL-19094
cy. 2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Contribution from the Department of Chemistry of the
University of California and the Inorganic Materials Research Division of
the Lawrence Radiation Laboratory, Berkeley, California 94720

Reactions of Disulfur Dinitride with Antimony Pentachloride

By R. Lyle Patton and William L. Jolly

ABSTRACT

Solutions of S_2N_2 in dichloromethane react with antimony pentachloride (in excess) to form a diadduct $S_2N_2(SbCl_5)_2$ which can further react with S_2N_2 to form a monoadduct $S_2N_2SbCl_5$. The monoadduct can be reconverted to the diadduct by treatment with $SbCl_5$. The physical and chemical properties of these compounds indicate that the S_2N_2 ring structure is maintained intact. The monoadduct $S_2N_2SbCl_5$ reacts irreversibly with S_2N_2 to form both the previously characterized $S_4N_4SbCl_5$ and, in lower yields, a less reactive material $(S_4N_4SbCl_5)_x$. Antimony pentachloride acts as a catalyst for the dimerization of S_2N_2.

Tetrasulfur tetranitride reacts with many Lewis acids such as antimony pentachloride,[1] boron trichloride[2] and boron trifluoride[2] to form 1:1 adducts in which the Lewis acid is coordinated to a nitrogen atom of S_4N_4. Although a change in the configuration of the S_4N_4 results, the basic 8-membered ring structure remains intact.[3,4] Of the dozen or so known adducts, only two have two molecules of Lewis acid associated with each molecule of S_4N_4. These are $S_4N_4(SO_3)_2$ [5] and $BCl_3 \cdot S_4N_4 \cdot SbCl_5$ [2] (which may have the ionic structure $[BCl_2S_4N_4]^+ [SbCl_6]^-$). Apparently when one of the nitrogen atoms of S_4N_4 is coordinated, the donor ability of the remaining nitrogens is markedly decreased. In this investigation, we have studied the reactions of disulfur dinitride, a planar four-membered ring with alternating S-N atoms,[6] with antimony pentachloride. Of particular interest to us was the ability of the S_2N_2 nitrogens to act as donor atoms and the effect of this donation on the stability and structure of the S_2N_2 ring.

Experimental Section

General.- The moisture-sensitivity of the materials required their manipulation in a vacuum line or in a polyethylene glove bag flushed with nitrogen or argon. Infrared spectra were recorded in the 400-4000 cm^{-1} range with Perkin Elmer Infracord spectrometers (Models 137 and 137B). Solid samples were prepared as Nujol mulls pressed between KBr plates; solution spectra were run in a 0.1-mm KBr cavity cell.

Melting points were obtained in argon-filled, sealed capillaries. Analyses for N, S, and Cl were performed in the microanalytical laboratory of this department by V. Tashinian.

Antimony pentachloride (J. T. Baker Co.) was purified by vacuum-line fractional condensation and was found to melt at 3° (literature,[7] 2.8°). Dichloromethane was refluxed over P_2O_5 for 6-10 hrs., distilled, and stored in a nitrogen atmosphere. Disulfur dinitride was prepared by subliming S_4N_4 through a plug of silver wool at 300° using a technique like that described by Becke-Goehring.[8] It was found that the best results were obtained using a 0.5-g. plug of silver wool in a 10-mm. i.d. glass tube. Initially almost all the silver reacted with S_4N_4 to form silver sulfide and nitrogen; the silver sulfide then acted as a catalyst for the conversion of S_4N_4 to S_2N_2 and small amounts of $(SN)_x$. The S_2N_2 was identified by its infrared spectrum[6] and its molecular weight in dichloromethane (observed, 93; calcd. 92.1). The material was observed to slowly polymerize to $(SN)_x$ and small amounts of S_4N_4 at room temperature.[9] A sample of S_2N_2 (0.059 g, 0.646 mmole) was held in a sealed tube at 250° for 1 hr, whereupon it decomposed to sulfur and 0.646 mmole of nitrogen.

The $SbCl_5$ and S_2N_2 were stored in tared bulbs equipped with stopcocks lubricated with Kel-F wax. Disulfur dinitride was measured out from its tared storage bulb by room-temperature vacuum sublimation into a reaction vessel held at -196°. This method of transfer also served as a purification method for S_2N_2, because the common contaminants, $(SN)_x$ and S_4N_4, are nonvolatile at room temperature. Enough CH_2Cl_2 was distilled into the reaction vessel to dissolve the S_2N_2 when warmed to 20°; then the solution was transferred to a glove bag. The formation of flocculent blue-black particles of $(SN)_x$ (always less than 1 mg) on dissolution necessitated filtration of these S_2N_2 solutions immediately before use. Because S_2N_2 polymerizes appreciably in the solid state at 20° and is reportedly

shock-sensitive, direct handling of the solid in the glove bag was avoided. Solutions of known amounts of $SbCl_5$ were prepared in a similar manner. Products were normally characterized by their infrared spectra and physical properties.

Formation of $S_4N_4SbCl_5$.- Antimony pentachloride (4.7 mmoles) in 10 ml of CH_2Cl_2 was slowly dripped into a solution of S_2N_2 (3.16 mmoles) in 40 ml of CH_2Cl_2. The reaction mixture immediately became red-black and opaque, and remained so throughout the $SbCl_5$ addition. After 1/2 hour at 20°, the suspension slowly cleared and lightened to a deep red solution over a dark red-brown solid. Filtration of this mixture yielded almost pure crystals of $S_4N_4SbCl_5$ as identified by its infrared spectrum[2] and melting point (observed, 160°; literature,[2] 160-2°). Evaporation of the solution to dryness also resulted in $S_4N_4SbCl_5$. The total yield was 93%.

Formation of $S_2N_2(SbCl_5)_2$.- The dropwise addition at 20° of a solution of S_2N_2 (1.72 mmoles) in 25 ml of CH_2Cl_2 to $SbCl_5$ (7.8 mmoles) in 10 ml of CH_2Cl_2 resulted in the immediate formation of fine yellow-white crystals which increased in quantity with continued S_2N_2 addition. Filtration of the mixture yielded 272 mg of extremely moisture-sensitive yellow-white crystals which quickly turned yellow, then black in moist air. Anal. Calcd. for $S_2N_2(SbCl_5)_2$: N, 4.06; S, 9.29; Cl, 51.37. Found: N, 3.96; S, 9.17; Cl, 51.36.

Cooling the filtrate to -15° yielded a further 522 mg of $S_2N_2(SbCl_5)_2$; the total yield was 67%. Evaporation of the final filtrate to dryness *in vacuo* produced only this diadduct; no $S_4N_4SbCl_5$, as observed when the reactants were mixed in the reverse order, was found. The diadduct is also formed (in 95-100% yields) when a frozen mixture of $SbCl_5$, S_2N_2 and

CH_2Cl_2 is slowly warmed to and held at 20°, followed by vacuum removal of the solvent and excess $SbCl_5$. The glove bag preparation, however, is more convenient.

The solubility of $S_2N_2(SbCl_5)$ in dichloromethane is ca. 1.5 g/100 ml at 20°, and it may be readily recrystallized by cooling a saturated solution. No decomposition on long standing in solution was detected, provided moisture was carefully excluded. The crystals slowly become opaque and red-black above 120°, before melting at 130-131° with decomposition. The infrared spectrum of $S_2N_2(SbCl_5)_2$ shows absorptions (in cm^{-1}) at 461 (s), 818 (s), and 898 (vw).

Formation of $S_2N_2SbCl_5$.- Disulfur dinitride (1.95 mmoles) in 30 ml of CH_2Cl_2 was slowly dripped into a solution of $SbCl_5$ (2.0 mmoles) in 10 ml of CH_2Cl_2. After addition of 4 ml of S_2N_2 solution, fine yellow-white crystals appeared which increased in quantity until half of the S_2N_2 solution had been added. As the remaining S_2N_2 was added, the crystals slowly dissolved, and the solution changed from light yellow to light orange in color. Complete addition of the S_2N_2 resulted in complete disappearance of the crystals. Concentration of this solution to 10 ml by distilling off CH_2Cl_2 *in vacuo* followed by cooling to -15° yielded long (up to 1 cm), clear, light orange needless of $S_2N_2SbCl_5$, of which 100 mg was filtered off. Further concentration and filtration, followed by final evaporation to dryness, yielded a total of 700 mg of this product which contained slight $S_2N_2(SbCl_5)_2$ contamination. Anal. Calcd. for $S_2N_2SbCl_5$: N, 7.16; S, 16.39; Cl, 45.32; mol wt, 391.2. Found: N, 7.26; S, 16.08; Cl, 45.38, mol wt (by vapor pressure lowering in CH_2Cl_2), 402.

The monoadduct $S_2N_2SbCl_5$ is very soluble in CH_2Cl_2 at 20°: ca. 10g/100 ml. These solutions may be heated briefly to 40° with no apparent decomposition. Darkening of solutions is observed, however, even at 20° on long standing. In a sealed capillary, $S_2N_2SbCl_5$ crystals turn black at 100°, then melt with decomposition at 108-109°. The crystals immediately blacken in moist air. The infrared spectrum shows the following peaks; 1005 (m), 922 (vw), 800 (s), 723 (mw), 627 (w), and 459 (s) cm^{-1}.

A sample of $S_2N_2SbCl_5$ (0.83 mmole) was placed in a vacuum sublimator with a water-cooled cold finger and heated while evacuating through a -196° trap with a Toepler pump. Above 55° a white material collected on the cold finger. After 4 hours at 70°, the products were examined. An orange-brown residue of $S_4N_4SbCl_5$ (0.12 mmole) remained in the sublimator. Nitrogen (0.008 mmole) was found in the Toepler pump buret, $SbCl_5$ (0.02 mmole) was found in the -196° trap, and a yellow-white solid was found on the cold finger. The infrared absorptions of this material corresponded to ca. 90% $S_2N_2SbCl_5$ and 10% $S_2N_2(SbCl_5)_2$. A slower sublimation at 60° produced proportionately much less $S_4N_4SbCl_5$ and $S_2N_2(SbCl_5)_2$.

Pyrolysis of $S_2N_2(SbCl_5)_2$.- When $S_2N_2(SbCl_5)_2$ (0.25 mmole) was heated at 90° for 1 hour in a vacuum sublimator with a 20° cold finger, $SbCl_5$ (0.22 mmole) was evolved, and yellow-white $S_2N_2SbCl_5$ (with slight $S_2N_2(SbCl_5)_2$ contamination) collected on the cold finger. A slight residue of $S_4N_4SbCl_5$ (0.01 mmole) remained in the sublimator.

Addition of S_2N_2 to $S_2N_2SbCl_5$ and $S_2N_2(SbCl_5)_2$.- Addition of S_2N_2 to $S_2N_2SbCl_5$ solutions in equimolar amounts yielded an opaque greenish-black suspension; precipitation began with the first drops of S_2N_2 added. On standing at 20° for 1-2 hours, the suspension gradually cleared and lightened

to an orange-red solution above a dark red-orange solid. Examination of this solid and that left after evaporation of the solvent showed it to be $S_4N_4SbCl_5$ slightly contaminated with a greenish-yellow substance. The $S_4N_4SbCl_5$ was identified by its infrared spectrum and melting point of 160° (literature,[2] 160-2°) after recrystallization from CH_2Cl_2. No S_2N_2 or S_4N_4 was recovered.

Addition of S_2N_2 to $S_2N_2(SbCl_5)_2$ solutions resulted in an initial color change corresponding to formation of $S_2N_2SbCl_5$ which, with further S_2N_2, again became dark and opaque followed by slow clearing to a red-orange solution and precipitation of $S_4N_4SbCl_5$. Addition of S_2N_2 in excess of the amount required to convert all the $S_2N_2SbCl_5$ to $S_4N_4SbCl_5$ resulted in a mixture of S_4N_4 and $S_4N_4 \cdot SbCl_5$ in the relative amounts expected from conversion of the excess S_2N_2 to S_4N_4. Tetrasulfur tetranitride was recrystallized from dichloromethane and identified by its infrared spectrum[10] and melting point of 182° (literature,[11] 187-187.5°). To test the catalytic effect of $S_4N_4SbCl_5$ on the dimerization of S_2N_2, 1.02 mmoles of disulfur dinitride was slowly added to a solution of 0.15 mmole of $S_4N_4SbCl_5$. The solution remained orange-red. Infrared analysis indicated that after 2 hrs approximately equal amounts of S_4N_4 and S_2N_2 were in the solution, and that after 15 hours only 10% of the original S_2N_2 remained.

A small amount (5-10%) of the greenish-yellow substance was always formed as a by-product of these reactions in which $S_4N_4SbCl_5$ was formed. The characterization of this material is described below.

$\underline{(S_4N_4SbCl_5)_x}$.- The small amount of greenish-yellow material formed in the above reactions was found to be insoluble in CH_2Cl_2 and could be isolated free of contamination by $S_4N_4SbCl_5$ and S_4N_4 by using enough solvent

to keep all the expected $S_4N_4SbCl_5$ and S_4N_4 in solution. When the opaque suspension cleared, as much as 50 mg of this microcrystalline substance remained and was filtered off. Although the color of different samples varied from yellow to a dull green, each sample melted at 193-194° (the green samples becoming yellow at 100-110°) and exhibited the same infrared absorptions at 1160 (m), 1115 (m), 975 (vw), 725 (w), and 535 (m) cm^{-1}.

In an attempt to prepare a large quantity of this material, S_2N_2 (3.64 mmoles) in 45 ml of CH_2Cl_2 was added with a dropping funnel during 1 hr. to a rapidly stirred solution of $SbCl_5$ (1.58 mmoles) in 5 ml of CH_2Cl_2. When half of the S_2N_2 had been added, the solution became dark and opaque; it cleared to a deep red solution above an orange-yellow solid only after the addition was complete. Several overnight leachings of the solid with CH_2Cl_2 removed all the $S_4N_4SbCl_5$ and left 200 mg of yellow solid having only the above infrared absorptions and a 193-193.5° m.p. Anal. Calcd. for $(S_4N_4SbCl_5)_x$: N, 11.6; S, 26.5; Cl, 36.7; Sb, 25.2. Found: N, 11.8; S, 26.1; Cl, 37.0; Sb (by difference), 25.1.

Approximately the same amounts of S_2N_2 and $SbCl_5$ as above, each in 40 ml of CH_2Cl_2, were mixed during 1 hr in the same manner as above but with the $SbCl_5$ slowly dripped into the S_2N_2 solution. In this reaction only a 1% yield of $(S_4N_4SbCl_5)_x$ was obtained, in contrast to the 26% yield resulting from the addition performed in the opposite order.

Like $S_4N_4SbCl_5$, $(S_4N_4SbCl_5)_x$ is stable to short exposure to moist air; however, its infrared spectrum, melting point, and insolubility in CH_2Cl_2 do not correspond to $S_4N_4SbCl_5$ or to any other known compound. When heated *in vacuo*, no sublimation of $(S_4N_4SbCl_5)_x$ occurred; slow decomposition began at 162°, and after several hours at 170° N_2, S_4N_4, $SbCl_3$, NSCl, and $S_3N_2Cl_2$

were found among the decomposition products. The infrared spectrum of the remaining residue showed only $(S_4N_4SbCl_5)_x$, with no evidence of thermal rearrangement to $S_4N_4SbCl_5$. $(S_4N_4SbCl_5)_x$ was also unaffected by treatment with liquid $SbCl_5$ at 20°.

Several attempts were made to filter the opaque suspensions which formed immediately after mixing solutions of S_2N_2 and $S_2N_2SbCl_5$. In each case, regardless of order or speed of mixing, a dull green-black powder was isolated whose infrared spectrum showed only weak bands due to $(S_4N_4SbCl_5)_x$. When heated *in vacuo* above 100°, this material became orange-yellow and evolved nitrogen and sulfur. In moist air, the green color slowly changed to yellow. Neither treatment significantly changed the infrared spectrum. All these properties indicate that the isolated solid was a mixture of $(S_4N_4SbCl_5)_x$ and $(SN)_x$.

Results and Discussion

Reactions of S_2N_2 with $SbCl_5$.- The addition of disulfur dinitride to excess antimony pentachloride in dichloromethane results in rapid, complete formation of the diadduct $S_2N_2(SbCl_5)_2$. When S_2N_2 is added to an equimolar amount of $SbCl_5$, $S_2N_2(SbCl_5)_2$ precipitates during the first half of the addition. This $S_2N_2(SbCl_5)_2$ then reacts with the S_2N_2 as it is added during the second half of the addition to form the monoadduct $S_2N_2SbCl_5$. The reactions may be written:

$$S_2N_2 + 2SbCl_5 \rightarrow S_2N_2(SbCl_5)_2 \qquad (1)$$

$$S_2N_2 + S_2N_2(SbCl_5)_2 \rightarrow 2S_2N_2SbCl_5 \qquad (2)$$

Addition of S_2N_2 to an equimolar amount of $S_2N_2SbCl_5$ in dichloromethane yields the previously-characterized adduct $S_4N_4SbCl_5$.

$$S_2N_2 + S_2N_2SbCl_5 \rightarrow S_4N_4SbCl_5 \qquad (3)$$

A relatively insoluble greenish-yellow material, having the empirical formula $(S_4N_4SbCl_5)_x$ but possessing physical properties entirely different from those of ordinary $S_4N_4SbCl_5$, forms as a significant by-product of reaction 3. This incompletely characterized material is definitely worthy of further study. During the course of reaction 3, the formation of a dark-colored intermediate is apparent. Inasmuch as attempts to isolate this intermediate yielded only the $(S_4N_4SbCl_5)_x$ by-product contaminated with what appeared to be $(SN)_x$, we tentatively suggest that the intermediate is an open-chain isomer of S_4N_4 or S_2N_2, perhaps coordinated to $SbCl_5$. This intermediate may be a precursor of $(S_4N_4SbCl_5)_x$.

When S_2N_2 is added to $S_2N_2SbCl_5$ in greater than a 1:1 ratio, the excess S_2N_2 is quickly converted to S_4N_4. Obviously some species in the reaction system catalyzes the dimerization of S_2N_2. The adduct $S_4N_4SbCl_5$ does catalyze the dimerization, but relatively inefficiently. Possibly the dark intermediate discussed above is involved in the rapid dimerization.

The adduct $S_4N_4SbCl_5$ is inert toward $SbCl_5$;[2] consequently this adduct is formed whenever $SbCl_5$ is added to a solution of S_2N_2, regardless of the final S_2N_2: $SbCl_5$ ratio. The fact that $S_2N_2SbCl_5$ in dichloromethane solution does not decompose to $S_4N_4SbCl_5$ proves that the following equilibria lie far to the left.

$$S_2N_2SbCl_5 \rightleftharpoons S_2N_2 + SbCl_5 \qquad (4)$$

$$2S_2N_2SbCl_5 \rightleftharpoons S_2N_2 + S_2N_2(SbCl_5)_2 \qquad (5)$$

Otherwise $S_4N_4SbCl_5$ would form irreversibly by reaction 3. Thus we conclude that S_2N_2 is a strong base toward $SbCl_5$, and that $S_2N_2SbCl_5$ is considerably weaker base. The principle reactions observed in dichloromethane solutions are summarized in the following diagram.

$$S_2N_2 + SbCl_5 \longrightarrow S_2N_2SbCl_5 \underset{S_2N_2}{\overset{SbCl_5}{\rightleftharpoons}} S_2N_2(SbCl_5)_2$$

$$\Big\downarrow (1 + x)S_2N_2$$

$$S_4N_4 + SbCl_5 \longrightarrow S_4N_4SbCl_5 + x/2\ S_4N_4$$

The inertness of $S_4N_4SbCl_5$ toward $SbCl_5$, the interconvertibility of $S_2N_2SbCl_5$ and $S_2N_2(SbCl_5)_2$, and the observed molecular weight of $S_2N_2(SbCl_5)_2$ support the formulation of the latter compounds as adducts of S_2N_2, not S_4N_4.

Thermal Stability of S_2N_2 Adducts.- The adduct $S_2N_2SbCl_5$ sublimes *in vacuo* at 60-70° with slight decomposition corresponding to reaction 6.

$$2S_2N_2SbCl_5(s) \xrightarrow{60\text{-}70^\circ} S_4N_4SbCl_5(s) + SbCl_5(g) \qquad (6)$$

The sublimed $S_2N_2SbCl_5$ is contaminated with $S_2N_2(SbCl_5)_2$ which forms by reaction 7 on the cold finger.

$$SbCl_5(g) + S_2N_2SbCl_5(s) \xrightarrow{20^\circ} S_2N_2(SbCl_5)_2(s) \qquad (7)$$

The adduct $S_2N_2(SbCl_5)_2$ dissociates *in vacuo* at 90° via the reaction

$$S_2N_2(SbCl_5)_2(s) \longrightarrow S_2N_2SbCl_5(g) + SbCl_5(g)\ . \qquad (8)$$

The slight impurity of $S_2N_2(SbCl_5)_2$ found in the condensed $S_2N_2SbCl_5$ probably resulted from recombination of $S_2N_2SbCl_5$ and $SbCl_5$ on the cold finger rather than from direct sublimation of $S_2N_2(SbCl_5)_2$.

The low sublimation temperature of $S_2N_2SbCl_5$ supports the formulation as an S_2N_2 adduct, because an $S_4N_4(SbCl_5)_2$ adduct would be expected to be much less volatile.

Crystals of $S_2N_2SbCl_5$ are stable indefinitely at -15°, but slowly darken at 20°, becoming black after 1-2 weeks. Crystals of $S_2N_2(SbCl_5)_2$ undergo no change on standing at 20° for several months in sealed ampules after an initial surface yellowing, presumably from reaction with traces of moisture present on the glass. The stability of these adducts, compared to the appreciable polymerization of S_2N_2 above -30°, indicates that the S_2N_2 ring is stabilized by coordination with Lewis acids. The rapid reaction of $S_2N_2SbCl_5$ with S_2N_2, however, shows that this coordination somehow increases the reactivity toward attack by S_2N_2.

Infrared Spectra.- The infrared spectrum and chemical properties of S_2N_2 have led to its formulation as a four-membered planar ring with alternating S-N atoms (D_{2h} point group).[6] The prediction of three infrared-active modes agrees with the observed absorptions at 795 (s), 663 (w), and 474 (s) cm^{-1}. In the adducts, modes due to the donor $SbCl_5$ groups are expected to lie below 400 cm^{-1}. Therefore structural changes in the S_2N_2 caused by coordination should be reflected in changes in the spectrum above 400 cm^{-1}.

The two strong bands at 818 and 461 cm^{-1} in the infrared spectrum of $S_2N_2(SbCl_5)_2$ correspond closely in shape and position to the two strong S_2N_2 bands. The very weak band at 898 cm^{-1} is possibly an overtone of the band at 460 cm^{-1}. The similarity of the absorptions to those of S_2N_2 and the number of modes (2 or 3) observed indicate that the S_2N_2 ring remains

intact and planar (puckering of the ring to C_{2v} symmetry should result in five observed modes) with symmetric arrangement of the two $SbCl_5$ groups.

$S_2N_2SbCl_5$ exhibits strong absorption at 800 and 459 cm^{-1} and a weak band at 627 cm^{-1} corresponding closely in intensity, shape, and position to the S_2N_2 modes. The additional observed bands at 1005 (m), 922 (vw) (possibly an overtone of the 459 cm^{-1} band), and 723 (mw) cm^{-1} are expected from the lowering of symmetry resulting from the non-equivalence of the nitrogen atoms. Distinction between the possible configurations of the S_2N_2 group: planar ring (C_{2v}), puckered ring (C_s), or chain ($C_{\infty v}$, C_s or C_1), is not possible inasmuch as five or six infrared-active modes are predicted for each configuration. That the ring is broken (chain), however, is unlikely because of the ease of interconversion of $S_2N_2SbCl_5$ and $S_2N_2(SbCl_5)_2$, the similarity to the S_2N_2 modes, and the lack of higher frequency bands expected for terminal N-S groups. The 1005 cm^{-1} band can be reasonably assigned to an S-N ring system, because bands have been observed[2] for the S-N ring system in $S_4N_4SbCl_5$ in the same region (at 1060 and 968 cm^{-1}).

The possibility that the compounds $S_2N_2(SbCl_5)_2$ and $S_2N_2SbCl_5$ are tetra- and diadducts of S_4N_4 was considered. However, the observed bands showed no correlation with those of S_4N_4 or $S_4N_4SbCl_5$, and the forms of highest symmetry for $S_4N_4(SbCl_5)_4$ (planar ring, D_{4h}) and $S_4N_4(SbCl_5)_2$ (planar ring D_{2h}), predict four and eight observed modes, respectively. The lack of correlation in frequency and number of observed modes makes these formulations unlikely.

The infrared and chemical evidence support the formulation of $S_2N_2SbCl_5$ and $S_2N_2(SbCl_5)_2$ as adducts of the difunctional Lewis base

S_2N_2 with the structures:



(puckered or planar ring)



(planar ring) .

A recent x-ray diffraction study of $S_2N_2(SbCl_5)_2$ confirms the second of these predicted structures.[12]

Acknowledgement.- This work was supported by the United States Atomic Energy Commission.

References

(1) H. Wöbling, Z. Anorg. Allgem. Chem., 57, 281 (1908).

(2) K. J. Wynne and W. L. Jolly, Inorg. Chem., 6, 107 (1967).

(3) D. Neubauer and J. Weiss, Z. Anorg. Allgem. Chem., 303, 28 (1960).

(4) M. G. B. Drew, D. H. Templeton, and A. Zalkin, Inorg. Chem., 6, 1906 (1967).

(5) M. Goehring, H. Hohenschutz, and R. Appel, Z. Naturforsch., 9b, 678 (1954).

(6) J. R. W. Warn and D. Chapman, Spectrochim. Acta, 22, 1371 (1966).

(7) C. D. Hodgman (Editor), "Handbook of Chemistry and Physics," 44th Edition, The Chemical Rubber Publishing Co., Cleveland, Ohio, 1963, pp. 537.

(8) M. Becke-Goehring, Inorg. Syntheses, 6, 123 (1960).

(9) M. Goehring and D. Voight, Z. Anorg. Allgem. Chem., 285, 181 (1956).

(10) E. R. Lippincott and M. C. Tobin, J. Chem. Phys., 21, 1559 (1953).

(11) M. Villena-Blanco, "A Study of the Reactions Between Gaseous Ammonia and Sulfur Chlorides," M.S. Thesis, University of California, Berkeley, California, October, 1963.

(12) Unpublished work of R. L. Patton and K. N. Raymond.

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

TECHNICAL INFORMATION DIVISION
LAWRENCE RADIATION LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720