Contribution to the Conference on
Hyperon Resonances, Duke University,
Durham, N. C., April 24-25, 1970

UCRL-19774
Preprint

c. 2

THE REACTION $K^+N \rightarrow K^*N$ IN THE ISOSPIN-O CHANNEL NEAR THRESHOLD

A. A. Hirata, G. Goldhaber, V. H. Seeger,
G. H. Trilling, and C. G. Wohl

April 1970

AEC Contract No. W-7405-eng-48

RECEIVED
LAWRENCE
RADIATION LABORATORY
JUL 9 1970
LIBRARY AND
DOCUMENTS SECTION

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

UCRL-19774

LAWRENCE RADIATION LABORATORY
UNIVERSITY of CALIFORNIA BERKELEY

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Contribution to the Conference on
Hyperon Resonances, Duke University,
Durham, N.C., April 24-25, 1970

UCRL-19774

THE REACTION $K^+N \rightarrow K^*N$ IN THE ISOSPIN-0 CHANNEL NEAR THRESHOLD

A. A. Hirata, G. Goldhaber, V. H. Seeger, G. H. Trilling, and C. G. Wohl*

Department of Physics and Lawrence Radiation Laboratory
University of California, Berkeley, California 94720

April 1970

In an earlier Letter,[1] partial cross sections for K^+d and isospin-0 KN reactions at K^+ momenta of 865, 970, 1210 and 1365 MeV/c were presented. In this paper we present a discussion of the isospin-0 $K^+N \rightarrow K\pi N$ channel based on further analysis of K^+d data at 1210 and 1365 MeV/c plus recently analyzed data at 1585 MeV/c from the same exposure of the 25-inch bubble chamber at the Bevatron. Our results show that above the K^*N threshold at 1080 MeV/c most of the $I = 0$ $K\pi N$ production proceeds via the K^*N channel. The K^* production and decay angular distributions indicate the presence of several partial waves even near threshold and suggest that the main production mechanism is t-channel pion exchange. Thus there is no apparent support for an s-channel resonance interpretation of the peak in the $I = 0$ KN total cross section around 1200 MeV/c.[2]

(1) Methods.--(a) As shown earlier the isospin-0 $KN \rightarrow K\pi N$ cross section is given by

$$\sigma_o(KN \rightarrow K\pi N) = 3[\sigma(K^+n \rightarrow K^o\pi^+n) + \sigma(K^+n \rightarrow K^+\pi^-p) - \sigma(K^+p \rightarrow K^+\pi^op)] \ . \qquad (1)$$

Since the same relation applies to angular distributions and invariant-mass spectra, the K^*N cross section could be obtained by making the appropriate fit to the $K\pi N$ Dalitz plot obtained from Eq. (1).

(b) Alternatively, the K^*N cross section can also be determined from

$$\sigma_o(KN \to K^*N) = 3[\sigma(K^+n \to K^{*+}n \to K^o\pi^+n) + \sigma(K^+n \to K^{*o}p \to K^+\pi^-p) - \frac{1}{2}\sigma(K^+p \to K^{*+}p \to K^o\pi^+p)] \quad , \qquad (2)$$

in which each appropriate deuterium channel is individually separated into K^*N and the $K^o\pi^+p$ hydrogen channel rather than the $K^+\pi^op$ state is used to subtract out the $I = 1$ $K^{*+}N$ part of the cross section.

(2) Results.--In Fig. 1 we give the energy dependence of the isospin-0 and isospin-1 $KN \to K^*N$ cross sections from threshold to 3.0 BeV/c. The isospin-1 cross sections at 1210, 1365 and 1585 MeV/c are taken from the work of Bland et al.[3] Higher momentum data are taken from the compilation of Price et al.[4] The isospin-0 cross sections have been calculated using Eq. (2). At 1210, 1365, and 1585 MeV/c we have used the $K^+p \to K^{*+}p \to K^o\pi^+p$ data of Bland et al.[3] and at 3.0 BeV/c we have used the published K^+d data of Bassompierre et al.[5] to calculate the isospin-0 cross section. A smooth curve has been drawn through the isospin-1 $KN \to K^*N$ cross sections and has been scaled upward by a factor 2.5 to give the smooth curve passing through the isospin-0 $KN \to K^*N$ cross sections. The data indicate that the energy dependency of the isospin-0 and isospin-1 $KN \to K^*N$ cross sections are quite similar in this energy region.

In Fig. 2 we give the production and decay angular distributions for the isospin-0 $KN \to K^*N$ reaction at 1210, 1365, and 1585 MeV/c. For the 1210 and 1365 MeV/c data procedure (a) was used, whereas at 1585 MeV/c, the availability of only the $K^o\pi^+p$ final state in hydrogen permitted only procedure (b). Cosine θ is the K^* production angle cosine $\hat{K}^+\cdot\hat{K}^*$ in the K^*N center of mass, cos α is the polar-decay-angle cosine $\hat{K}^+\cdot\hat{K}$ in the K^* c.m., and angle φ is the Treiman-Yang azimuthal decay angle.

The K^* angular distributions in Fig. 2 are strikingly different from those produced in the isospin-1 state.[6] In the isospin-0 channel, the production angular distributions peak sharply in the forward direction, the polar decay angular distributions vary mainly as $\cos^2 \alpha$, and the azimuthal decay angular distributions are flat. In the isospin-1 channel (Ref. 6), the production distributions peak much less sharply, the polar decay distributions vary mainly as $\sin^2 \alpha$, and the azimuthal decay distributions vary mainly as $\sin^2 \varphi$. In the first case, the distributions are characteristic of exchange in the t-channel of a light pseudoscalar meson, whereas in the second case they are characteristic of exchange of a heavier vector meson. All the main features point to π exchange being the dominant K^* production mechanism in the isospin-0 channel, and ω,ρ exchange being dominant in the isospin-1 channel.

As is the case with $K\Delta$ and K^*N production in the $I = 1$ channel, the major inelastic process in the $I = 0$ channel in the region of the total cross-section bump, namely K^*N production, exhibits behavior which can be interpreted in terms of t-channel exchanges. Furthermore as is evident from Fig. 2 this process is highly peripheral even very close to the bump and therefore does not suggest dominance of one partial wave. There is therefore no evidence of resonant behavior near 1200 MeV/c at least insofar as the inelastic channels are concerned.

We gratefully acknowledge the contributions made to this experiment by the Bevatron and bubble-chamber crews, by the F.S.D. staff under the direction of Howard White, and by our scanning and programming groups. We especially thank Dr. Roger W. Bland for data and numerous discussions.

REFERENCES

*Present address: Nuclear Physics Laboratory, Oxford, England.

1. A. A. Hirata, C. G. Wohl, G. Goldhaber, and G. H. Trilling, Phys. Rev. Letters 21, 1485 (1968); Errata, Phys. Rev. Letters 21, 1728 (1968).
2. R. L. Cool, G. Giacomelli, T. F. Kycia, B. A. Leontic, K. K. Li, A. Lundby, and J. Teiger, Phys. Rev. Letters 17, 102 (1966); D. V. Bugg et al., Phys. Rev. 168, 1466 (1968).
3. R. W. Bland, M. G. Bowler, J. L. Brown, J. A. Kadyk, G. Goldhaber, S. Goldhaber, V. H. Seeger, and G. H. Trilling, Nucl. Physics B13, 595 (1969).
4. L. R. Price, N. Barash-Schmidt, O. Benary, R. W. Bland, A. H. Rosenfeld, and C. G. Wohl, UCRL-20000 K^+N, September 1969.
5. G. Bassompierre et al., Nucl. Physics B16, 125 (1970).
6. R. W. Bland, M. G. Bowler, J. L. Brown, G. Goldhaber, S. Goldhaber, J. A. Kadyk, V. H. Seeger, and G. H. Trilling, Nucl. Physics B17 (in press).

FIGURE CAPTIONS

Fig. 1. Energy dependence of the isospin-0 and isospin-1 $KN \rightarrow K^*N$ cross sections.

Fig. 2. Production and decay angular distributions of the isospin-0 $KN \rightarrow K^*N$ reaction at 1210, 1365, and 1585 MeV/c. Angles are defined in text.



XBL705-2756

Fig. 1



XBL 705-2753

Fig. 2

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

TECHNICAL INFORMATION DIVISION
LAWRENCE RADIATION LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720