Submitted to Nuclear Data

UCRL-19900
Preprint

c.2

RECEIVED
LAWRENCE
RADIATION LABORATORY

JUN 23 1970

LIBRARY AND
DOCUMENTS SECTION

TABLES OF THE CLEBSCH-GORDAN COEFFICIENTS FOR ODD- AND EVEN-MASS DEFORMED NUCLEI*

D. J. Gorman and F. Asaro

May 1970

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545

UCRL-19900 cy 2

LAWRENCE RADIATION LABORATORY
UNIVERSITY of CALIFORNIA BERKELEY

37

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

TABLES OF THE CLEBSCH-GORDAN COEFFICIENTS FOR ODD- AND EVEN-MASS DEFORMED NUCLEI*

D. J. Gorman and F. Asaro

Lawrence Radiation Laboratory
University of California
Berkeley, California 94720

May 1970

A set of tables has been prepared as a convenient source of the Clebsch-Gordan Coefficients commonly used in the calculation of nuclear transition probabilities for deformed nuclei. Similar, less extensive tables have been prepared by other authors (Si 54, Lu 61, and Ya 66).

The transition probability for a nuclear transition of multipole order L between an initial state I and a final state F is characterized by the reduced transition probability B(L, II → IF).

If the nuclear motion is separated into two modes--rotational and intrinsic--the reduced transition probability can be written as a product of a geometrical factor, depending only on the angular momenta, and a factor involving integrations over the intrinsic wave function of the initial and final states.†

If one compares the reduced transition probability for the emission of a given multipole radiation from a state I to different members F,F' ··· of a rotational family, the factors involving the intrinsic wave functions are the same. One thus obtains a ratio which depends only on the geometrical factors.

---

*Available as UCRL-18975 from Lawrence Radiation Laboratory, University of California, Berkeley, California 94720.

†This and most of the following discussion is taken from Ref. Al .

$$\frac{B(L,II \rightarrow IF)}{B(L,II \rightarrow IF')} = \frac{|\langle II\ L\ KI\ KF\text{-}KI | IF\ KF \rangle|^2}{|\langle II\ L\ KI\ KF\text{-}KI | IF'\ KF \rangle|^2}, \qquad (1)$$

where $\langle II\ L\ KI\ KF\text{-}KI | IF\ KF \rangle$ is the Clebsch-Gordan Coefficient (Co 35).

II = The spin of the initial state.

IF = The spin of the final state.

KI = The K of the initial state, where K is the projection of the nuclear angular momentum on the nuclear symmetry axis.

KF = The K of the final state

L = The angular momentum carried off by the radiation emitted during the transition from the initial to the final state.

The relation (1) holds when the initial and final states belong to the same or different rotational bands.

In special cases when $L \geqslant KI + KF$, symmetrization of the wave function may cause the transition matrix elements to become a sum of two products of geometrical and intrinsic factors. The ratio of reduced transition probabilities can then be written in the form.

$$\frac{B(L,II \rightarrow IF)}{B(L,II \rightarrow IF')} = \left[\frac{\langle II\ L\ KI\ KF\text{-}KI | IF\ KF \rangle + b(-)^{IF+KF} \langle II\ L\ KI\ -KF\text{-}KI | IF\ -KF \rangle}{\langle II\ L\ KI\ KF\text{-}KI | IF'\ KF \rangle + b(-)^{IF'+KF} \langle II\ L\ KI\ -KF\text{-}KI | IF'\ -KF \rangle}\right]^2 \qquad (2)$$

where b is a parameter depending on the intrinsic wave function.

The reduced transition probabilities are also important in Coulomb excitation experiments. The cross section for Coulomb excitation of particular states is proportional to $B(L,II \rightarrow IF)$.

THE ARRANGEMENT OF THE TABLES

The compilation is arranged into four tables covering integral and half-integral values of the spin (I) and positive and negative values of KF. The range of the arguments used is shown in the table below.

Table I. Range of Arguments for Clebsch-Gordan Coefficients

Table	KI	KF	II	L	IF
1	1/2 - 31/2	1/2 - 31/2	1/2 - 31/2	0 - 6	1/2 - 43/2
2	1/2 - 11/2	1/2 - (-11/2)	1/2 - 31/2	0 - 6	1/2 - 43/2
3	0 - 16	0 - 16	0 - 16	0 - 6	0 - 22
4	0 - 5	-1 - (-6)	0 - 16	0 - 6	0 - 22

Some sample pages from the tables are reproduced below. The tables are arranged such that the right hand argument IF varies the fastest, followed by L, II, KF, and KI.

KI	KF	II	L	IF	(CLER)**2	CLER
1/2	1/2	5/2	0	5/2	1.00000000	1.00000000
1/2	1/2	5/2	1	3/2	.40000000	-.63245553
				5/2	.02857143	.16903085
				7/2	.57142857	.75592895
1/2	1/2	5/2	2	1/2	.20000000	.44721360
				3/2	.05714286	-.23904572
				5/2	.22857143	-.47809144
				7/2	.03809524	.19518001
				9/2	.47619048	.69006556
1/2	1/2	5/2	3	1/2	.14285714	.37796447
				3/2	.11428571	.33806170
				5/2	.07619048	-.27602622
				7/2	.19047619	-.43643578
				9/2	.04329004	.20806259
				11/2	.43290043	.65795169
1/2	1/2	5/2	4	3/2	.19047619	.43643578
				5/2	.09523810	.30860670
				7/2	.08658009	-.29424494
				9/2	.17316017	-.41612519
				11/2	.04662005	.21591676
				13/2	.40792541	.63869039
⋮	⋮	⋮		⋮	⋮	⋮
5/2	3/2	5/2	1	3/2	.66666667	.81649658
				5/2	.28571429	.53452248
				7/2	.04761905	.21821789
5/2	3/2	5/2	2	3/2	.28571429	.53452248
				5/2	.42857143	.65465367
				7/2	.23809524	.48795004
				9/2	.04761905	.21821789
5/2	3/2	5/2	3	3/2	.07142857	.26726124
				5/2	.28571429	.53452248
				7/2	.38095238	.61721340
				9/2	.21645022	.46524211
				11/2	.04545455	.21320072
5/2	3/2	5/2	4	3/2	.00793651	.08908708
				5/2	.09523810	.30860670
				7/2	.28860029	.53721531
				9/2	.36075036	.60062498
				11/2	.20396270	.45162230
				13/2	.04351204	.20859541
⋮	⋮	⋮		⋮	⋮	⋮
1/2	-1/2	3/2	5	7/2	.36363636	-.60302269
				9/2	.04545455	-.21320072
				11/2	.22377622	.47304992
				13/2	.36713287	.60591490
1/2	-1/2	3/2	6	9/2	.36713287	-.60591490
				11/2	.05594406	-.23652496
				13/2	.20769231	.45573272
				15/2	.36923077	.60764362
1/2	-1/2	5/2	1	3/2	.20000000	.44721360
				5/2	.51428571	.71713717
				7/2	.28571429	.53452248
1/2	-1/2	5/2	2	1/2	.13333333	-.36514837
				3/2	.23809524	-.48795004
				5/2	.00000000	.00000000
				7/2	.31111111	.55777335
				9/2	.31746032	.56343617
1/2	-1/2	5/2	3	1/2	.19047619	.43643578
				3/2	.00952381	-.09759001
				5/2	.22857143	-.47809144
				7/2	.01587302	-.12598816
				9/2	.23088023	.48049998
				11/2	.32467532	.56980288

Fig. 1

EXAMPLES OF THE USE OF THE TABLES

A. An E2 transition from a K = 5/2 to a K = 3/2 rotational band.



We can obtain the relative transition probabilities from the spin 5/2 member of the K = 5/2 rotational band to the different rotational members of the K = 3/2 band using formula 1. In this case II = 5/2, KI = 5/2, KF = 3/2, L = 2, and IF = 9/2, 7/2, 5/2 and 3/2.

$$B(2,5/2 \rightarrow 3/2) \;\alpha\; |\langle 5/2\ 2\ 5/2\ -1|3/2\ 3/2\rangle|^2 = 0.28571429$$

$$B(2,5/2 \rightarrow 5/2) \;\alpha\; |\langle 5/2\ 2\ 5/2\ -1|5/2\ 3/2\rangle|^2 = 0.42857143$$

$$B(2,5/2 \rightarrow 7/2) \;\alpha\; |\langle 5/2\ 2\ 5/2\ -1|7/2\ 3/2\rangle|^2 = 0.23809524$$

$$B(2,5/2 \rightarrow 9/2) \;\alpha\; |\langle 5/2\ 2\ 5/2\ -1|9/2\ 3/2\rangle|^2 = 0.04761905$$

B. An M1 transition within a K = 1/2 rotational band with a decoupling constant (a) of -1.5.



In this case, to find the relative transition probabilities for the transitions from the 5/2 to the 7/2 and 3/2 members of the K = 1/2 band we must use formula 2. Here we have II = 5/2, KI = KF = 1/2, L = 1, and IF = 7/2 and 3/2.

$$B(1,5/2 \rightarrow 7/2) \;\alpha\; [\langle 5/2\ 1\ 1/2\ 0|7/2\ 1/2\rangle + b\langle 5/2\ 1\ 1/2\ 0|7/2\ -1/2\rangle]^2$$

$$= [0.75592895 + b\ (0.53452248)]^2$$

$$B(1,5/2 \rightarrow 3/2) \;\alpha\; [\langle 5/2\ 1\ 1/2\ 0|3/2\ 1/2\rangle + b\langle 5/2\ 1\ 1/2\ 0|3/2\ -1/2\rangle]^2$$

$$= [-0.63245553 + b\ (0.44721360]^2$$

THE COMPUTER PROGRAM

The computer program consists of a main program, which is slightly different for each of the four tables, and a subroutine CLEB which remains the same. The program is written in Chippewa Fortran for the CDC 6600.

The main program for calculating the coefficients for half-integral values of I and negative KF is reproduced below.

```
                PROGRAM IDA(INPUT,OUTPUT,TAPE2=INPUT,TAPE3=OUTPUT)
          C
          C     THIS PROGRAM CALCULATES CLEBSCH GORDON COEFFICIENTS FOR
          C     HALF-INTEGRAL ARGUMENTS
          C     AND NEGATIVE VALUES OF KF
          C
          C     X(1) = II
          C     X(2) = L
          C     X(3) = IF
          C     X(4) = KI
          C     X(5) = KF - KI
          C     X(6) = KF
          C
000043          DATA LINE /0/
000043          DATA CUT/0.0000000001/
000043          DIMENSION X(6),M(6)
000043          DIMENSION FMT(4)
000043          DATA NPAGE/1/
000043          FMT(1) = 10H(1H1,5X,*-
000045          FMT(3) = 10H0X,*UCPI-1
000046          FMT(4) = 10H8975*)
000050          DIGIT1 = 10H*,I1,*-*,5
000051          DIGIT2 = 10H*,I2,*-*,5
000053          DIGIT3 = 10H*,I3,*-*,5
000054          DIGIT4 = 10H*,I4,*-*,5
000056          PRINT 1
000061        1 FORMAT(6H1*CVF*)
000061          ICOUNT = 0
000062          IF(NPAGE.LE.9999) FMT(2) = DIGIT4
000067          IF(NPAGE.LE.999) FMT(2) = DIGIT3
000073          IF(NPAGE.LE.99) FMT(2) = DIGIT2
000077          IF(NPAGE.LE.9) FMT(2) = DIGIT1
000103          WRITE(3,FMT) NPAGE
000111          NPAGE = NPAGE + 1
000113          WRITE(3,3003)
000116          WRITE(3,3000)
000121          WRITE(3,3010)
000124          ICOUNT = ICOUNT + 5
000126          DO 80 I4 = 1,11,2
000127          X(4) = FLOAT(I4)/2.
000131          PMAX = (6. - X(4))*2.
000133          KMAX = IF(X(PMAX)
000135          DO 60 I6 = 1,KMAX,2
000136          X(6) = -FLOAT(I6)/2.
000140          X(5) = X(6) - X(4)
000142          DO 50 I1 = I4,31,2
000144          X(1) = FLOAT(I1)/2.
000146          DO 40 I2 = 1,7
000147          LINE = 0
000150          X(2) = FLOAT(I2) - 1.
000152          IF(ABS(X(5)).GT.X(2)) GO TO 40
000157          JMIN = ABS(X(1) - X(2)) * 2
000163          JMAX = ABS(X(1) + X(2)) * 2
000167          KMIN = ABS(X(6)) * 2
000172          IF(JMIN.LT.KMIN) JMIN = KMIN
000174          DO 30 I3 = JMIN,JMAX,2
000176          X(3) = FLOAT(I3)/2.
000200          Y = CLER(X)
000202          IF(ABS(Y).EQ.0) Y=CUT
000205          Z = Y**2
000207          LINE = LINE + 1
000210          M(1) = 2 * X(1)
000213          M(2) = X(2)
000215          M(3) = 2 * X(3)
000220          M(4) = 2 * X(4)
000223          M(5) = X(5)
000224          M(6) = 2 * X(6)
000227          IF(LINE.GT.1) GO TO 20
000232          WRITE(3,3001) M(4),M(6),M(1),M(2),M(3),Z,Y
000253          ICOUNT = ICOUNT + 2
000255          IF(ICOUNT.GE.60) GO TO 100
000257          GO TO 30
000257       20 WRITE(3,3002) M(3),Z,Y
000270          ICOUNT = ICOUNT + 1
000272          IF(ICOUNT.GE.60) GO TO 100
000274       30 CONTINUE
000277       40 CONTINUE
000301       50 CONTINUE
000303       60 CONTINUE
000306       80 CONTINUE
000310          CALL EXIT
000311      100 ICOUNT = 0
000312          IF(NPAGE.LE.9999) FMT(2) = DIGIT4
000317          IF(NPAGE.LE.999) FMT(2) = DIGIT3
000323          IF(NPAGE.LE.99) FMT(2) = DIGIT2
000327          IF(NPAGE.LE.9) FMT(2) = DIGIT1
000333          WRITE(3,FMT) NPAGE
000341          NPAGE = NPAGE + 1
000343          WRITE(3,3003)
000346          WRITE(3,3000)
000351          WRITE(3,3010)
000354          ICOUNT = ICOUNT + 5
000356          LINE = 0
000357          GO TO 30
000357     3010 FORMAT(1H0)
000357     3000 FORMAT(1H0,7X,2HKI,6X,2HKF,6X,2HII,6X,1HL,6X,2HIF,7X,9H(CLER)**2,
               110X,4HCLER)
000357     3001 FORMAT(1H0,5X,I3,2H/2,3X,I3,2H/2,3X,I3,2H/2,3X,I3,4X,I3,2H/2,3X,
               1F12.8,5X,F12.8)
000357     3002 FORMAT(1H ,36X,I3,2H/2,3X,F12.8,5X,F12.8)
000357     3003 FORMAT(1H )
000357          END
```

Fig. 2

```
                FUNCTION CLEB(X)
000535          DIMENSION X(6)
000535          DIMENSION FACT(101)
000535          DATA EPS/10.E-5/
000535          DATA(FACT(I),I=1,30)/
               X 1.000000000000000E+00, 1.000000000000000E+00,
               X 1.414213562373341E+00, 2.449489742783178E+00,
               X 4.898979485566850E+00, 1.095445115010980E+01,
               X 2.683281573000885E+01, 7.099295739772260E+01,
               X 2.007984063683618E+02, 6.023952191048888E+02,
               X 1.904940943967628E+03, 6.317974359925955E+03,
               X 2.188610518115346E+04, 7.891147465084575E+04,
               X 2.952597012400218E+05, 1.143535905864376E+06,
               X 4.574143623458277E+06, 1.885967730676345E+07,
               X 8.001483428549577E+07, 3.487765766345658E+08,
               X 1.559776266828958E+09, 7.147792818187560E+09,
               X 3.352617008237675E+10, 1.607856235454208E+11,
               X 7.870854713229575E+11, 3.938427356461453E+12,
               X 2.008211794424299E+13, 1.043497458090628E+14,
               X 5.521669535670754E+14, 2.973510046011428E+15/
000535          DATA(FACT(I),I=31,60)/
               X 1.624654327168638E+16, 9.067986906786461E+16,
               X 5.129628026880119E+17, 2.966746955319335E+18,
               X 1.718233974286175E+19, 1.016520927790088E+20,
               X 6.099125566743515E+20, 3.709953246495951E+21,
               X 2.286968774306135E+22, 1.428211541793825E+23,
               X 9.032802905216805E+23, 5.788159721435645E+24,
               X 3.748341123414035E+25, 2.457951684890475E+26,
               X 1.630420674175555E+27, 1.093719437813105E+28,
               X 7.417966136206525E+28, 5.085501166729755E+29,
               X 3.523338699653905E+30, 2.466337089757225E+31,
               X 1.743963680858765E+32, 1.245439180883175E+33,
               X 8.980989654291025E+33, 6.538259159770855E+34,
               X 4.806619674253915E+35, 3.563201279846415E+36,
               X 2.666455677111365E+37, 2.013129899111718E+38,
               X 1.533156046814455E+39, 1.177637969751375E+40/
000535          DATA(FACT(I),I=61,90)/
               X 9.121944481667935E+40, 7.124466393154765E+41,
               X 5.609810447782435E+42, 4.452649004111665E+43,
               X 3.562119203289335E+44, 2.871872314708405E+45,
               X 2.333120097790115E+46, 1.909741105955415E+47,
               X 1.574812859487115E+48, 1.308137807824095E+49,
               X 1.094466613004625E+50, 9.222139602145115E+50,
               X 7.825244940320435E+51, 6.685892207809145E+52,
               X 5.751421947193555E+53, 4.940877514150825E+54,
               X 4.362228346865215E+55, 3.810289910567215E+56,
               X 3.365156932149475E+57, 2.991016905770175E+58,
               X 2.675244849262255E+59, 2.407722164336145E+60,
               X 2.180285150366085E+61, 1.986334104600975E+62,
               X 1.820505461261825E+63, 1.678423103484875E+64,
               X 1.556505553574135E+65, 1.451911729640345E+66,
               X 1.361920123400395E+67, 1.284832787452825E+68/
000535          DATA(FACT(I),I=91,101)/
               X 1.218899489063045E+69, 1.162756005204145E+70,
               X 1.115276380735025E+71, 1.075533591778915E+72,
               X 1.042769505767755E+73, 1.016365017495115E+74,
               X 9.958302741108335E+74, 9.807790764431755E+75,
               X 9.709217501187775E+76, 9.660549437804065E+77,
               X 9.660549437794465E+78/
000535          CLEB=0.
000536          NA=2.*X(1)+EPS
000541          NB=2.*X(2)+EPS
000543          NC=2.*X(3)+EPS
000546          ND=SIGN(ABS(2.*X(4))+EPS,X(4))
000553          NE=SIGN(ABS(2.*X(5))+EPS,X(5))
000561          NF=-SIGN(ABS(2.*X(6))+EPS,X(6))
000567          IF(ABS(X(4)+X(5)-X(6)) .GT. EPS) RETURN
000576          L3=X(2)+X(5)+EPS
000602          L8=X(1)+X(4)+EPS
000605          L10=X(3)-X(6)+EPS
000610          IF(ABS(X(2)+X(5)-FLOAT(L3)) .GT. EPS) RETURN
000617          IF(ABS(X(1)+X(4)-FLOAT(L8)).GT. EPS) RETURN
000626          IF(ABS(X(3)-X(6)-FLOAT(L10)) .GT. EPS) RETURN
000636          L0=L3+L8+L10
000641          IF(L0 .GT. 99) GO TO 2
000644          IF(ABS(ND) .GT. NA  .OR.
               X   ABS(NE) .GT. NB  .OR.
               X     ABS(NF) .GT. NC) RETURN
000663          L1=L0-NC
000665          L2=L8-ND
000667          L9=L3-NF
000671          L11=L10-NF
000673          L6=L0-NB
000675          L7=L0-NA
000677          L4=L6-L2
000701          L5=L7-L3
000703          L20=NC+1
000705          KMIN=MAX0(0,-L4,-L5)
000712          KMAX=MIN0(L1,L2,L3)
000720          IF(KMAX .LT. KMIN)RETURN
000724          COI=FACT(L1+1)*FACT(L6+1)*FACT(L7+1)*FACT(L8+1)
               X    *FACT(L2+1)*FACT(L3+1)
               X*FACT(L9+1)*FACT(L10+1)*FACT(L11+1)/FACT(L0+2)
               X  *FACT(L20+1)/FACT(L20)
000747          SGN=1.
000751          DO 1 K=KMIN,KMAX
000763          L12=L1-K+1
000765          L13=L2-K+1
000767          L14=L3-K+1
000770          L15=L4+K+1
000771          L16=L5+K+1
000773          DEN=FACT(K+1)*FACT(L12)*FACT(L13)*FACT(L14)*FACT(L15)*FACT(L16)
001002          CLEB=CLEB+SGN*COI/DEN**2
001006          SGN=-SGN
001006        1 CONTINUE
001016          IF(MOD(KMIN,2) .NE. 0) CLEB=-CLEB
001022          RETURN
001024        2 CLEB=0./0.
001026          RETURN
001027          END
```

Fig. 2

To calculate the coefficients for positive values of KF, the following changes should be made.

DO 80 I4 = 1,11,2	becomes	DO 80 I4 = 1,31,2
PMAX = (6. - X(4))* 2.	becomes	PMAX = (X(4) + 6.)* 2.
X(6) = -FLOAT(I6)/2.	becomes	X(6) = FLOAT(I6)/2.

The variable NPAGE which is set in a data statement gives the initial page number. This enables one to calculate the four tables successively and keep consecutive page numbers.

The statements [PRINT 1 / 1 FORMAT(6H1*OVF*)] suppress the automatic skip to a new page on the printer.

For calculating the coefficients with integral arguments and negative KF, the lines from 000126 through 000224 should be replaced by the following sequence.

```
000126        DO 80 I4 = 0,6
000127        X(4) = FLOAT(I4)
000130        PMAX = (6. - X(4))
000132        KMAX = IFIX(PMAX)
000133        DO 60 I6 = 1,KMAX
000135        X(6) = -FLOAT(I6)
000136        X(5) = X(6) - X(4)
000140        DO 50 I1 = I4,16
000142        X(1) = FLOAT(I1)
000143        DO 40 I2 = 0,6
000145        LINE = 0
000146        X(2) = FLOAT(I2)
000147        IF(ABS(X(5)).GT.X(2)) GO TO 40
000154        JMIN = ABS(X(1) - X(2))
000157        JMAX = ABS(X(1) + X(2))
000162        KMIN = ABS(X(6))
000164        IF(JMIN.LT.KMIN) JMIN = KMIN
000166        DO 30 I3 = JMIN,JMAX
000170        X(3) = FLOAT(I3)
000171        Y = CLEH(X)
000174        IF(ABS(Y).EQ.0) Y = CUT
000177        Z = Y**2
000201        LINE = LINE + 1
000202        DO 10 J = 1,6
000207     10 M(J) = X(J)
```

Fig. 3

For positive KF and integral arguments make the following substitutions.

DO 80 I4 = 0,6	becomes	DO 80 I4 = 0,16
PMAX = 6. - X(4)	becomes	PMAX = X(4) + 6.
DO 60 I6 = 1,KMAX	becomes	DO 60 I6 = 0,KMAX
X(6) = -FLOAT(I6)	becomes	X(6) = FLOAT(I6)

The subroutine CLEB was written by Marjory Simmons of the Math and Computing division at LRL Berkeley.

REFERENCES

Al G. Alaga, K. Alder, A. Bohr, and B. R. Mottelson, Dan. Mat. Fys. Medd. 29, NO. 9.

Co 35 E. U. Condon and G. H. Shortley, "The Theory of Atomic Spectra," The University Press, Cambridge (1935).

Lu 61 J. K. Lum, J. H. Light, and F. Asaro, "Clebsch-Gordan Coefficients for Nuclear Transition Probabilities: I Odd-Mass Nuclides," Lawrence Radiation Laboratory Report UCRL-9679, August, 1961.

Si 54 A. Simon, "Numerical Tables of the Clebsch-Gordan Coefficients," Oak Ridge Report ORNL-1718, July, 1954.

Ya 66 T. Yamazaki, "Clebsch-Gordan Coefficients for Nuclear Transition Probabilities for Deformed Nuclei," Lawrence Radiation Laboratory Report UCRL-16784, March, 1966; Nucl. Data A, 1, No. 5, July, 1966.

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

TECHNICAL INFORMATION DIVISION
LAWRENCE RADIATION LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720