Submitted to Physical Review Letters

UCRL-19929
Preprint

c. 2

RECEIVED
LAWRENCE
RADIATION LABORATORY
JUL 29 1970
LIBRARY AND
DOCUMENTS SECTION

THE MASS OF ^{13}O AND THE ISOBARIC MULTIPLET MASS EQUATION

Robert Mendelson, G. J. Wozniak,
A. D. Bacher, J. M. Loiseaux, and Joseph Cerny

July 1970

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

UCRL-19929

LAWRENCE RADIATION LABORATORY
UNIVERSITY of CALIFORNIA BERKELEY

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

THE MASS OF ^{13}O AND THE ISOBARIC MULTIPLET MASS EQUATION*

Robert Mendelson, G. J. Wozniak, A. D. Bacher, J. M. Loiseaux,† and Joseph Cerny

Lawrence Radiation Laboratory and Department of Chemistry
University of California
Berkeley, California 94720

July 1970

Abstract:

The mass excess of ^{13}O has been determined to be (23.107±0.015) MeV. This yields a value for d, the coefficient of T_z^3 in the isobaric multiplet mass equation, of (-0.8±3.4) keV for the A = 13 quartet. Additional values for d, obtained from new measurements of the masses of ^{17}Ne, ^{21}Mg and ^{25}Si, are also consistent with zero.

Although the level energies of several isobaric multiplets with T = 3/2 have been determined,[1] only the A = 9 quartet has been measured with sufficient accuracy to observe a significant deviation from the quadratic form of the isobaric multiplet mass equation, $\Delta M = a + bT_z + cT_z^2$. If this deviation is paramaterized by an additional term dT_z^3 in the mass equation, then d has a value[2] of (9.2±3.7) keV. The non-zero value for d indicates that a first-order perturbation treatment of the isospin-nonconserving operator (assuming charge-dependent forces of tensorial rank two or less) is inadequate to describe the A = 9 quartet. Several theoretical attempts to evaluate the magnitude of d have led to conflicting results. A second-order perturbation

calculation,[3] using Coulomb-energy systematics to estimate the mixing of $T = 1/2$ states with the $T = 3/2$ states in ^{9}Be and ^{9}B, gives a d coefficient comparable to that observed. However, further considerations based on the widths of the perturbing $T = 1/2$ states indicate that the $T = 1/2$ admixture is small and not consistent with the large value observed for d.[4] Henley and Lacy,[5] applying a simple nonperturbative model to the $A = 4n + 1$ isospin quartets, find generally small values for d ($\lesssim 1$ keV) and, in particular, obtain d = 0.06 keV for A = 9.

In order to investigate systematics that might appear in the d coefficient, we have accurately remeasured the mass of ^{13}O and we report new or improved values for the masses of ^{17}Ne, ^{21}Mg, and ^{25}Si. In all cases the values obtained for d are essentially consistent with zero. In particular, using the well-known masses for the other members of the A = 13 multiplet,[1] no deviation from the quadratic mass equation is observed within an error comparable to the uncertainty in the A = 9 determination.

The experiments were performed with ^{3}He and ^{4}He beams from the Berkeley 88-inch variable-energy cyclotron. The ^{13}O mass measurement employed semiconductor counter-telescopes to observe the $^{16}O(^{3}He,^{6}He)^{13}O$ reaction and the calibration reactions $^{12}C(^{3}He,^{6}He)^{9}C$ and $^{12}C(\alpha,^{6}He)^{10}C$. ^{3}He ions, bombarding a CO_2 gas target, produced ^{6}He nuclei corresponding to the ^{9}C and ^{13}O ground states. Since the peaks due to ^{9}C and ^{13}O were separated by approximately 1.4 MeV, the known mass[2] for the ^{9}C ground state provided an accurate energy calibration. The slope of the energy scale was determined by observing the ground state and first-excited state (E_x = 3.3527 MeV)[6] in ^{10}C produced by alpha-particle bombardment of ethane.

Two similar four-counter telescopes (consisting of 114μ, 90μ, 300μ and 500μ counters) and electronic systems were simultaneously employed at equal angles on opposite sides of the beam axis.[7] Two particle-identifications were performed and compared using the signals from the two successive differential-energy-loss (ΔE) detectors and the third E detector. Sub-nanosecond pulse-shape discrimination was employed to reduce pile-up due to ^{3}He's and deuterons from the same beam burst which can produce a false signal in the ^{6}He region of the particle-identifier spectrum. Pulser calibrations following each event were used to correct the slope and zero-point of the energy scale for each system. A split Faraday cup and the observation of elastically-scattered particles were used to monitor the beam position.

Five parameters were recorded for each event: the total energy, two ΔE losses, particle identification, and pile-up. Corrections were made and limits were placed on these parameters using an off-line computer. Peak centroids were determined using Chauvenet's theorem[8] to systematically set limits on the extent of the ^{6}He peaks. A typical set of spectra are shown in Fig. 1. The resolution was 200-300 keV.

The ^{13}O mass was obtained by averaging the Q-values measured in each detector system. Two independent runs at $E_{^3He}$ = 62.6 MeV, θ_{Lab} = 11.65 deg. and at $E_{^3He}$ = 66.3 MeV, θ_{Lab} = 12.19 deg. were combined to yield an ^{13}O mass excess of (23.107±0.015) MeV based on a ^{9}C mass excess of (28.906±0.004) MeV.[2] The uncertainty in the mass excess is primarily due to statistical errors and the uncertainty in the beam energy. Errors due to uncertainties in the scattering angle, the slope of the energy scale, counter dead-layers, gas-target foil corrections, gas-pressure variations and energy fluctuations in the

beam were small because of the differential nature of the measurement. Small corrections were made for the shifts in each peak position due to the angular distribution of the reaction products across the solid angle (5×10^{-5} sr) of the detectors. The mass excess obtained is in excellent agreement with the previously-measured value, (23.110±0.070) MeV.[9]

The masses of ^{17}Ne, ^{21}Mg and ^{25}Si, measured using similar techniques but with less precision, are given in Table 1.[10] For the ^{17}Ne measurement, a gas target containing a mixture of 97% enriched ^{20}Ne gas and CO_2 was used. Carbon-backed ^{24}Mg (99% enriched) and natural SiO_2 targets were used for the ^{21}Mg and ^{25}Si measurements, respectively. The overall resolution for the solid targets was ≈ 175 keV.

The mass of ^{17}Ne (16.479±0.050 MeV) is found to be in good agreement with a previous value[11] of (16.47±0.25) MeV. The new mass of ^{21}Mg, (10.889±0.040) MeV, supersedes the previous measurement of (10.62±0.12) MeV from this laboratory.[12] For the ^{25}Si measurement the group observed could be composed of both the ground and first-excited states which are expected to lie ≈ 30 keV apart. If the first-excited state is populated to a significant extent, the d coefficient will be slightly more positive than that given in Table 1, but it will still be consistent with zero. Values of a, b, c, and χ^2 determined by a linear-least-squares fit to the quadratic mass equation are also given in Table 1. With the exception of A = 9 all are consistent with a value of $\chi^2 \lesssim 1$. Values for d are given in the last column of Table 1.

Our value for d, (−0.8±3.4) keV, for the A = 13 quartet is significantly less than the value, (9.2±3.7) keV, previously determined for A = 9. Since additional values of d for the quartets with A > 13 are also essentially

consistent with zero, there does not appear to be a systematic increase in d with increasing average charge of the multiplet. This suggests that the non-zero d coefficient for the A = 9 quartet is not due primarily to charge-dependent forces of tensorial rank 3 or higher.

It is of particular interest to understand the variation in magnitude of the d coefficient in the more-accurately measured A = 9 and A = 13 quartets. In the absence of higher-order, charge-dependent forces, a non-zero value for d might arise from level shifts due to nearby thresholds for isospin-allowed decays[13] or, in a perturbation treatment (where d depends on the product of off-diagonal vector and tensor Coulomb matrix elements[1,3]), it might be due to mixing of the ($T = 3/2$, $T_z = \pm 1/2$) levels with other $J^{\pi} = 3/2^-$ levels with either $T = 3/2$ or $T = 1/2$. A reliable test of the perturbation approach requires a knowledge of all of the parameters of the appropriate $T = 3/2$ and $T = 1/2$ levels in the $T_z = \pm 1/2$ nuclei. In both the A = 9 and A = 13 systems it is therefore necessary to identify and characterize the ($3/2^-$, $T = 1/2$) states predicted[14] to lie in the energy range close to the analogue levels. In addition, for the A = 9 system a more accurate determination of the width of the ^9B analogue state is needed.

The authors wish to thank Penny Fink for writing the off-line computer analysis program.

FOOTNOTES AND REFERENCES

*Work performed under the auspices of the U. S. Atomic Energy Commission.

†NATO Fellow; permanent address: I.P.N., Orsay, France.

1. J. Cerny, Ann. Rev. Nucl. Sci. 18, 27 (1968). All values of masses and excitation energies are those given by G. T. Garvey, in Nuclear Isospin, ed. by J. D. Anderson, S. D. Bloom, J. Cerny, and W. W. True (Academic Press, New York, 1969), p. 703, unless explicitly stated otherwise.
2. J. M. Mosher, R. W. Kavanagh and T. A. Tombrello, Bull. Am. Phys. Soc. 14, 1167 (1969).
3. J. Jänecke, Isospin in Nuclear Physics, ed. by D. H. Wilkinson (North-Holland Publishing Company, Amsterdam, 1969), Chap. 8.
4. J. Hardy, J. M. Loiseaux, J. Cerny and G. T. Garvey, to be published.
5. E. M. Henley and C. F. Lacy, Phys. Rev. 184, 1228 (1969).
6. R. A. Paddock, S. M. Austin, W. Benenson, I. D. Proctor, and F. St. Amant, Phys. Rev. 182, 1104 (1969).
7. The system used, with the exceptions discussed in the text, was identical to that reported in J. Cerny, R. A. Mendelson, Jr., G. J. Wozniak, John E. Esterl, and J. C. Hardy, Phys. Rev. Letters 22, 612 (1969).
8. This theorem is discussed in H. Jeffreys, Theory of Probability (Oxford University Press, Oxford, 1950), p. 357.
9. J. Cerny, R. H. Pehl, G. Butler, D. G. Fleming, C. Maples, and C. Detraz, Phys. Letters 20, 35 (1966).
10. The values of the measured masses are relevant to the possibility of an induced isotensor interaction in beta decay as discussed by D. H. Wilkinson, Phys. Letters 31B, 477 (1970). Although the values we obtain are essentially

the same as those used by Wilkinson , the somewhat smaller errors obtained here reinforce his thesis.

11. R. A. Esterlund, R. McPherson, A. M. Poskanzer, and P. L. Reeder, Phys. Rev. 156, 1094 (1967); A. M. Poskanzer, private communication.

12. G. W. Butler, J. Cerny, S. W. Cosper, and R. L. McGrath, Phys. Rev. 166, 1096 (1968).

13. Garvey, Ref. 1, estimates that this effect should be small for the lowest T = 3/2 states but, to our knowledge, no explicit evaluation of this effect has been made for the T = 3/2 levels in the A = 9 or A = 13 systems.

14. S. Cohen and D. Kurath, Nucl. Phys. 73, 1 (1965).

FIGURE CAPTION

Fig. 1. Typical spectra representing approximately 25% of the total data. The short arrows indicate the region used in calculating the centroids of the peaks. The long arrows mark the positions of these centroids.

Table 1. Completed $4n + 1$, $T = 3/2$ Multiplets. Values of a, b and c are obtained from a least squares fit to $\Delta M = a + bT_z + cT_z^2$. d is the coefficient of an additional T_z^3 term in ΔM. Errors, in parentheses, are in keV.

A	$T_z = -3/2$ Mass Excess (MeV)	IMME Prediction[c] (MeV)	a (MeV)	b (MeV)	c (MeV)	χ^2	d (keV)
9	28.906 (4)[a]	28.961 (22)	26.343 (4.0)	-1.315 (2.0)	0.263 (2.4)	6.16	9.2 (3.7)
13	23.107 (15)[b]	23.102 (14)[d]	19.257 (2.2)	-2.181 (3.4)	0.256 (2.7)	0.06	-0.8 (3.4)
17	16.479 (50)[b]	16.508 (23)[e]	11.651 (3.7)	-2.878 (5.4)	0.238 (8.1)	0.28	4.8 (9.2)
21	10.889 (40)[b]	10.940 (24)	4.897 (4.8)	-3.658 (6.8)	0.241 (6.4)	1.18	8.5 (7.8)
25	3.817 (50)[b]	3.796 (42)	-3.262 (8.1)	-4.387 (10.6)	0.216 (10.8)	0.10	-3.5 (10.9)
37	-13.230 (50)[c]	-13.198 (91)	-22.886 (11.6)	-6.181 (14.6)	0.174 (14.6)	0.10	5.3 (17.3)

[a] Ref. 2.

[b] Present work. For the ^{25}Si measurement we have assumed that only the ground state is populated.

[c] Ref. 1.

[d] An improved value for the ^{13}N state (M. J. Levine and P. D. Parker. Phys. Rev. 186, 1021 (1969)) and for the ^{13}C state (K. A. Snover, E. G. Adelberger and F. Riess, Bull. Am. Phys. Soc. 13, 1662 (1968)) has been included in the weighted average.

[e] A new value for the ^{17}O state (C. Detraz and H. H. Duhm, Phys. Letters 29B, 29 (1969)) has been included in the weighted average.



XBL706-3042

Fig. 1.

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

TECHNICAL INFORMATION DIVISION
LAWRENCE RADIATION LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720