Submitted to Physics Letters A

UCRL-19972
Preprint

c.2

THE HYPERFINE STRUCTURE ANOMALY OF ^{241}Pu AND ^{239}Pu AND THE NUCLEAR MOMENT OF ^{241}Pu

N. Edelstein

September 1970

RECEIVED
LAWRENCE
RADIATION LABORATORY
SEP 22 1970
LIBRARY AND
DOCUMENTS SECTION

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

UCRL-19972

c.2

LAWRENCE RADIATION LABORATORY
UNIVERSITY of CALIFORNIA BERKELEY

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

THE HYPERFINE STRUCTURE ANOMALY OF ^{241}Pu AND ^{239}Pu AND THE NUCLEAR MOMENT OF ^{241}Pu*

N. Edelstein

Lawrence Radiation Laboratory
University of California
Berkeley, California 94720

September 1970

The hfs anomaly of an s electron for the ^{241,239}Pu isotopes is shown to be $^{241}\Delta^{239} = -6.3 \pm 0.2\%$. The value for $\mu_N(^{241}\text{Pu})$ is found to be -0.718 ± 0.017 nm.

— — —

The hyperfine structure (hfs) constants for a number of levels in the Pu I and Pu II optical spectra have been measured by Fabry-Perot interferometry for the isotopes ^{241}Pu (I = 5/2, Z = 94, A = 147) and ^{239}Pu (I = 1/2, A = 145) [1]. The measured ratio of the hfs constants differed from that found earlier by Bleaney et al. [2] in electron paramagnetic resonance (epr) experiments of $^{241,239}PuO_2^{++}$ (Pu VII) diluted in a crystal of $UO_2Rb(NO_3)_3$. In the course of ENDOR and epr measurements on $^{241,239}Pu^{3+}$ (Pu IV) in CaF_2 [3] this ratio has been remeasured. The data from the various measurements are collected in Table I.

In a recent analysis of the hfs of Pu IV in CaF_2 [4] it was shown that approximately 5% of the total hyperfine interaction comes from unpaired s electrons through the mechanism of core polarization. The ground configuration of

*Work performed under the auspices of the U. S. Atomic Energy Commission.

Pu II, f^6s, is fairly pure and can be described by coupling the two lowest levels of the f^6 core ($^7F_{0,1}$) with an s electron and then mixing these two J = 1/2 states [5,6]. The hfs of the J = 1 level of Pu I (f^6s^2) has been shown to be extremely small [7], so almost all the hfs of the ground state of Pu II can be attributed to the s electron. Therefore the difference in the hfs ratios between the optical measurement of the lowest level of Pu II and the resonance measurements of Pu IV is attributed to the hfs anomaly $^{241}\Delta^{239}$ of the s electron [8]. The hfs splittings of the excited states of Pu I and Pu II listed in Table I also are probably due to s and $p_{1/2}$ electrons but since many close lying configurations interact further discussion is dependent upon the detailed analysis of the optical spectra [6]. Varying the percentage s electron contribution to the total hfs constant will cause changes in the hfs ratio of ^{241}Pu to ^{239}Pu. This may be the reason for the discrepancy in the ratio between Pu VII and Pu IV.

Since the s electron contribution to A(Pu IV) is small

$$\frac{A(^{241}\text{Pu IV})}{A(^{239}\text{Pu IV})} \simeq \frac{g_I(^{241}\text{Pu})}{g_I(^{239}\text{Pu})} \quad . \tag{1}$$

Then the hfs anomaly is

$$^{241}\Delta^{239} = \frac{A(^{241}\text{Pu II})}{A(^{239}\text{Pu II})} \times \frac{A(^{239}\text{Pu IV})}{A(^{241}\text{Pu IV})} - 1 = -6.3 \pm 0.2\% \quad . \tag{2}$$

The Bohr-Weisskopf theory [8] allows an estimate of $^{241}\Delta^{239}$ assuming $g_s = g$ (neutron), $g_\ell = \frac{Z}{A}$, and the coefficient b ≃ 5.5% extrapolated from their table. Then

$$^{241}\Delta^{239} \simeq 0.3b \frac{g_s g_\ell}{g_s - g_\ell} \left(\frac{1}{g_I(^{241}\text{Pu})} - \frac{1}{g_I(^{239}\text{Pu})} \right) = -5.4\% \quad . \tag{3}$$

The agreement between the theory and experiment is surprisingly good, which was not true for the stable odd-neutron isotopes of Yb(Z = 70, A = 101,103) [9].

A better value for μ_N of ^{241}Pu may be calculated from the hfs ratio, Eq. (1), and $\mu_N(^{239}\text{Pu}) = +0.200 \pm 0.004$ nm [10]. The new value for $\mu_N(^{241}\text{Pu}) = -0.718 \pm 0.017$ nm where the sign is obtained from the optical data.

I wish to thank J. Conway and M. Fred for many valuable discussions on the optical spectra of Pu I and Pu II and W. Kolbe and G. Kaindl for their help and advice.

References

1. S. Gerstenkorn and F. S. Tomkins, Physica 42 (1969) 581.
2. B. Bleaney, P. M. Llewellyn, M. H. L. Pryce, and G. R. Hall, Phil. Mag. 45 (1954) 991.
3. W. Kolbe and N. Edelstein, (unpublished results).
4. N. Edelstein and R. Mehlhorn, Phys. Rev. B (Sept. 1, 1970).
5. B. R. Judd, Phys. Rev. 125 (1962) 613.
6. M. Fred, (private communication).
7. J. Bauche and B. R. Judd, Proc. Phys. Soc. (London) 83 (1964) 145.
8. A. Bohr and V. F. Weisskopf, Phys. Rev. 77 (1950) 94.
9. B. Budick and J. Snir, Phys. Letters 24B (1967) 276.
10. J. Faust, R. Marrus, and W. A. Nierenberg, Phys. Letters 16 (1965) 71.

Table I. Hfs constants for various levels of Pu ions.

Ion	Level (cm^{-1})	A (10^{-3} cm^{-1})	$\frac{A(^{241}Pu)I(^{241}Pu)}{A(^{239}Pu)I(^{239}Pu)}$	Reference
^{241}Pu I	23766	-58.25 ± 0.2	-3.367 ± 0.02	1
^{239}Pu I	23766	86.5 ± 0.2		
^{241}Pu II	0	-167.3 ± 0.5	-3.362 ± 0.02	1
^{239}Pu II	0	248.8 ± 0.5		
^{241}Pu II	22038	- 32.7 ± 1	- 3.41 ± 0.2	1
^{239}Pu II	22038	48 ± 1		
^{241}Pu IV	0	|4.807| ± 0.010	|3.590| ± 0.010	3
^{239}Pu IV	0	|6.695| ± 0.003		
^{241}Pu VII	0	|245| ± 1 gauss	|3.53| ± 0.02	2
^{239}Pu VII	0	|347| ± 1 gauss		

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

TECHNICAL INFORMATION DIVISION
LAWRENCE RADIATION LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720