Submitted to the Journal "Nuovo Cimento"

UCRL-20087
Preprint

c. 2

COMMENTS ON THE BROKEN EXCHANGE-DEGENERATE REGGE POLE MODEL FOR np AND $p\bar{p}$ CHARGE EXCHANGE

RECEIVED
LAWRENCE
RADIATION LABORATORY

SEP 30 1970

LIBRARY AND
DOCUMENTS SECTION

B. K. Pal

August 1970

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

UCRL-20087

LAWRENCE RADIATION LABORATORY
UNIVERSITY of CALIFORNIA BERKELEY

3 c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

COMMENTS ON THE BROKEN EXCHANGE-DEGENERATE REGGE POLE MODEL FOR np AND $p\bar{p}$ CHARGE EXCHANGE*

B. K. Pal

Lawrence Radiation Laboratory
University of California
Berkeley, California 94720

August 1970

ABSTRACT

We point out that formulas given by Pal on np and $p\bar{p}$ CEX for exchange-degenerate and broken exchange-degenerate Regge pole models can be better rationalized by considering π as simultaneously an M = 0 singlet and an M = 1 parity doublet and satisfying threshold behavior.

We have seen that np and $p\bar{p}$ CEX (charge-exchange) and charged-pion photoproduction differential cross section produce sharp forward peaks.[1] The pion conspiracy was popular and attractively economical in fitting those processes, it certainly is not unique,[2] and it causes difficulties in other processes via factorization.[3] We want to show here that the factorization difficulties resulting from the conspiracy hypothesis can be avoided if we take into consideration the pion as an M = 0 singlet and an M = 1 parity doublet simultaneously. The M = 0 pion is an evasive one with the usual pion-nucleon coupling constant at the pion pole, and an M = 1 is a parity doublet satisfying the conspiracy relation. An M = 0 pion singlet and an M = 1 parity doublet were considered

simultaneously in np CEX.[4] Only an M = 1 pion parity doublet was considered by Pal[5] in the np and $p\bar{p}$ CEX broken exchange-degenerate Regge pole model. Now an M = 0 evasive pion should occur in invariant amplitudes $G_i(s,t,u)_{t=0,1}$[6] in the following manner:

$$G(s,t,u)_{T=0,1} = G_2(s,t,w)_{T=0,1} = G_5(s,t,w)_{T=0,1}$$
$$= -\frac{g^2}{\pi}\alpha'(t)\,\Gamma[-\alpha(t)]\,(1+e^{-i\pi\alpha(t)})\,(-\tfrac{3}{2},\tfrac{1}{2})\,(\frac{s}{s_0})^{\alpha(t)},$$
$$G_3(s,t,u)_{T=0,1} = G_4(s,t,u)_{T=0,1} \tag{1}$$
$$= \frac{g^2}{\pi}\alpha'(t)\,\Gamma[-\alpha(t)]\,(1+e^{-i\pi\alpha(t)})\,(-\tfrac{3}{2},\tfrac{1}{2})\,(\frac{s}{s_0})^{\alpha(t)}$$

for the pion exchange in the t channel, and therefore,

$$\phi_2(s,t)_{T=0,1} = \phi_4(s,t)_{T=0,1}$$
$$= -\frac{t}{2\sqrt{s}}\frac{g^2}{\pi}\alpha'(t)\,\Gamma[-\alpha(t)]\,(1+e^{-i\pi\alpha(t)})\,(-\tfrac{3}{2},\tfrac{1}{2})\,(\frac{s}{s_0})^{\alpha(t)} \tag{2}$$

and all other $\phi_i(s,t)$ are zero. For the pion exchange in the u-channel,

$$G_1(s,u,t)_{T=0,1} = G_4(s,u,t)_{T=0,1} = G_5(s,u,t)_{T=0,1}$$
$$= -\frac{g^2}{\pi}\alpha'(u)\Gamma[-\alpha(u)]\,(1+e^{-i\pi\alpha(u)})\,(\tfrac{3}{2},\tfrac{1}{2})\,(\frac{s}{s_0})^{\alpha(u)} \tag{3}$$

$$G_2(s,u,t)_{T=0,1} = G_3(s,u,t)_{T=0,1}$$

$$= \frac{g^2}{\pi}\,\alpha'(u)\Gamma[-\alpha(u)]\,(1+e^{-i\pi\alpha(u)})\,(\frac{s}{s_0})^{\alpha(u)}\,(\frac{3}{2},\frac{1}{2}) \quad , \tag{4}$$

and therefore

$$\phi_2(s,u)_{T=0,1} = -\frac{u}{2\sqrt{s}}\,\frac{g^2}{\pi}\,\alpha'(u)\Gamma[-\alpha(u)]\,(1+e^{-i\pi\alpha(u)})\,(\frac{s}{s_0})^{\alpha(u)}\,(\frac{s}{s_0})^{\alpha(u)}\,(\frac{3}{2},\frac{1}{2}),$$

$$\phi_3(s,u)_{T=0,1} = \frac{u}{2\sqrt{s}}\,\frac{g^2}{\pi}\,\alpha'(u)\,\Gamma[-\alpha(u)]\,(1+e^{-i\pi\alpha(u)})\,(\frac{s}{s_0})^{\alpha(u)}\,(\frac{3}{2},\frac{1}{2}) \quad , \tag{5}$$

and all other $\phi_i(s,u)$ are zero.

In Ref. 5, the residues were not properly rationalized, and therefore they did not satisfy correct threshold behavior for $\phi_5(s,t,u)$ only. Choosing the same invariant amplitudes in Ref. 5 for the t-channel pion exchange, one has conspiracy π_c and its exchange degenerate B_c

$$\phi_5(s,t)_{T=0,1} \underset{\text{fixed } t}{\overset{\text{large } s}{\sim}} -(-\frac{3}{2},\frac{1}{2})\,\frac{\beta(-t)^{1/2}(s+t-4m^2)^{1/2}}{m}\,\frac{(bs)^{\alpha(t)-1}}{\alpha(0)\Gamma[\alpha(t)]^{\sin\pi\alpha(t)}} \tag{6}$$

instead of

$$\phi_5(s,t)_{T=0,1} \underset{\text{fixed } t}{\overset{\text{large } s}{\sim}} -(-\frac{3}{2},\frac{1}{2})\,\beta_t(1+\frac{s}{m^2})^{1/2}\,\frac{m(bs)^{\alpha(t)-1}}{\sin\pi\alpha(t)\,\alpha(0)\Gamma[\alpha(t)]}$$

In Ref. 5, it does not satisfy correct threshold behavior[7] $\propto (-t)^{1/2}$ for small t. For the u-channel pion exchange, one has

$$\phi_5(s,u)_{T=0,1} \underset{\text{fixed } u}{\overset{\text{large } s}{\sim}} -\left(\frac{3}{2}, \frac{1}{2}\right) \beta(-u)^{1/2} \frac{(s+u-4m^2)^{1/2}}{m} \frac{(bs)^{\alpha(u)-1}}{\alpha(0)\,\Gamma[\alpha(u)]\,\sin\pi\alpha(u)}. \quad (7)$$

It satisfies the correct threshold behavior $\propto (-4)^{1/2}$ for small u.

ACKNOWLEDGMENTS

It is a pleasure to thank Professor G. F. Chew for warm hospitality at the Lawrence Radiation Laboratory.

FOOTNOTE AND REFERENCES

*Work done under auspices of the U. S. Atomic Energy Commission.

1. G. Manning et al., Nuovo Cimento 61A, 167 (1966); A. M. Boyarski et al., Phys. Rev. Letters 20, 300 (1968).
2. K. Dietz and W. Kerth, Phys. Letters 26B, 394 (1968); D. Amati et al., Phys. Letters 26B, 510 (1968).
3. M. Le Bellac, Phys. Letters 25B, 524 (1967); M. Aderholz et al., Phys. Letters 27B, 174 (1968).
4. B. K. Pal, Phys. Rev. 1D, 1405 (1970).
5. B. K. Pal, Venziano Model for np → pn and $p\bar{p} \to n\bar{n}$ Scattering and Parity Doublet, UCRL-19441, 1970, and Neutron-Proton Charge-Exchange Polarization, UCRL-19820, 1970.
6. I. J. Muzinich, Phys. Rev. 130, 1571 (1963).

7. M. L. Goldberger, M. T. Grisaru, S. W. MacDowell, and D. Y. Wong, Phys. Rev. 120, 2250 (1960).

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

TECHNICAL INFORMATION DIVISION
LAWRENCE RADIATION LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720