Submitted to the Logistics Review

UCRL-20137
Preprint

c. 2

LIBRARY AND
DOCUMENTS SECTION

THE SPACE OF CUBIC SPLINES WITH SPECIFIED KNOTS

Jonathan D. Young

October 1970

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

UCRL-20137 c. 2

LAWRENCE RADIATION LABORATORY
UNIVERSITY of CALIFORNIA BERKELEY

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UCRL-20137

THE SPACE OF CUBIC SPLINES WITH SPECIFIED KNOTS

Jonathan D. Young

Lawrence Radiation Laboratory
University of California
Berkeley, California

October 1970

ABSTRACT

For a given closed finite interval, which has been partitioned into a finite number of subintervals, we consider all cubic splines[1] which have knots at the endpoints and at each partition point. A finite basic subset may be readily chosen such that any spline in the parent set can be expressed as a unique linear combination of the basic splines.

Any function defined on the interval whose values at the knots and whose first derivatives at the endpoints are known can be approximated by a unique cubic spline, hence as a linear combination of basic splines.

Where a relatively large number of functions are to be approximated, it becomes practical to first compute the basic splines and then determine the required linear combination to approximate each of the functions.

INTRODUCTION

For a given interval, $[a,b]$ with partitioning:

$$a = x_1 < x_2 < \text{----} \; x_n = b,$$

a cubic spline, s (x), defined on $[a,b]$ with knots at the x_i has the following properties:

(I) The function, s, is a polynomial of degree three (or less) in x on each subinterval,

$$[x_i, x_{i+1}]; \; i = 1, n-1$$

(II) The function s has a continuous second derivative on the whole interval $[a,b]$.

Obvious examples of cubic splines are:

$$s = \text{constant (including zero)}$$

$$s = x$$

$$s = x^2$$

$$s = x^3$$

and any linear combination of the above. However these splines do more than meet the requirements (I), (II), they have constant (hence continuous) third derivatives over the whole interval. In general, cubic splines with knots at x_i may have finite discontinuities in the third derivative at interior knots. As an example consider:

$$s = 0 \quad x \le x_2$$

$$s = (x-x_2)^3 \quad x \ge x_2.$$

Any function f (x) defined on $[a,b]$ for which values at the x_i are known and values of its first derivative at a and b are known can be readily approximated by a unique cubic spline with knots at the x_i. The cubic spline fit can be computed directly for the function[1] or where several functions are to be fitted

a basic set of splines can be determined and the fitting splines be computed for each function as a unique linear combination of the basis.

Direct Cubic Spline Fit. Suppose we are given a set of points,

$$\left\{x_i, f_i\right\}; \; i = 1, n$$

with the assumption that there is a function , $f(x)$, defined on x_i, x_n such that

$$f(x_i) = f_i \quad ; \quad i = 1, n$$

with specified terminal derivatives

$$f'(x_1) = f'_1$$

and

$$f'(x_n) = f'_n .$$

The cubic spline s which approximates f on $[a,b]$ must be an exact fit of the above data, i.e.

$$s_i = f_i \quad ; i = 1, n$$
$$s'_1 = f'_1$$
$$s'_n = f'_n$$

Th$^{\underline{m}}$ 1 There is a unique s which fits f.

Proof:

(a) Existence

From properties (I) and (II) we have

$$s'_{i-1} + 4s'_i + s'_{i+1} = 3\left[\frac{(x_i - x_{i-1})(s_{i+1} - s_i)}{x_{i+1} - x_i} + \frac{(x_{i+1} - x_i)(s_i - s_{i-1})}{x_i - x_{i-1}}\right]$$

for $i = 2, n - 1$

In terms of the fitting requirements

$$4s'_2 + s'_3 = 3\left[\frac{(x_2 - x_1)(f_3 - f_2)}{x_3 - x_2} + \frac{(x_3 - x_2)(f_2 - f_1)}{x_2 - x_1}\right] - f_1'$$

$$s'_{i-1} + 4s'_i + s'_{i+1} = 3\left[\frac{(x_i - x_{i-1})(f_{i+1} - f_i)}{x_{i+1} - x_i} + \frac{(x_{i+1} - x_i)(f_i - f_{i-1})}{x_i - x_{i-1}}\right]$$

$$s'_{n-2} + 4s'_{n-1} = 3\left[\frac{(x_{n-1} - x_{n-2})(f_n - f_{n-1})}{x_n - x_{n-1}} + \frac{(x_n - x_{n-1})(f_{n-1} - f_{n-2})}{x_{n-1} - x_{n-2}}\right] - f'_n$$

The above linear system of n-2 equations is tridiagonal with diagonal dominance, hence has a unique solution for s'_i ; $i = 2, n-1$.

(b) Uniqueness

On any subinterval $[x_i, x_{i+1}]$; $i = 1, n-1$ the values of s_i and s_{i+1} are determined by

$$s_i = f_i \quad \text{and} \quad s_{i+1} = f_{i+1}$$

and the values of s'_i and s'_{i+1} are uniquely determined by the linear system above. These values uniquely determine a cubic segment on $[x_i, x_{i+1}]$. Consequently s is uniquely determined for the whole interval $[a,b]$.

<u>The Cubic Spline Space.</u> Let S be the set of all cubic splines with knots at x_i ; $i = 1, n$. We have already shown the S is not empty. In fact it contains as a subset the linear space of all polynomials of degree three or less[2]. With the obvious definitions of addition

$$u = s + t \quad : \quad u(x) = s(x) + t(x) \quad ; \quad s,t \in S$$

and scalar multiplication

$$v = as \qquad v(x) = a\, s(x), \qquad s \in S$$

it may be readily shown that S is itself a linear space.

Since S contains a linear space of dimension 4 , its dimension must be greater than or equal 4. Further since any s in S is completely determined by its n-1 cubic segments, the dimension of S is less than or equal 4(n-1). Note that for n=2, the dimension of S is necessarly 4.

The dimension of S is readily established by the following theorem:

Th$^{\underline{m}}$ 2. Any spline s in S is uniquely determined by the value of s at the x_i and s' at x_1 and x_n.

Proof: (Indirect)

Suppose there are two such splines, s and s* satisfying the specified values. Then both s and s* are spline approximations of s and by Th$^{\underline{m}}$ 1 s = s*.

Since n+2 values are necessary and sufficient to determine any s in S, the dimension of S must be n+2.

We now construct a convenient basis $\{t_j\}$; j = 1, n+2 for S. Let t_j be cubic splines in S such that

$$t_j'(x_1) = 0 \quad t_j'(x_n) = 0 \quad t_j(x_i) = \delta_{ij} \quad \text{for } j = 1, n \; ; \; i = 1, n$$

$$t_{n+1}'(x_1) = 1 \quad t_{n+1}'(x_n) = 0 \quad t_{n+1}(x_i) = 0 \quad \text{for } i = 1, n$$

$$t_{n+2}'(x_1) = 0 \quad t_{n+2}'(x_n) = 1 \quad t_{n+2}(x_i) = 0 \quad \text{for } i = 1, n.$$

It is obvious that no one of the t_j is a linear combination of the others and, further, that for any s in S specified in accordance with Th$^{\underline{m}}$ 2 we have

$$s(x) = \sum_{i=1}^{n} s(x_i)\, t_i(x) + s'(x_i)\, t_{n+1}(x) + s'(x_n)\, t_{n+2}(x)$$

<u>Practical Applications</u> For any function f defined on [a,b] with terminal first derivatives and function values at the x_i known, the basic splines, t_j , may be generated (this consists of computing $t_j'(x_i)$ for i = 2,n-1) then the fitting spline s for f is determined by

$$s'(x_i) = \sum_{j=1}^{n} f'(x_i')t_j'(x_i) + f'(x_1)\, t_{n+1}'(x_i) + f'(x_n)t_{n+2}'(x_i)$$

for $i = 2, n-1$ together with

$$s'(x_1) = f'(x_1) \quad , \quad s'(x_n) = f'(x_n)$$

and $$s(x_i) = f(x_i) \; ; \; i = 1, n.$$

Obviously this method is not practical for fitting one function, f , or even for fitting n+2 such functions since the generation of the basic splines would require as much computation as the direct computation of the fitting splines and the final vector multiplication would be required. If we assume (perhaps conservatively) that each matrix inversion requires approximately n times as much computation as the vector multiplication, then the basic computation is worthwhile when

$$m \geqslant (n^2+2n) \; / \; (n-1)$$

where m is the number of functions to be fitted.

Methods for obtaining numerical solutions to second-order differential equations in the spline space as linear combinations of basic splines will be discussed in subsequent articles.

ACKNOWLEDGEMENT

This work was done in part under the auspices of the Atomic Energy Commission.

REFERENCES

1. Birkhoff, G. and de Boor, C. R., "Piecewise Polynomial Interpolation and Approximation," Approximation of Functions, Henry L. Garabedian, Ed., (Elsevier Publishing Co., Amsterdam, 1965), 164 - 168

2. Halmos, P. R., Finite - Dimensional Vector Spaces, (D. Van Nostrand Co., Princeton, 1958), 1-14.

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

TECHNICAL INFORMATION DIVISION
LAWRENCE RADIATION LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720