UCRL-20622
UC-34 Physics
TID-4500 (57th Ed.)

c. 2

RECEIVED
LAWRENCE
RADIATION LABORATORY

AND
DOCUMENTS SECTION

A2 DUPLICITY: A COMPILATION OF THE AMOUNT OF DIPOLE STRUCTURE OBSERVED IN A2 MESON EXPERIMENTS

Gerald R. Lynch

March 10, 1971

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

LAWRENCE RADIATION LABORATORY
UNIVERSITY of CALIFORNIA BERKELEY

UCRL-20622 c.2

34a

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

A2 DUPLICITY: A COMPILATION OF THE AMOUNT OF DIPOLE STRUCTURE OBSERVED IN A2 MESON EXPERIMENTS*

Gerald R. Lynch

Lawrence Radiation Laboratory
University of California
Berkeley, California 94720

March 10, 1971

In an attempt to find a way to compare various experiments that have looked for splitting in the A2 meson, I have fit these experiments to a distribution of the form

$$\delta \text{ (dipole)} + (1-\delta) \text{ (Breit-Wigner)}.$$

The parameter δ, which I call the duplicity (a name that was suggested to me by Alan Krisch), measures the amount of dipole structure observed in an experiment. A δ greater than 1 means that the data show a dip greater than possible for a dipole smeared out by experimental resolution.

The procedure was first to find the best fit (highest likelihood) to the data for a dipole and also the best fit for a Breit-Wigner, in each case with a linear background under the resonance. (One experiment, the CERN A_2^0 experiment, required a quadratic background). Then with the resonance parameters fixed at their best-fit values, but with the background parameters free, a fit was made to the above form.

The fits were made with the same fitting routines and the same resonance shapes that were used in our π-82 experiment.[5] For some experiments the data were obtained by reading the values off of published graphs, a procedure that is subject to error. (This is especially true for the CERN A_2^0 experiment.)

Perhaps the best way of determining the number of standard deviations of preference an experiment has for one or the other of the models is to quote $S = \sqrt{\Delta(2 \ln L)}$, the square root of twice the difference in the logarithm of the likelihoods for the two fits. The sign of S is arbitrarily defined to be negative if the dipole fit is best. The quantities δ and S, as well as the resolution that I used in the calculations, are tabulated below, and the values of δ are plotted.

I feel that these numbers, δ and S, provide a better method of comparing the results of various experiments than other methods that have been used. In particular they are better than confidence intervals, which are so dependent upon the interval in which the fit was done.

However, these parameters have their drawbacks. The obvious limitation of this parametrization is that it singles out the dipole structure to test, a structure that has little theoretical justification and is chosen only because the missing mass spectrometer data were observed to fit it well. This parametrization is useless in looking for other types of structure.

A more subtle failing of this parametrization is that even though the dipole form has the same number of parameters as does the Breit-Wigner form, it nevertheless has more

flexibility in fitting the data because the mass of the dipole can be adjusted to put the dip in the prediction where a dip happens to occur in the data. In an attempt to get a feeling of the magnitude of this effect, I generated 13 Monte Carlo experiments according to the best Breit-Wigner fit to the BDNPT data and analyzed each of these fake experiments by the method that I have used to analyze the real experiments. The result was that for 9 of the 13 random experiments the dipole fit better than did the Breit-Wigner! Four of the 13 experiments had $S < -2$ and one had $S = -2.7$, the value for the BDNPT experiment. Thus for a relatively low statistics experiment with a 1-to-1 signal-to-noise ratio such as the BDNPT experiment or Crenell et al., the natural bias of preferring a dipole is considerable and an S of -2.7 standard deviations is not as significant as it would be in a well-behaved system. This fact, coupled with the fact that many experiments that have looked at the A2 have not published their data, makes it difficult to estimate how many of the experiments with $|S|$ in the 2.5 to 3 level are significant.

Footnote and References

*Work done under the auspices of the U. S. Atomic Energy Commission.

1. MMS+CBS; G. E. Chikovani et al., Phys. Letters 25B, 44 (1967); H. Benz et al., Phys. Letters 28B, 233 (1968).
2. BNL; D. J. Crennell, U. Karshon, K. W. Lai, J. M. Scarr, and I. O. Skillicorn, Phys. Rev. Letters 20, 1318 (1968).
3. BDNPT; K. Bockman et al., Bonn-Durham-Nijmegen-Paris (E. P.)-Torino Collaboration, Nucl. Phys. B16, 221 (1970).
4. CBS($K\bar{K}$); R. Baud et al., Phys. Letters 31B, 397 (1970).
5. LRL-Group A; M. Alston-Garnjost et al., Phys. Letters 33B, 607 (1970).
6. LRL-TG Group; K. W. Barnham et al., UCRL-20292, to be published.
7. CERN A_2^0; M. Basile et al., Lett. Nuovo Cimento 4, 838 (1970).
8. CERN-Munich; G. Grayer et al., Phys. Letters 34B, 4, 333 (1971).
9. BNL($K\bar{K}$); K. J. Foley et al., Phys. Rev. Letters 26, 413 (1971).
10. M. Aguilar-Benitez et al., CERN-Paris (C. d. F.)-Liverpool Collaboration, Phys. Letters 29B, 62 (1969). This experiment has not been included here because it reports the A2 mass spectrum in mass-squared rather than mass bins.

Experiment	Resolution $\Gamma/2$ (MeV)	Duplicity[a]	$S = \sqrt{\Delta(-2 \ln L)}$
1. MMS + CBS	5.9	1.04 ± 0.20	-6.1
2. BNL	5	1.7 ± 0.6	-2.9
3. BDNPT	5	1.02 ± 0.37	-2.7
4. CBS ($K\bar{K}$)	9.4	1.14 ± 0.36	-3.1
5. LRL-Group A	3.8, 6.7, 9.2	0.17 ± 0.15	5.7
6. LRL-TG Group	7	0.78 ± 0.31	-2.5
7. CERN A_2^0	9.4	1.2 ± 0.5	-2.8
8. CERN-Munich	5.9	0.01 ± 0.13	7.6
9. BNL ($K\bar{K}$)	5.5	0.15 ± 0.14	5.2

[a] Fit was to $dn/dm = \delta(\text{dipole}) + (1-\delta)(\text{Breit-Wigner}) + a + bm$.



XBL713-3046

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE RADIATION LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720