

UCRL 2617

UNCLASSIFIED

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

The Reaction $p+p \rightarrow n^+ + d$ with Polarized Protons*†

UC-2017

F. S. Crawford, Jr. and M. L. Stevenson

Radiation Laboratory, University of California, Berkeley, California

June 17, 1954

Marshak and Messiah¹ have pointed out that if mesons are present in both S and P states in the above reaction, then it is possible to obtain, through interference, a large azimuthal asymmetry in the angular distribution, provided that a polarized proton beam (or target) is used. If a substantial amount of such an asymmetry is indeed observed, then the relative amounts and phases of S and P waves can be estimated. Carvalho, et al.,² looked for an asymmetry in meson production in the course of examining possible asymmetric background in the scattering from hydrogen of 439 Mev polarized protons. They obtained a null result of $Q = .04 \pm .06$.

Following the Rochester technique,³ Chamberlain and co-workers⁴ have obtained an external beam of 315 Mev protons from the Berkeley cyclotron having a polarization⁵ estimated to be $|P| = |2\langle S_y \rangle| = .73$. With this beam, we have made measurements of the left right asymmetry of meson production in the plane perpendicular to the proton polarization, using coincidence detection of the meson and deuteron, and a liquid hydrogen target. Measurements were made at incident proton energies of 317 and 310 Mev. The results of the raw data are $R-L/R+L = 0.16 \pm 0.03$, and 0.23 ± 0.04 at 317 and 310 Mev respectively. R and L refer to the direction of the produced meson, as seen by the incident proton. (The polarized proton beam is produced by a "left" scatter.⁴)

In the notation of Marshak and Messiah, $R-L/R+L = PQ \frac{A \sin \theta}{A + \cos^2 \theta}$.

Here, $A + \cos^2 \theta$ is the angular dependence of the differential cross section in the C.M. system for unpolarized protons, and Q is the interesting quantity, involving the relative amplitudes and phases of the transitions involved. It is apparent that the angular factor becomes unity, independent of A , at $\theta = 90^\circ$. However, measurements were made at 69° because of meson energy loss in the target. We used $A = 0.40$ and 0.49^6 at 317 and 310 Mev, respectively, to average the angular factor. With $|P| = 0.73$ this gives $|Q| = 0.37 \pm .06$ and $0.42 \pm .07$ at 317 and 310 Mev, respectively. Analysis⁷ of the excitation function shows that Q is expected to vary by less than ten percent between 310 and 317 Mev, so that we may conveniently combine these data to obtain an average $|Q| = 0.41 \pm .05$ at an average proton energy of 314 Mev. $(KE_\pi)_{cm} = 11.3$ Mev.

We satisfied ourselves as to the following points: (1) Uniformity of counter sensitivity. (2) Insensitivity of counters to stray fields. (3) Target and counter alignment. (4) Exclusion of $p+p \rightarrow \pi^+ + p + n$. (5) Reproducibility during many interchanges of counter positions. (6) As an overall check, the experiment was repeated using the ordinary "scattered" beam, which has been shown⁴ to be unpolarized. The beam energy was degraded with a Be absorber to correspond to the measured energy of the polarized beam. At 317 and 310 Mev, $R-L/R+L = -0.02 \pm .06$ and $0.04 \pm .04$ were obtained, respectively. These results have convinced us that no hidden asymmetries were present in the equipment. The average of the left and right measurements with the polarized beam agreed with the results obtained with the unpolarized beam, and with our previous measurements.⁶

Rosenfeld⁷ has compiled data on pion production from nucleons and compared it with the phenomenological theory of Watson and Brueckner.⁸

In his notation, $Q = \sqrt{2} \eta_c \eta (\eta^2 + \eta_c^2)^{-1} \sin(\Psi - \tau_1)$, where $\eta = p_R/m_\pi c$, and η_c is obtained by fitting the 90° (CM) excitation function data to $\ln d\sigma_{10}(90^\circ)/d\Omega = a_{10}(\eta + \eta^3 \eta_c^{-2})$. The terms in η and η^3 come from S and P mesons, respectively.

$\frac{S_0}{S_2} + \sqrt{\frac{1}{2}} = \left| \frac{S_0}{S_2} + \sqrt{\frac{1}{2}} \right| \exp i\Psi$, and $S_1/S_2 = |S_1/S_2| \exp i\tau_1$, where S_0 and S_2 are the complex transition amplitudes giving p state meson production from 1S_0 and 1D_2 protons, respectively, and S_1 is the amplitude for S state production from 3P_1 protons. Our data⁶ yields $\eta_c = 0.62 \pm .16$. With the measured Q , at $\eta = 0.41$, this gives $|\sin(\Psi - \tau_1)| = 0.61 \pm .10$.

The authors wish to thank Dr. Hugh Bradner, Mr. Robert E. Donaldson, Dr. ~~Melvin H. Good~~ ^{Myron L. Good}, Mr. H. Brook Knowles, Mr. R. Tripp, and Mr. T. Ypsilantis for assistance during the measurements.

* This work has been performed under the auspices of the U.S. Atomic Energy Commission.

† This work was reported as a post deadline paper at the Minneapolis meeting of the American Physical Society, June 28-30, 1954. (Bull. Am. Phys. Soc. 29, No. 5)

1. R. E. Marshak and A. M. L. Messiah, *Il Nuovo Cimento*, 11, 337 (1954).
2. Carvalho, Heiberg, Marshall, and Marshall, *Phys. Rev.* (in press).
3. Oxley, Cartwright, Rouvina, Baskir, Klein, King, and Skillman, *Phys. Rev.* 91, 419 (1953).
4. O. Chamberlain, E. Segre, R. Tripp, C. Wiegand, and T. Ypsilantis, *Phys. Rev.* 93, 1430 (1954)
5. O. Chamberlain, E. Segre, R. Tripp, C. Wiegand, and T. Ypsilantis, *Phys. Rev.* (in press).

6. F. S. Crawford, Jr. and M. L. Stevenson, UCRL 2187, UCRL 2188, and to be published.
7. A. H. Rosenfeld - "Production of Pions in Nucleon-Nucleon Collisions at Cyclotron Energies", to be published.
8. K. M. Watson and K. A. Brueckner, Phys. Rev. 83, 1 (1951).