UCRL 2621

UNCLASSIFIED

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UCRL-2621

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

EMISSION OF HIGH-ENERGY GAMMA LINE FROM C^{12}

David Cohen, Burton J. Moyer, Harlan Shaw, and Charles Waddell

June 11, 1954

Berkeley, California

EMISSION OF HIGH-ENERGY GAMMA LINE FROM C^{12}

David Cohen, Burton J. Moyer, Harlan Shaw, and Charles Waddell

Radiation Laboratory, Department of Physics,
University of California, Berkeley, California

June 11, 1954

In connection with studies of proton bremsstrahlung, in which a 180° pair spectrometer is used to analyze the photon spectrum, a gamma line at 15.2 ± 0.2 Mev has been observed which is believed to arise from the excited C^{12} nucleus. Such a line was suggested in R. Wilson's[1] experimental bremsstrahlung data on proton bombardment of carbon, but his methods did not permit resolution of the line.

Figure 1 shows this gamma emission revealed by the pair spectrometer as it appears superimposed upon the bremsstrahlung, viewed at 90° from the beam direction, in the bombardment of carbon by 340-Mev protons in the Berkeley cyclotron. The cyclotron target could be pushed in to a radius where the proton energy was 30 Mev; and at this energy the bremsstrahlung is negligible, leaving the line emission as shown in Fig. 2.

The breadth of the line as here shown is essentially entirely instrumental, and a more precise evaluation of its true energy awaits calculation of the instrumental line shape. Calibration of the spectrometer was accomplished by the floating-wire technique, and a calibration test of the spectrometer upon the 17.6-Mev gamma line from Be^{8*} indicates that the energy error is within the limits given above.

The assignment of the line to C^{12*} is based upon the following facts.

A. It is present with a carbon target for proton energies ranging from 30 to 340 Mev, and is not observed in proton bombardment of Be, B and O.

B. It has been observed in deuteron bombardment of B^{11} in which deuterons of 18, 30, and 50 Mev were incident on a thick target. It is not observed in the deuteron bombardment of B^{10}. Recent work[2] at Indiana gives definite indication of high-energy (about 15-Mev) gamma emission from 11-Mev deuteron bombardment of a thick boron target (presumably natural boron which is 81 percent B^{11}).

C. A level below 15.95 Mev in C^{12} cannot decay by neutron or proton emission, as is made evident by Fig. 3. Consequently such a level must either disintegrate into three alpha particles or into Be^8 plus an alpha, or decay by gamma emission. If the gamma process is to predominate, a selection rule must operate which suppresses the alpha processes in the decay from this level.

Absolute values of the differential cross section for producing these gamma photons into unit solid angle at 90° were roughly determined by a thermocouple monitoring of target temperature. This is related to beam energy loss in the target, and thus, at a known energy, to beam traversal of the target. The absolute efficiency of the spectrometer is calculated from established pair-production cross sections and geometry. In Fig. 4 is illustrated the dependence of the yield of the level in question upon energy of bombarding protons, with a carbon target.

It seems reasonable to propose tentatively[3] that the alpha disintegration may be considerably forbidden by isotopic spin considerations, which would be the case if this C^{12*} level were a T = 1 state. This allows dipole gamma emission in the transition to the C^{12} ground state.[4] Upon this basis the B^{12}, C^{12*}, and N^{12} would form the isotopic triplet. The 15.09-Mev C^{12*} level listed by Ajzenberg and Lauritsen[5] may be the one here in question.

Attempts to produce this gamma emission by the alpha bombardment of Be^9, employing 170-Mev alphas upon a Be target of sufficient thickness to stop them, have not given a detectable yield. This apparent failure to produce the required state by Be^9 (α, η) C^{12*} may be understood in terms of the T = 1 proposal advanced above; since a process which amounts essentially to the coalescence of three alpha particles, with the release of the loosely-bound neutron, could not develop a T = 1 state.

Helpful discussions with Professor M. A. Ruderman are acknowledged.

This work was done under the auspices of the Atomic Energy Commission.

REFERENCES

1. Richard Wilson, Phys. Rev. 85, 568 (1952).
2. V. K. Rasmussen, private communication.
3. This suggestion has come to us by private communication from the group at the California Institute of Technology.
4. L. Radicati, Phys. Rev. 87, 521 (1952).
5. F. Ajzenberg and T. Lauritsen, Revs. Modern Phys. 24, 321 (1952).

FIGURE CAPTIONS

Fig. 1. Photon spectrum from 340-Mev proton bombardment of carbon.

Fig. 2. Photon spectrum from 30-Mev proton bombardment of carbon.

Fig. 3. Energy level diagram.

Fig. 4. Yield of the C^{12} state in question vs. proton energy.



MU-7736

22408-2



MU-7735

22407-2

ENERGY LEVEL DIAGRAM



MU-7733



MU-7741

22413-2