UCRL 3869

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UCRL-3869

UNIVERSITY OF CALIFORNIA

Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

DISPERSION RELATIONS FOR SCATTERING IN THE PRESENCE OF A COULOMB FIELD

Geoffrey F. Chew and H. Pierre Noyes

July 10, 1957

Printed for the U.S. Atomic Energy Commission

UCRL-3869

DISPERSION RELATIONS FOR SCATTERING IN THE PRESENCE OF A COULOMB FIELD*

Geoffrey F. Chew

Radiation Laboratory and Department of Physics
University of California, Berkeley, California

and

H. Pierre Noyes

Radiation Laboratory
University of California
Livermore, California

July 10, 1957

ABSTRACT

A dispersion relation is derived for short-range potential scattering in the presence of a Coulomb field, and on the basis of this result a conjecture is made as to the Coulomb modifications of the Goldberger relations for pion-nucleon scattering. The new relations contain in addition to Coulomb phase shifts only amplitudes that are directly measurable experimentally, the assumption of charge independence not being required. Estimates are made to show that the Coulomb phases that appear explicitly are of no practical importance.

---

* This work was performed under the auspices of the U.S. Atomic Energy Commission.

DISPERSION RELATIONS FOR SCATTERING IN THE PRESENCE OF A COULOMB FIELD

Geoffrey F. Chew

Radiation Laboratory and Department of Physics
University of California, Berkeley, California

and

H. Pierre Noyes

Radiation Laboratory
University of California
Livermore, California

July 10, 1957

I. INTRODUCTION

1. It has been pointed out by Puppi and Stanghellini[1] that the experimentally determined meson-nucleon forward-scattering amplitudes do not quantitatively satisfy the Goldberger dispersion relations.[2] Since if this discrepancy is real it constitutes the first concrete evidence against the validity of local field theory, the most careful scrutiny of both theory and experiment here is required.

The Goldberger relations are incomplete in that they take account only of strong interactions that satisfy charge independence. The very weak Fermi interactions may be safely ignored, but the electromagnetic interaction, which is only moderately weak, requires a closer study. Agodi, Cini, and Vitale[3] have estimated the corrections due to the production of photons both in the physical and nonphysical region. They find nothing large enough to account for the Bologna discrepancy. In this paper we address ourselves to another possible source of trouble: the Coulomb field.

2. The conventional approach to the Coulomb problem is to analyze the experimental angular distributions so as to extract the so-called "nuclear scattering amplitude," which is defined as the difference between the complete amplitude and the pure Coulomb amplitude. Except for some fairly trivial phases this amplitude is then assumed to be identical with the amplitude one would obtain in the absence of the Coulomb field. Actually it is not identical, and at very low energies the deviation is large.

---

[1] G. Puppi and A. Stanghellini, Nuovo cimento 5, 1257 (1957).

[2] Goldberger, Miyazawa, and Oehme, Phys. Rev. 99, 986 (1955).

[3] Agodi, Cini, and Vitale, Phys. Rev. (to be published).

Experience with the Coulomb effect in nucleon-nucleon scattering suggests that the energy at which important Coulomb corrections to the meson-nucleon interaction appear is sufficiently small that Puppi and Stanghellini were justified in ignoring them. The importance of the Bologna discrepancy is so great, however, that this possibility of trouble, even if small, should be pursued. One way to study the effect we are interested in is to calculate the phase shifts with and without the Coulomb field on the basis of some model for the pion-nucleon interaction. This was the approach used by Noyes.[4] The difficulty here is that no good model exists for the S-wave part of the interaction, which is dominant at low energies, so that the results of such calculations can never be completely convincing.

In this paper a different approach is adopted: We make no attempt to find a "pure" meson-nucleon amplitude to insert into the Goldberger relation, but instead seek a modified dispersion relation involving quantities that are more or less directly measurable. That such a relation should exist seems a priori likely, since the electromagnetic interaction is microscopically causal. However, we do not pretend to derive the relation we shall write down. To do so would require at least the heavy artillery of Bogoliubov[5] and no doubt in addition theorems not yet discovered. We shall instead arrive at our relation on the basis of plausibility arguments, starting with the nonrelativistic potential scattering problem, in which fairly rigorous statements can be made.

3. The form for our conjectured dispersion relations turns out to be identical with that of Goldberger,[2] so long as the latter is expressed in terms of real and imaginary parts of a "forward" scattering amplitude. What we propose to change is the definition of the appropriate amplitude. In particular, if the complete forward amplitude in the presence of the (screened) Coulomb field is decomposed according to angular momentum,*

$$f = \frac{k}{q^2} \sum_{\ell} \left[\ell\, e^{i\delta_{\ell-}} \sin \delta_{\ell-} + (\ell + 1) e^{i\delta_{\ell+}} \sin \delta_{\ell+} \right] , \tag{3.1}$$

---

4 H. P. Noyes, Phys. Rev. 101, 320 (1956).

5 Bogoliubov, Medvedev, and Polivanov, Institute for Advanced Study Notes, Princeton, 1956.

* We do not assume charge independence. The dispersion relations for negative and positive meson scattering depend on crossing symmetry but not on charge independence.

where k is the laboratory wave number of the pion and q the wave number in the barycentric system, and the subscripts $(\ell+)$ and $(\ell-)$ refer to states of total angular momentum $\ell+\frac{1}{2}$ and $\ell-\frac{1}{2}$, respectively, then the partial nuclear phase shifts $\delta_{\ell\pm}^{N}$ may be defined as the difference between the full phase shifts $\delta_{\ell\pm}$ and the corresponding Coulomb phase shifts $\delta_{\ell\pm}^{c}$. We conjecture that the correct "forward" amplitude to use in the Goldberger relation is then

$$f' = \frac{k}{q^2} \sum_{\ell=0} e^{2i\sigma_\ell} \left[\ell e^{i\delta_{\ell-}^{N}} \sin \delta_{\ell-}^{N} + (\ell+1) e^{i\delta_{\ell+}^{N}} \sin \delta_{\ell+}^{N} \right], \tag{3.2}$$

where the Coulomb phase shifts σ_ℓ are sufficiently well given by

$$\sigma_\ell \approx \arg \Gamma(\ell + 1 + i\eta) , \tag{3.3}$$

for $\eta = e^2/v$, with v the laboratory velocity of the pion.

4. The presence of the Coulomb phase shifts in Eq. (3.2) produces more complicated expressions for the real and the imaginary parts of f' than are normally used, the difference being important whenever $2\sigma_\ell$ is comparable in size to the corresponding nuclear phase shifts. From zero up to 5 or 10 Mev pion energy, $2\sigma_\ell$ is of the same order of magnitude or larger than the S-wave nuclear phase shifts, so that this energy region should be studied with care. In particular the zero-kinetic-energy scattering lengths, introduced by Goldberger when he makes subtractions in his equations, must be re-examined.

We shall find it desirable not to use the scattering-length concept at all but to make the necessary subtraction at zero total energy rather than zero kinetic energy, thus avoiding emphasis of a point which from the Coulomb point of view is singular. That is, we propose to use dispersion relations for the positive- and negative-pion forward amplitudes in the form

$$\mathrm{Re}\, f'^{(\pm)}(\nu) = \mp \frac{2f^2}{\nu \mp 1/2M} + \frac{\nu^2}{\pi} P \int_1^\infty \frac{d\nu'}{\nu'^2} \left\{ \frac{\mathrm{Im}\, f'^{(\pm)}(\nu')}{\nu' - \nu} + \frac{\mathrm{Im}\, f'^{(\mp)}(\nu')}{\nu' + \nu} \right\}$$

$$+ c_1 \pm \nu c_2 , \tag{4.1}$$

---

6 U. Haber-Schaim, Phys. Rev. 104, 1113 (1956).

where ν is the laboratory pion energy in units of the pion rest mass and C_1 and C_2 are two constants. The symbol P means principal value. This is essentially the form used by Haber-Schaim[6] to determine f^2, the Yukawa coupling constant; it has the advantage of treating the experimental information on low-energy S-wave scattering on the same basis as the rest of the data. The original Goldberger form can be reached from Eq. (4.1) by straightforward manipulation.

In Part II of this paper the expression (3.2) is derived for potential scattering in the presence of a screened Coulomb field. In Part III plausibility arguments are given for the extension to relativistic field theory, while numerical estimates of Coulomb effects in the dispersion relation (4.1) will occupy Part IV.

II. POTENTIAL SCATTERING

5. It has been shown by N. N. Khuri that the scattering amplitude for all potentials that fall off sufficiently rapidly at large distances and are not too singular at the origin satisfy a simple dispersion relation.[7] Consequently, if we assume that the Coulomb potential vanishes beyond some screening radius r_s, we can immediately say that the forward Coulomb scattering amplitude $f^c(E)$ for like charges satisfies the dispersion relation

$$\mathrm{Re}\, f^c(E) = -\frac{M}{2\pi}\tilde{V}_c + \frac{P}{\pi}\int_0^\infty dE' \frac{\mathrm{Im}\, f^c(E')}{E' - E}, \tag{5.1}$$

where $\tilde{V}_c$ is the volume integral of the potential, E is the relative kinetic energy and M the reduced mass. Similarly, if there is a short-range nuclear interaction present in addition, which gives rise to no bound states, the scattering amplitude for this case obeys

$$\mathrm{Re}\, f(E) = -\frac{M}{2\pi}(\tilde{V}_c + \tilde{V}_N) + \frac{P}{\pi}\int_0^\infty dE' \frac{\mathrm{Im}\, f(E')}{E' - E}, \tag{5.2}$$

where $\tilde{V}_N$ is the volume integral of the added nuclear potential. Therefore we can <u>define</u> a "nuclear" scattering amplitude $f^N \equiv f - f^c$, which (subtracting Eq. (5.1) and (5.2)) obeys

---

[7] N. N. Khuri, Phys. Rev. (to be published).

$$\mathrm{Re}\, f^N(E) = -\frac{M}{2\pi}\,\tilde{V}_N + \frac{P}{\pi}\int_0^\infty dE'\,\frac{\mathrm{Im}\, f^N(E')}{E'-E} \quad . \tag{5.3}$$

If the charges are of opposite sign, there will be a sum of terms $R_j^c/(E_j^c - E)$, where E_j^c are the bound-state energies of the (screened) Coulomb field, appearing in both Eqs. (5.1) and (5.2), but if the nuclear potential introduces no appreciable level shifts in these states, the poles do not appear in Eq. (5.3). One may perhaps worry that the coefficients R_j^c may be changed by the presence of a short-range interaction even though the energy levels are not. It can be shown, however, that these coefficients are even less sensitive to modifications of the potential at short distances than are the binding energies.

We can also define "nuclear" phase shifts by $\delta_\ell^N = \delta_\ell - \delta_\ell^c$, and hence

$$\begin{aligned} f^N(E) &= \sum_\ell \frac{(2\ell+1)}{2ik}\left\{(e^{2i\delta_\ell} - 1) - (e^{2i\delta_\ell^c} - 1)\right\} \\ &= \sum \frac{(2\ell+1)}{k}\, e^{2i\delta_\ell^c}\, e^{i\delta_\ell^N}\, \sin\delta_\ell^N \quad , \end{aligned} \tag{5.4}$$

where k is the wave number. Note that instead of the usual identification

$$\mathrm{Im}\, f(E) = \sum (2\ell+1)\,\frac{\sin^2\delta_\ell}{k} = \frac{k}{2\pi}\,\sigma_{\mathrm{total}}(E) \quad , \tag{5.5}$$

we have

$$\begin{aligned} \mathrm{Im}\, f^N(E) &= \sum (2\ell+1)\,\frac{\sin(\delta_\ell^N + 2\delta_\ell^c)\,\sin\delta_\ell^N}{k} \\ &= \frac{k}{4\pi}\int d\Omega\left\{\frac{d\sigma(\theta)}{d\Omega} - \frac{d\sigma_c(\theta)}{d\Omega}\right\} \quad . \end{aligned} \tag{5.6}$$

At first sight the last line, which tells us to remove the Coulomb scattering from the total cross section (but to leave the Coulomb-nuclear interference), looks like a simple prescription to apply. It can be used, however, only if the difference $\frac{d\sigma}{d\Omega} - \frac{d\sigma_c}{d\Omega}$ can be extrapolated to the forward direction without ambiguity; in practice this means that except at high energy the alternate expression in terms of phase shifts must be used. Thus we need to know the Coulomb phase shifts $\delta_\ell{}^c$ explicitly to correct both the real part of the forward scattering amplitude and the imaginary part.

Provided that we are at a high enough energy to have $kr_s >> 1$, and do not have to analyze for partial waves such that $\ell \gtrsim kr_s$, the Coulomb phase shifts are

$$\delta_\ell{}^c = \sigma_\ell - \eta \log 2kr_s \quad , \tag{5.7}$$

where

$$\eta = \pm e^2 M/k \qquad \text{and} \qquad \sigma_\ell = \arg \Gamma(1 + \ell + i\eta). \tag{5.8}$$

Were we interested in energies or angles such that these conditions would be violated, we would have to explicitly evaluate $\delta_\ell{}^c$ for the charge distribution under consideration and use the dispersion relation (5.3) together with (5.4). However, such is not the case in the present application to pion-nucleon scattering, and all explicit reference to the screening may be removed, as we now show.

6. Note that at the energies and for the partial waves of interest the effect of the screening is simply to multiply the scattering amplitude f^N by an energy-dependent phase factor, and that this factor--although it has an essential singularity at the origin--is analytic in the upper half of the complex k plane and approaches unity for large k. The behavior at the origin is physically incorrect, since for low enough energy the S phase must go as $a(r_s)k$, where $a(r_s)$ is the scattering length for our screened Coulomb field. Consequently, it can do no harm to displace the essential singularity below the origin by an amount $k_o \sim 1/r_s$. Then the function $\exp(2iS)(f^N - \tilde{V}_N)$, with

$$S = e^2 M \frac{\log 2(k + ik_o)r_s}{k + ik_o} \quad , \tag{6.1}$$

is analytic in the upper half plane and goes to zero as k becomes large. Hence we have

$$\mathrm{Re}\, e^{2iS(E)}(f^N(E) - \tilde{V}_N) = \frac{P}{\pi}\int_0^\infty dE' \frac{\mathrm{Im}\, e^{2iS(E')}(f^N(E') - \tilde{V}_N)}{E' - E} . \tag{6.2}$$

But $e^{2iS} - 1$ satisfies the same conditions, so that one obtains

$$\mathrm{Re}(e^{2iS(E)} - 1)\tilde{V}_N = \frac{P}{\pi}\int dE' \frac{\mathrm{Im}(e^{2iS(E')} - 1)\tilde{V}_N}{E' - E} \tag{6.3}$$

and by subtracting Eq. (6.3) from (6.2) we see that the dispersion relation (5.3) is also satisfied by the quantity

$$f' = e^{2iS} f^N = \sum_\ell (2\ell + 1) e^{2i\sigma_\ell} e^{i\delta_\ell^N} \sin \delta_\ell^N , \tag{6.4}$$

where σ_ℓ is the usual Coulomb phase for $\ell << kr_s$ and otherwise is $\delta_\ell^c + S$.

Since the Klein-Gordon equation leads to the same radial equation as the Schroedinger equation, if we interpret η as e^2/v and ℓ in the centrifugal term as $((\ell + \frac{1}{2})^2 - e^4)^{\frac{1}{2}} - \frac{1}{2}$ (and the nuclear interaction is a world scalar), we can immediately extend our results to this case, if these modifications cause no difficulty. The modification of σ_ℓ is clearly trivial, but the phase factor no longer goes to zero for large k. However if S is multiplied by $iK/(k + iK)$, where K is much larger than any wave number of interest, we have restored this property without destroying the analyticity in the upper half plane, and the proof still stands.

III. EXTENSION TO RELATIVISTIC FIELD THEORY

7. In the absence of electromagnetic interaction the dispersion relations for potential scattering bear a striking resemblance to the Goldberger relations for pion-nucleon scattering in local field theory. It is this resemblance which emboldens us to make a conjecture as to the probable Coulomb modifications of the Goldberger relations.

In the potential problem the fundamental relation between the real and imaginary parts of the forward amplitude was not altered by Coulomb effects, so that we conjecture first of all that the same will be true in the relativistic field theoretical problem. It is hard to think of possible modifications that could occur in the physical region, $\nu > \mu$; and the crossing relation,[8] which tells us what to do for $\nu < -\mu$, remains valid in the presence of an electromagnetic field. In the nonphysical region between $-\mu$ and $+\mu$ the electromagnetic interaction will of course have effects, but we suppose these to be accounted for as follows: (a) Contributions from states containing one or more photons should be classified as radiative rather than Coulomb corrections.[3] (b) In the case of π^-, p scattering there are Coulomb bound-state contributions, but these should be removed when we take the difference between the full amplitude and the Coulomb amplitude, just as for potential scattering. (c) The form of the single-nucleon-state contribution, except for the over-all coefficient, is determined by kinematical considerations, and electromagnetic effects on the coefficient simply amount to an unobservable renormalization of the Yukawa constant. This single-nucleon contribution may be looked upon as the analogue of the volume integral of the potential, which also was not changed by Coulomb corrections.

8. If one were to attempt a real proof of our conjecture, at least two difficulties would certainly arise: (a) The correct method of introducing screening into a field theory is obscure. (b) The nonphysical region is extended when photons are included, so that the Bogoliubov method of analytical continuation[5] cannot be applied. Nevertheless we think it extremely likely that a dispersion relation does hold for the exact forward scattering, including all interactions, weak and strong. Since it is possible to construct a hypothetical field-theory scattering problem involving particles of pionic and nucleonic mass but having <u>only</u> an electromagnetic interaction, one should, by taking a difference, be able to arrive at the relation (4.1), to be obeyed by the amplitude (3.2). Of course the Coulomb phase shifts are accurately given by Eq. (3.3) only for low pion velocities, but at high velocities they are negligible in any case.

We have not developed a detailed argument to show that the screening phase can be removed from the amplitude in the field theoretical problem, as it was removed in the problem of scattering by a potential.

---

[8] M. L. Goldberger, Phys. Rev. <u>99</u>, 979 (1955).

Here our reasoning is simply that we can see nothing in the field theoretical problem to make the value of the screening radius significant if it is insignificant for potential scattering.

IV. NUMERICAL ESTIMATE OF COULOMB PHASE-SHIFT CORRECTIONS

9. If the argument in Section III is accepted, the only change introduced into the pion-nucleon dispersion relations by the presence of the Coulomb field, outside of tiny corrections associated with the mesic-hydrogen-atom level shifts, is to replace the forward scattering amplitude by

$$f'^{(\pm)} = \frac{k}{q^2} \sum_{\ell} e^{2i\sigma_\ell^{(\pm)}} \left[\ell\, e^{i\delta_{\ell-}^{N(\pm)}} \sin \delta_{\ell-}^{N(\pm)} + (\ell+1) e^{i\delta_{\ell+}^{N(\pm)}} \sin \delta_{\ell+}^{N(\pm)} \right] , \tag{9.1}$$

where the phase shifts for positive pions are real, but those for negative pions must be taken to be complex in order to account for charge-exchange scattering and radiative capture. (The superscripts ($\pm$) distinguish between positive and negative pions.) Here in order to estimate the order of magnitude of the correction we may use the usual charge-independent real phase shifts $\delta_{\ell\pm}^{NI}$, where $I = 1$ or 3,* so that we have

$$\delta_{\ell\pm}^{N(+)} = \delta_{\ell\pm}^{N3} \quad \text{and}$$

$$e^{i\delta_{\ell\pm}^{N(-)}} \sin \delta_{\ell\pm}^{N(-)} \approx \frac{1}{3} e^{i\delta_{\ell\pm}^{N3}} \sin \delta_{\ell\pm}^{N3} + \frac{2}{3} e^{i\delta_{\ell\pm}^{N1}} \sin \delta_{\ell\pm}^{N1} . \tag{9.2}$$

---

* It should be said that we are not at all convinced it is safe to assume charge independence in testing the dispersion relations. However, all we are doing in this section is estimating the order of magnitude of explicit Coulomb phase-shift effects. Any assumption about the nuclear phase shifts that gives them a reasonable size should suffice for this purpose.

We see that for each term of the form $\exp(i\,\delta_\ell)\sin\delta_\ell$ in the uncorrected scattering amplitude, the (additive) correction to the real part is

$$-\,2\sin\sigma_\ell\ \sin(\delta_\ell+\sigma_\ell)\sin\delta_\ell \tag{9.3}$$

and the correction to the imaginary part is

$$2\sin\sigma_\ell\ \cos(\delta_\ell+\sigma_\ell)\sin\delta_\ell\ . \tag{9.4}$$

Evidently these corrections, being proportional to $\sin\sigma_\ell$, are important only when the Coulomb phase becomes substantial in absolute value, that is, at very low energies where the nuclear phases may be approximated by

$$\delta_{\ell\pm}^{NI} \approx a_{\ell\pm}^{I}\, q^{2\ell+1}\ . \tag{9.5}$$

10. Considering first the correction to the real part of the forward scattering amplitude, we find that so long as σ_ℓ itself is also small, the fractional correction is

$$\frac{2\sin\sigma_\ell\,\sin(\delta_\ell+\sigma_\ell)}{\cos\delta_\ell} \approx 2\sigma_\ell\,(a_{\ell\pm}^{I}\,q^{2\ell+1}+\sigma_\ell)\ , \tag{10.1}$$

with

$$\sigma_\ell^{(\pm)} \approx \pm\,(1-\gamma+\sum_{p=1}^{\ell}\frac{1}{p})\,\frac{e^2}{k} \tag{10.2}$$

(using the pion rest mass as our energy unit). Since e^2/k is only 0.05 at 1.5 Mev, not rising to 0.5 until the energy is less than 15 kev, and the scattering amplitude is never measured directly at such a low energy, this approximation in effect can always be used. Further, since the S-wave scattering lengths are ~ -0.1, ~ 0.17 and the largest P-wave scattering length is 0.235, we see that the Coulomb correction to the real part of the amplitude can be safely ignored.

Turning to the imaginary part of the forward scattering amplitude, and using the dispersion relations in the form (4.1), we can again show the correction to be small. From Eq. (9.4), the fractional correction at low energies is

$$\frac{{}^{2}\sigma_{\ell}}{{}^{a}{}^{I}_{\ell\pm}\, q^{2\ell+1}} , \tag{10.3}$$

which becomes comparable to unity for S waves at about 5 Mev and for P waves at about 20 Mev. However, the contribution to the dispersion integral in Eq. (4.1) from the region below 50 Mev is generally less than 1%. Consequently, except for fine details such as finding the precise energy at which the real part of the forward scattering amplitude vanishes, the Coulomb phase-shift effects must be negligible. We have verified this fact by direct calculation.

11. If instead of using Eq. (4.1) we had made our subtraction at zero kinetic energy, that is, had used the dispersion relations in the Goldberger form

$$\mathrm{Re}\, f^{(\pm)}(\nu) = \pm \frac{2k^2 f^2}{\nu \mp 1/2M} \frac{1}{1 - \frac{1}{4M^2}} + \frac{k^2}{\pi} \int_1^\infty \frac{d\nu'}{k'^2} \left\{ \frac{\mathrm{Im}\, f^{(\pm)}(\nu')}{\nu' - \nu} + \frac{\mathrm{Im}\, f^{(\mp)}(\nu')}{\nu' + \nu} \right\} + c_1' \pm \nu c_2' , \tag{11.1}$$

this circumstance would not have been quite so obvious. In fact, because of the k'^2 in the denominator of the dispersion integral of Eq. (11.1), the Coulomb correction appears to be enormous. However, the region of ν' affected is so close to the lower limit that we can clearly take the denominators $\nu' \pm \nu$ outside the integrals as $1 \pm \nu$, and since the integrals are multiplied by $k^2 = \nu^2 - 1$, we see that this Coulomb contribution effectively changes the constants C'_1 and C'_2 to C''_1 and C''_2. Of course, since the Bologna analysis identified these constants with "scattering lengths," one must check to see if the use of the data was correct. To make this check, note that at 2 Mev, for example, the Coulomb correction, Eq. (10.1), to the real part of the forward scattering amplitude is still negligible, while the integral and the f^2 term of Eq. (11.1), being multiplied by k^2, contribute less than 1% to the right-hand side; thus the identification of C''_1 and C''_2 with scattering lengths can indeed be made to the required accuracy. Except for this redefinition of the constants, then, the Coulomb corrections to Eq. (11.1) are no greater than to the form (4.1).

V. CONCLUSION

12. In a sense the conclusions of this paper are negative: We have been unable to find any Coulomb corrections to the pion-nucleon dispersion relations larger than 1% or 2%. However, there is also a positive aspect. If our conjectured modification of the Goldberger relations is correct, one can stop worrying about the difference between a "pure" scattering amplitude, generated only by charge-independent strong interactions, and the actual amplitude that is measured. It is possible to use the measured amplitude directly as a test of microscopic causality. We have of course not derived our proposed modification but, to paraphrase a well known remark,[9] <u>if</u> the result is correct someone should be able to prove it.

It must be emphasized again that we have not shown (nor do we believe) that the failure of charge independence leads to negligible effects. The amplitudes used in the dispersion relations in a convincing test of microscopic causality must be obtained from experiment without the assumption of isotopic spin conservation. Puppi and Stanghellini[1] avoided the use of charge independence to a considerable extent but not completely.* It remains to be seen whether the Bologna discrepancy will persist when an analysis entirely free from the charge-independence assumption is carried out.

---

[9] M. L. Goldberger, private but widely circulated communication. The statement, made in reference to the original dispersion relations before the work of Bogoliubov[5] was, "<u>Since</u> the result is correct it can be proved."

* For example, they used the Orear scattering lengths (J. Orear, Nuovo cimento <u>4</u>, 856 (1956)), which depend on charge exchange as well as elastic scattering measurements.