UCRL 443
Copy 2

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

TWO-WEEK LOAN COPY

This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545

BERKELEY, CALIFORNIA

Copy 2

UCRL-443

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

UNCLASSIFIED

RANGE ENERGY RELATION FOR PROTONS IN NUCLEAR EMULSIONS

Hugh Bradner, Frances M. Smith, Walter H. Barkas, A. S. Bishop

September 9, 1949

Berkeley, California

Range Energy Relation for Protons in Nuclear Emulsions

Hugh Bradner, Frances H. Smith, Walter H. Barkas*, A. S. Bishop

Radiation Laboratory
University of California
Berkeley, California

September 9, 1949

ABSTRACT

An experimental range-energy relation in Ilford C-2 emulsion has been obtained for protons up to 39.5 Mev. In the region from 17 to 33 Mev the relation for dry emulsion is fitted by the empirical equation $E_{(Mev)} = 0.251\, R_{(\mu)}^{0.581}$. Variations in water content due to changes in atmospheric humidity make several percent difference in range. The range in Ilford glass is found to be 18 ± 4 percent longer than in dry C-2 emulsion.

*Office of Naval Research, San Francisco

Introduction

The ranges of protons of energies up to 13.1 Mev in nuclear track emulsions were measured by Lattes, Fowler and Cuer[1], with 50μ thick Ilford

[1] C. M. G. Lattes, P. H. Fowler, and P. Cuer, Proc. Phys. Soc. 59, 883 (1947).

B1 emulsion. Extrapolations[2] of those data were used by Bishop[3] to calculate

[2] V. Camerini and C. M. G. Lattes, Ilford technical data, revised March 27, 1948.

[3] A. S. Bishop, Phys. Rev. 75, 1468 (1949).

masses of mesons whose range in emulsion and initial momentum are known. The region of interest corresponded to proton ranges of approximately 4000 μ, or 31 Mev energy. For tracks of this length the extrapolation gave energy values accurate to only ±8 percent.

In the present study the ranges of protons up to 39.5 Mev were measured in Ilford nuclear emulsions. C2 plates used in this study are stated by Ilford Ltd. to have identical chemical composition to B1 plates.

Method

Monoenergetic protons were obtained from the circulating beam of the 184-inch Berkeley cyclotron in an arrangement shown in Figure 1. Protons which struck a 1/8" x 1/16" x 3" copper ribbon target at, say, 30 inches from the cyclotron center scattered in all directions with a variety of energies. A C2 plate placed behind a short channel at 80 inches radius recorded those protons which left the target in a backward direction with such an energy that their paths had approximately 25 inches radius of curvature in the cyclotron magnetic field. With an accurate knowledge of the field it was

possible to calculate the energy of a proton entering normal to the edge of the plate. Plates were put at 80 inch radius for all exposures, and the target radius was varied to obtain different proton energies.

Plates were tilted approximately 3° from horizontal so that protons would enter the surface of the emulsion. Distortion of the emulsion upon processing prevents accurate determination of entrance angle if one uses particles coming through the edge of the emulsion. This tilt, plus scattering of the protons in the emulsion made it difficult to find tracks in 200 μ plates which stayed in the emulsion for 7000μ. Hence, the study was not extended beyond ~39 Mev.

Plates were processed in 3:1 D-19 at 68° F for 20 to 25 minutes, followed by shortstop and commercial hardening fixer. Plates suffered 50 ± 5 percent thickness shrinkage in processing, but negligible transverse shrinkage. Track lengths were measured both by eyepiece reticle and by dial indicator attached to the microscope stage. Corrections were made for angle of dip of protons in the emulsion.

Results

Results of the measurements are presented in Figures 2 and 3. It will be noted that the data fit a straight line on log-log plot between 17 and 35 Mev and fit the data of Lattes at 8 Mev. A least squares analysis gives for this portion the relation:

$$E_{(Mev)} = 0.251\ R_{(\mu)}^{0.581} \qquad (1)$$

where the energy E is expressed in Mev and the range R is measured in microns.

The original values predicted by Camerini and Lattes[4] and a

---

[4] Ilford Technical Data - 1947

recent relativistically correct extrapolation made by Aron[5], are indistinguish-

---

[5] W. Aron - Private communication

---

able from the present experimental results up to 35 Mev.

The effect of widely different humidity conditions on plates before exposure is indicated by the 3 points at 33.5 Mev, and the 2 points at 17.6 Mev. It was necessary to evacuate plates for at least 6 minutes before exposure so that moisture conditions were not accurately known. No difference was found between 6 minutes and 6 hours evacuation of 100μ emulsions which had previously come to equilibrium in a normal Berkeley atmosphere (75° F, 60% R.H.) and these plates are therefore called "dry" in the present study. However, the 17.6 Mev point shows difference between 6 minutes evacuation of such a plate and 6 minutes evacuation of a plate which had been exposed to several hours of rainy weather (95% R.H.). Large changes occurred as shown at the 33.5 Mev point, when 200μ plates were given 6 minutes evacuation after 24 hours exposure to 93 percent R.H. at 75° F in a $NH_4H_2PO_4$ constant humidity chamber. The 80 percent humidity point was obtained with a plate which had been maintained for 24 hours over wet NH_4CL.

An approximate value for the range of protons in the glass backing of C2 plates was found by measuring the length of tracks which had travelled most of their range in the glass. Protons from the monoenergetic (±100 Kev) beam of the Berkeley linear accelerator were used in this experiment. The range of the protons of approximately 30 Mev energy was found to be 18 ± 4 percent larger in glass than in dry C2 emulsion.

Discussion

The numbers used above to illustrate the calculation of momentum from target and plate radius are only approximate because the protons did not

follow a circular orbit in the radially decreasing cyclotron field. Protons leaving the target at a cyclotron radius r_1 at a small angle θ_1 from the backwards direction, enter the plate (r_2) at a different small angle θ_2 to perpendicular, and have a momentum given by the expression

$$\overline{Hr} = \frac{1}{r_1 \cos \theta_1 + r_2 \cos \theta_2} \int_{r_1}^{r_2} Hrdr \quad .$$

Stepwise integrations is required to evaluate the integral. Figure 4 shows a group of values of θ_1 determined graphically by stepwise trajectory plotting starting with various values of θ_2. In the range-energy study only tracks entering the emulsion at $0^\circ \pm 5^\circ$ were measured, so that the difference between $\cos \theta_1$ and $\cos \theta_2$ was negligible, and the expression for $\overline{Hr}$ could be written

$$\overline{Hr} = \frac{\sec \theta_2}{r_1 + r_2} \int_{r_1}^{r_2} Hrdr \quad .$$

The value of the integral was obtained by numerical integration using absolute field calibration at 4 points by nuclear induction to 4 gauss accuracy in the 14,000 gauss field, and flip coil measurements[6] of relative fields at 1 inch

---

[6] Private communication from D. Sewell

---

intervals along a radius. The relative values at the different points in the flip coil measurements were estimated to be accurate to 0.1 percent. The field is shown as a function of cyclotron radius in Figure 5. It was assumed, in the integration, that the cyclotron field had azimuthal symmetry.

The chief limit to the accuracy of the results was due to scatter in range of supposedly monoenergetic protons. The mean square scatter of 1 percent to 3 percent in ranges was partly caused by inaccuracies in measurement of entrance angle, partly by "skipping" of protons along the emulsion so that the

actual beginning in the emulsion was missed, and sometimes caused by surface fog which prevented exact determination of the beginning of a track. Approximately half, however, could be accounted for theoretically by straggling. Probable error of the mean due to scatter in range was less than 0.7 percent for each energy value. However, it seems unwise to claim better than ±2 percent accuracy for the range-energy relation because of incomplete control of emulsion moisture, and ignorance of possible variations in emulsion composition.

The point at 16.4 Mev which lies 2% above the curve may be caused by "slipping" or perhaps an error in the radius to which the target was set. All other points lie within 1/2% of the curve defined by the relation (1). The relation is probably accurate to better than 2% for any batch of dry B1, B2, C2, C3 or C5 plates.

The authors wish to express their appreciation of the stimulating interest of Prof. E. O. Lawrence, and of the cooperation of the cyclotron crew.

The work described in this paper was done under the auspices of the Atomic Energy Commission.

Information Division
0-12-49/lw

Figure 1

Schematic of arrangement for range-energy experiment.


P BEAM
PROTONS EJECTED FROM TARGET IN BACKWARD DIRECTION
CENTER OF CYCLOTRON
TARGET
PLATE
SHIELDING
30"
40"
50"
60"
70"
80"

13775-1

Figure 2

Range of protons vs. energy.


E (MEV) R (μ)
7.8 389 ± 2.7
16.4 1358 8.1
17.6 1465 4.5
17.6 1497 6.0
22.3 2244 8.9
25.6 2849 7.1
28.2 3369 10.0
33.5 4597 13.8
33.5 4762 24.0
33.5 4996 10.0
39.5 6123 18.0
X DRY EMULSION
o ~ 80 % R.H.
Δ ~ 90 % R.H.
+ DATA OF LATTES
FULL LINE IS DATA OF L, F, & C PLOTTED BELOW 13 MEV.
EXTRAPOLATION BY W. ARON PLOTTED ABOVE 13 MEV
PROTON RANGE — MICRONS (FOR ILFORD C-2 EMULSION)
ENERGY MEV
0
1000
2000
3000
4000
10
20
30
4000
4200
4400
4600
4800
32
34
36
38
40
9/8/49
14066-1

Figure 3

Power law approximation for range-energy relation.


PROTON RANGE - MICRONS (FOR ILFORD C-2 EMULSION)
ENERGY MEV
x DRY EMULSION
+ DATA OF LATTES
BEST FIT STRAIGHT LINE FROM
17 - 39 MEV $E_{(MEV)} = .251\ R^{.581}_{(\mu)}$
KEUFFEL & ESSER CO., N. Y. NO. 359-110
Logarithmic, 2 X 2 Cycles.
MADE IN U.S.A.

9/8/49

Figure 4

Emission angle vs. entrance angle.

ENTRANCE ANGLE OF PROTONS IN PLATES



R_2	θ_2	R_1	θ_1
80"	0	27.3	-1° 45'
80"	-3°	27.4	+1° 15'
80"	+3°	27.6	-5° 0'
80"	0	33.0	-2° 35'
80"	-3°	33.1	+1° 25'
80"	+3°	33.2	-4° 45'
80"	0	37.7	-1° 45'
80"	-3°	37.7	+1° 25'
80"	+3°	37.8	-4° 40'

Figure 5

Cyclotron magnetic field vs. radius.


x BISHOP & BRADNER (PROTON MOMENT)
o SEWELL (FLIP COIL)
MAGNETIC FIELD OF CYCLOTRON
% OF 15000 GAUSS
96
95
94
93
58
62
66
70
74
78
82
RADIUS FROM CYCLOTRON CENTER — INCHES