

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

COPY 2

UNCLASSIFIED

UCRL-528

Unclassified Distribution

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

MASS SPECTROGRAPHIC IDENTIFICATION OF
RADIOACTIVE LANTHANUM ISOTOPES

R. A. Naumann, F. L. Reynolds, and I. Perlman

November 28, 1949

Berkeley, California

<u>INSTALLATION</u>	<u>No. of Copies</u>
Argonne National Laboratory	8
Armed Forces Special Weapons Project	1
Atomic Energy Commission, Washington	2
Battelle Memorial Institute	1
Brookhaven National Laboratory	8
Bureau of Medicine and Surgery	1
Bureau of Ships	1
Carbide & Carbon Chemicals Corp. (K-25)	4
Carbide & Carbon Chemicals Corp. (Y-12)	4
Chicago Operations Office	1
Cleveland Area Office	1
Columbia University (Dunning)	1
Columbia University (Failla)	1
Dow Chemical Company	1
General Electric Company, Richland	6
Hanford Operations Office	1
Idaho Operations Office	1
Iowa State College	2
Kansas City	1
Kellex Corporation	2
Knolls Atomic Power Laboratory	4
Los Alamos	3
Mallinckrodt Chemical Works	1
Massachusetts Institute of Technology (Gaudin)	1
Massachusetts Institute of Technology (Kaufmann)	1
Mound Laboratory	3
National Advisory Committee for Aeronautics	2
National Bureau of Standards	2
Naval Radiological Defense Laboratory	2
NEPA Project	2
New Brunswick Laboratory	1
New York Operations Office	5
North American Aviation, Inc.	1
Oak Ridge National Laboratory	8
Patent Advisor, Washington	1
Rand Corporation	1
Sandia Base	1
Sylvania Electric Products, Inc.	1
Technical Information Branch, ORE	15
U. S. Public Health Service	1
UCLA Medical Research Laboratory (Warren)	1
University of California Radiation Laboratory	5
University of Rochester	2
University of Washington	1
Western Reserve University (Friedell)	2
Westinghouse Electric Corp.	4
Total	119

Information Division
 Radiation Laboratory
 Univ. of California
 Berkeley, California

Mass Spectrographic Identification of
Radioactive Lanthanum Isotopes

R. A. Naumann,* F. L. Reynolds, and I. Perlman
Radiation Laboratory and Department of Chemistry
University of California, Berkeley, California

Two neutron deficient isotopes of lanthanum, one new and the other previously tentatively assigned, have been given mass assignments by the technique of dispersing the isotopes on a photographic plate with a mass spectrograph. These and other light isotopes of lanthanum were prepared and some of their radioactive characteristics determined by absorption methods and with a low-resolution beta-ray spectrometer.

In some experiments, pure CsNO_3 was irradiated with 30-Mev alpha-particles and this was followed by the isolation of the lanthanum fraction. The mass spectrograph showed a single radioactivity at mass number 135; and by following the decay of this line on the photographic plate through a slit arrangement, a 19-hour half-life was obtained. This confirms the mass assignment previously made by other methods.¹ Since this irradiation should have produced La^{136} and La^{134} as well as La^{135} , it was inferred that La^{136} and La^{134} have half-lives less than about 20 minutes since it required several hours to make the exposure on the mass spectrograph. The isotope La^{136} had previously been reported as a 2-hour positron emitter,¹ but the observed activity has since been shown to be F^{18} which partially went through the particular chemical separation employed. A search was made for La^{136} , and it proved to be the 10-minute period reported by Maurer² from

*Now at Princeton University, Princeton, New Jersey.

¹J. B. Chubbuck and I. Perlman, *Phys. Rev.* 74, 982 (1948).

²W. Maurer, *Z. für Naturforschung* 2a, 586 (1947).

the irradiation of barium with deuterons. The assignment was made to La^{136} since the ratio of the 10-minute period to the 19-hour period increased with decreasing alpha-particle energy on cesium in the range below 30 Mev, as would be expected for the (α, n) reaction in relation to the $(\alpha, 2n)$ reaction. Furthermore, the preparation of this activity in good yield by Maurer² using 5.8-Mev deuterons on barium would now point to La^{136} in view of the present knowledge of La^{138} , La^{137} , and La^{135} , unless one assumes the 10-minute activity to be a metastable state of one of these isotopes. Very recently Robertson, Carss, and Pool³, using separated stable barium isotopes as targets, have definitely proved this period to be associated with La^{136} . The characteristics of La^{136} which we determined are a half-life of 9.5 minutes, positron of 2.1 Mev determined with the spectrometer, and x-rays in high abundance from which it is estimated that about 1/3 of the decay is by positron emission and 2/3 by electron-capture.

The irradiation of cesium with 50-Mev alpha-particles resulted in a new lanthanum activity of 4.0-hours half-life. This activity has positrons of about 1.2 Mev, a conversion electron line of 0.26 Mev determined with the spectrometer, a harder gamma-ray of 0.8 Mev measured by lead absorbers, and K x-rays in high yield. This activity apparently decays predominantly by electron-capture since the positrons were only a few percent as abundant as the x-rays.

The 4-hour period was assigned to La^{133} by the mass spectrograph method. Figure 1 shows the original photographic plate (A) and two transfer

³B. E. Robertson, W. L. Carss, and M. L. Pool, Bull. Am. Phys. Soc. 24, No. 7, 14 (1949).

Figure 1: Mass spectrographic analysis of radioactive lanthanum isotopes by transfer technique. (Exposures for transfer plates T_1 and T_2 given in the text.)




FIG. 1

plates (T_1 and T_2) made by exposing successively blank photographic plates with the receiver plate. The upper picture, A, shows the receiver plate with the line for stable lanthanum at mass number 139 (actually measured as the singly ionized oxide), and two other lines at 135 and 133. The middle plate, T_1 , is the first transfer plate exposed for 24 hours after analysis showing that the lines at 133 and 135 are due to radioactivities. The bottom plate, T_2 , made by a 72-hour exposure following the first one, shows only the 19-hour La^{135} since the 4-hour La^{133} had decayed.

Our thanks are due to Mr. B. Rossi, Mr. J. Vale, and the members of the groups operating the 60-inch and 184-inch cyclotrons, and to Mr. J. G. Conway for showing the CsNO_3 used to be spectrographically pure. The work described in this paper was performed under the auspices of the Atomic Energy Commission.