UCRL-8517

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Pub. in Phys. Rev.

UCRL-8517

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

ALPHA-DECAY BARRIER PENETRABILITIES WITH AN EXPONENTIAL NUCLEAR POTENTIAL
II. ODD-MASS NUCLEI

John O. Rasmussen

March 1959

Printed for the U. S. Atomic Energy Commission

ALPHA-DECAY BARRIER PENETRABILITIES WITH AN EXPONENTIAL NUCLEAR POTENTIAL*

II. ODD-MASS NUCLEI

John O. Rasmussen

Lawrence Radiation Laboratory and Department of Chemistry
University of California, Berkeley, California

March 1959

ABSTRACT

Calculation of barrier penetrabilities, reduced widths, and hindrance factors for odd-mass alpha-particle emitters is made by using the diffuse exponential nuclear potential derived from optical-model analysis of alpha elastic scattering data. The calculations are made on the same basis as for even-even alpha emitters, reported in a previous publication.

*Work performed under the auspices of the U. S. Atomic Energy Commission

ALPHA-DECAY BARRIER PENETRABILITIES WITH AN EXPONENTIAL NUCLEAR POTENTIAL
II. ODD-MASS NUCLEI

John O. Rasmussen

Lawrence Radiation Laboratory and Department of Chemistry
University of California, Berkeley, California

March 1959

In a previous publication,[1] hereafter referred to as I, the calculation of alpha-decay barrier-penetration factors by using an exponential nuclear potential was discussed in detail and numerical results for all measured alpha transitions of even-even nuclei were given. The nuclear potential used was the real part of a potential by Igo to fit alpha elastic scattering data.[2]

$$V(r) = -1100 \exp\left\{\frac{1.17\ A^{1/3} - r}{0.574}\right\} \quad \text{Mev.}$$

Details of the IBM-650 computer calculation of the barrier-penetration factor P based on experimental alpha-decay energies are given in I. After calculation of P the experimental alpha-decay rate information is used to calculate a "reduced alpha-emission width" δ^2 from the expression

$$\lambda = \delta^2\ P/h,$$

where λ is the decay constant and h is Planck's constant.

The computer program permits inclusion of the centrifugal potential for any desired value of angular momentum. In contrast to the even-even nuclei of spin zero, where only one value of ℓ is allowed for a given decay group, one usually expects a mixture of two or more ℓ values in the alpha groups of odd nuclei with $I > 1/2$. In only a few cases is this admixture experimentally known, so that we have chosen, in the spirit of the "hindrance factor" concept, to calculate all barrier penetrabilities here with $\ell = 0$.

The experimental data used and the calculated P and δ^2 values are given in Table I. The data are taken mainly from the 1958 Table of Isotopes.[3] Table I also lists in the last column the hindrance factor, F, and the basis of calculation of F needs further discussion.

The hindrance factor is the factor by which a given alpha group decays more slowly than would be calculated by simple barrier-penetration theory, normalized to the ground-state transition rates of neighboring even-even nuclei. The hindrance factor is to alpha decay what the $f_o t$ value is to beta decay. There is an arbitrariness in the process of determining the normal unhindered rate from even-even neighbors. The previous extensive tabulations of hindrance factors (or "departure factors" as they were earlier called) have usually normalized to some average behavior of even-even nuclei over a region.[4,5] Asaro has made calculations of F for odd-A nuclei based solely on the average rate behavior of the two nearest even-even neighbors. It is this latter basis that was chosen for the F calculations of Table I, i.e.

$$F = \frac{\delta_1^2 + \delta_2^2}{2\,\delta_{odd}^2},$$

where δ_1^2 and δ_2^2 are the reduced widths for ground-state transitions of the nearest neighboring even-even nuclei (given in Table I of Ref. 1). In those cases for which the data are unknown for one or both of the nearest even-even neighbors, the basis for the unhindered rate was taken from the one (or more) next nearest neighbor(s).

The three-significant-figure accuracy to which calculated results are given in Table I is seldom justified on an absolute basis because of experimental uncertainties in energies or half lives. Such accuracy is often necessary to afford significant comparison of <u>relative</u> hindrance factors of different alpha groups of the same alpha emitter, and such relative comparisons to 1% accuracy are often justified by accuracy of the data, since only the relative intensities and energy differences of compared states are of major importance to the ratio of hindrance factors.

The hindrance factors given in this paper will usually differ only slightly from those calculated by earlier methods, and the difference will seldom be significant from an absolute standpoint. The three-figure accuracy of the computer calculation should, however, make possible more careful comparisons of relative hindrance factors in future theoretical studies.

For some purposes it may be desirable to know the effect of a nonzero angular momentum on the calculated values of Table I, and we therefore offer the following approximate formula:

$$P_L/P_0 = \exp\,[-2.027\ L(L+1)Z^{-1/2}\ A^{-1/6}].$$

This formula uses the calculated[1] centrifugal barrier factor P_ℓ/P_4 of 5.917 for $\ell = 4$ on a ${}_{88}Ra^{224}$ group of energy 5.481 Mev to normalize the approximate relationship given in Eq. 22.10 of Ref. 3. The Z and A in the formula refer to atomic number and mass number of the daughter nucleus, respectively.

ACKNOWLEDGMENTS

I wish to acknowledge the valuable aid of Dr. Richard Chasman, who applied a computer calculation of screening energies to facilitate preparation of the input-data cards for the main calculations. Thanks are due Miss Claudette Everson and Miss Edna Williams for the actual running of most of the computer calculations. I am also grateful to Mrs. Mab Tocher for aid in preparing and checking the table of results in this paper.

REFERENCES

1. J. O. Rasmussen, "Alpha Decay Barrier Penetrabilities with an Exponential Nuclear Potential. I. Even-Even Nuclei" Phys. Rev. (in press) UCRL-8452.
2. G. Igo, Phys. Rev. Letters 1, 72 (1958).
3. Strominger, Hollander, and Seaborg, Revs. Modern Phys. 30, 585 (1958).
4. Perlman, Ghiorso, and Seaborg, Phys. Rev. 77, 26 (1950). I. Perlman and F. Asaro, Ann. Rev. of Nucl. Sci. 4, 157 (1954). Bohr, Fröman, and Mottelson, Dan. Mat. Fys. Medd. 29, No. 10 (1955). C. J. Gallagher, Jr. and J. O. Rasmussen, J. Inorg. Nucl. Chem. 3, 333 (1957). P. O. Fröman, Mat. Fys. Skr. Dan. Vid. Selsk. 1, No. 3 (1957). I. Perlman and J. O. Rasmussen, "Alpha Radioactivity" Handbuch der Physik, Vol. 42 (1957).

Table I. Alpha Groups of Odd-Mass Nuclei

Alpha Decay Data[a] Assumed for Computation						Calculated Results		
Atomic no.	Mass no.	Neutron number	$E_\alpha + E_{sc}$ (Mev)[b]	Partial α half life (sec)	α group intensity (%)	Barrier penetration factor P	Reduced width δ^2(Mev)	Hindrance factor F
62	147	85	2.201	4.12 (18)	100	1.378(-39)	5.047(-1)	3.01(-2)
63	147	84	2.921	2.075(11)	100	1.394(-31)	9.910(-2)	5.07(-1)
64	149	85	3.021	1.15 (11)	100	2.612(-31)	9.541(-2)	8.93(-1)
65	149	84	3.972	1.48 (5)	100	1.692(-24)	1.145(-2)	7.44 (0)
	151	86	3.422	2.4 (10)	100	1.947(-28)	6.133(-4)	1.39 (2)
83	199	116	5.501	1.578 (7)	100	6.001(-25)	3.027(-4)	8.26 (1)
	201	118	5.181	1.260 (8)	100	1.288(-26)	1.767(-3)	1.41 (1)
	203	120	4.881	6.310(11)	100	2.465(-28)	1.843(-5)	1.63 (3)
	211	128	6.651	1.300 (2)	82.6	9.879(-20)	1.843(-4)	3.872(2)
			6.304		17.4	4.296(-21)	8.930(-4)	7.993(1)
	213	130	5.891	1.382 (5)	100	7.601(-23)	2.728(-4)	4.061(2)
84	205	121	5.232	7.260 (6)	100	8.493(-27)	4.648(-2)	4.99(-1)
	207	123	5.132	1.893 (8)	100	2.496(-27)	6.066(-3)	3.11 (0)
	209	125	4.909	3.156 (9)	99.4	1.279(-28)	7.057(-3)	1.22 (0)
			4.652		.6	2.896(-30)	1.881(-3)	4.57 (0)
	211	127	7.472	5.200(-1)	99.0	2.540(-17)	2.149(-4)	1.82 (2)
			6.912		.53	3.302(-19)	8.848(-5)	4.42 (2)
			6.5		.50	2.125(-20)	1.297(-3)	3.01 (1)
	211	127	8.732	2.500 (1)	7.0	8.220(-14)	9.763(-11)	4.01 (8)
			7.882		2.5	4.437(-16)	6.460(-9)	6.05 (6)
			7.171		90.5	2.654(-18)	3.910(-5)	1.00 (3)
	213	129	8.382	4.200(-6)	100	1.143(-14)	5.970(-2)	1.526(0)
	215	131	7.412	1.830(-3)	100	1.892(-17)	8.276(-2)	1.327(0)

Table I (cont'd.)

Alpha Decay Data[a] Assumed for Computation						Calculated Results		
Atomic no.	Mass no.	Neutron number	$E_\alpha+E_{sc}$ (Mev)[b]	Partial α half life (sec)	α group intensity (%)	Barrier penetration factor P	Reduced width δ^2(Mev)	Hindrance factor F
	217	133	6.572	1.000 (1)	100	2.215(-20)	1.294(-2)	8.840(0)
85	207	122	5.782	7.200 (4)	100	2.006(-24)	1.984(-2)	6.57 (0)
	209	124	5.674	4.320 (5)	100	6.411(-25)	1.035(-2)	1.372(0)
	211	126	5.894	6.590 (4)	100	7.926(-24)	5.487(-3)	1.70 (0)
	215	130	8.032	1.00 (-4)	100	5.773(-16)	4.965(-2)	2.332(0)
	217	132	7.082	1.800(-2)	100	6.231(-19)	2.556(-1)	8.437(-1)
	219	134	6.302	5.400 (1)	100	6.863(-22)	7.734(-2)	1.965(0)
86	207	121	6.173	1.620 (4)	100	4.180(-23)	4.232(-3)	2.26 (0)
	209	123	6.070	1.080 (4)	100	1.565(-23)	1.696(-2)	8.119(-1)
	211	125	5.880	2.230 (5)	33.5	2.220(-24)	1.940(-3)	7.702(0)
			5.812		64.5	1.047(-24)	7.915(-3)	1.887(0)
			5.646		2.0	1.577(-25)	1.630(-3)	9.168(0)
	217	131	7.773	1.000(-3)	100	4.392(-17)	6.526(-2)	4.937(0)
	219	133	6.840	3.92 (0)	69	3.337(-20)	1.512(-2)	1.675(1)
			6.575		15	3.162(-21)	3.469(-2)	7.300(1)
			6.450		12	9.866(-22)	8.893(-2)	2.847(0)
			6.230		4	1.162(-22)	2.517(-1)	1.006(0)
	221	135	6.033	7.56 (3)	100	1.660(-23)	2.284(-2)	7.553(0)
87	219	132	7.333	2.00(-2)	100	7.244(-19)	1.978(-1)	1.629(0)
	221		6.363	2.88 (2)	84	1.612(-22)	5.186(-2)	3.108(0)
			6.153		16	1.953(-23)	8.153(-2)	1.977(0)
	223		5.373	2.64 (7)	100	2.727(-27)	3.982(-2)	3.852(0)

Table I (cont'd.)

Alpha Decay Data[a] Assumed for Computation						Calculated Results		
Atomic no.	Mass no.	Neutron number	$E_\alpha + E_{sc}$ (Mev)[b]	Partial α half life (sec)	α group intensity (%)	Barrier penetration factor P	Reduced width δ^2(Mev)	Hindrance factor F
88	221	133	6.744	3.00 (1)	100	2.065(-21)	4.627(-2)	2.984(0)
	223	135	5.894	1.17 (6)	1	4.322(-25)	5.668(-5)	7.505(3)
			5.767		9	1.043(-25)	2.114(-3)	6.717(1)
			5.738		53	7.276(-26)	1.784(-2)	7.958(0)
			5.626		24	1.931(-26)	3.045(-2)	4.664(0)
			5.559		9	8.559(-27)	2.576(-2)	5.513(0)
			5.521		2	5.359(-27)	9.142(-3)	1.513(1)
			5.452		3	2.261(-27)	3.251(-2)	4.368(0)
89	223	134	6.675	1.32 (2)	100	4.244(-22)	5.116(-2)	2.699(0)
	225	136	5.853	8.64 (5)	54	9.504(-26)	1.885(-2)	7.745(0)
			5.817		28	6.282(-26)	1.478(-2)	9.878(0)
			5.756		9.5	3.086(-26)	1.021(-2)	1.430(1)
			5.748		2.6	2.808(-26)	3.071(-3)	4.754(1)
			5.707		.8	1.728(-26)	1.535(-3)	9.54 (1)
			5.662		3.8	1.008(-26)	1.251(-2)	1.167(1)
			5.634		.6	7.182(-27)	2.771(-3)	5.269(1)
			5.605		.7	5.042(-27)	4.606(-3)	3.170(1)
			5.578		.07	3.618(-27)	6.418(4)	2.275(1)
	227	138	4.977	5.68(10)	100	1.171(-30)	4.310(-2)	3.202(1)
90	225	135	6.605	5.40 (2)	100	8.471(-23)	6.266(-2)	2.31 (1)
	227	137	6.071	1.62 (6)	23	3.848(-25)	1.057(-3)	1.277(2)
			6.042		2.8	2.780(-25)	1.770(-4)	7.627(1)

Table I (cont'd.)

Alpha Decay Data[a] Assumed for Computation						Calculated Results		
Atomic no.	Mass no.	Neutron number	$E_\alpha + E_{sc}$ (Mev)[b]	Partial α half life (sec)	α group intensity (%)	Barrier penetration factor P	Reduced width δ^2(Mev)	Hindrance factor F
			6.011		24	1.985(-25)	2.139(-3)	6.311(1)
			5.993		3.5	1.625(-25)	3.811(-4)	3.542(1)
			5.949		.9	9.912(-26)	1.606(-4)	8.406(1)
			5.900		3.0	5.679(-26)	9.347(-4)	1.444(2)
			5.840		1.0	2.842(-26)	6.225(-4)	2.169(2)
			5.828		.3	2.471(-26)	2.148(-4)	6.288(2)
			5.796		.3	1.699(-26)	3.125(-4)	4.321(2)
			5.790		21	1.583(-26)	2.348(-2)	5.750(0)
			5.747		5.0	9.507(-27)	9.304(-3)	1.451(1)
			5.743		8.7	9.063(-27)	1.698(-2)	7.95 (0)
			5.734		4.0	8.139(-27)	8.694(-3)	1.553(1)
			5.727		1.5	7.484(-27)	3.546(-3)	3.807(1)
			5.702		1.9	5.541(-27)	6.067(-3)	2.225(1)
	229	139	5.055	2.316(11)	10	1.128(-30)	1.097(-3)	1.144(2)
			4.975		20	3.467(-31)	7.138(-3)	1.760(1)
			4.885		70	8.881(-32)	9.754(-2)	1.288(0)
91	227	136	6.496	2.70 (3)	100	1.104(-23)	9.617(-2)	1.30 (0)
	229	138	5.726	5.05 (7)	100	2.507(-27)	2.264(-2)	5.47 (0)
	231	140	5.081	1.08(12)	10	5.095(-31)	5.209(-4)	2.29 (2)
			5.052		23	3.329(-31)	1.834(-3)	6.51 (1)
			5.036		24	2.628(-31)	2.424(-3)	4.93 (1)
			5.006		2.3	1.681(-31)	3.631(-4)	3.29 (2)

Table I (cont'd.)

Alpha Decay Data[a] Assumed for Computation						Calculated Results		
Atomic no.	Mass no.	Neutron number	$E_\alpha + E_{sc}$ (Mev)[b]	Partial α half life (sec)	α group intensity (%)	Barrier penetration factor P	Reduced width δ^2(Mev)	Hindrance factor F
			4.973		22	1.024(-31)	5.703(-3)	2.09 (1)
			4.956		2.8	7.914(-32)	9.389(-4)	1.27 (2)
			4.874		1.4	2.241(-32)	1.658(-3)	7.20 (1)
			4.757		11	3.496(-33)	8.350(-2)	1.43 (0)
			4.731		1.4	2.291(-33)	1.621(-2)	7.37(0)
			4.702		2.1	1.401(-33)	3.978(-2)	3.002(0)
92	227	135	6.836	7.8 (1)	100	1.042(-22)	3.527(-1)	2.70(-1)
	229	137	6.456	1.8 (4)	100	2.784(-24)	5.720(-2)	1.91 (0)
	231	139	5.486	6.63 (9)	100	4.071(-29)	1.062(-2)	1.11 (1)
	233	141	4.852	5.11(12)	83.5	4.645(-33)	1.008(-1)	1.11 (0)
			4.809		14.9	2.335(-33)	3.579(-2)	3.14 (0)
			4.752		1.6	9.248(-34)	9.704(-3)	1.16 (1)
			4.525		.03	1.937(-35)	8.687(-3)	1.29 (1)
	235	143	4.595	2.24(16)	6.7	7.062(-35)	1.214(-4)	8.871(2)
			4.556		2.7	3.584(-35)	9.638(-5)	1.116(3)
			4.502		0.9	1.381(-35)	8.338(-5)	1.292(3)
			4.407		29	2.469(-36)	1.503(-2)	7.162(0)
			4.394		43	1.942(-36)	2.833(-2)	3.800(0)
			4.375		12	1.365(-36)	1.125(-2)	9.573(0)
			4.212		5.8	5.991(-38)	1.239(-1)	8.695(-1)

Table I (cont'd.)

Alpha Decay Data[a] Assumed for Computation						Calculated Results		
Atomic no.	Mass no.	Neutron number	$E_\alpha + E_{sc}$ (Mev)[b]	Partial α half life (sec)	α group intensity (%)	Barrier penetration factor P	Reduced width δ^2(Mev)	Hindrance factor F
93	231	138	6.316	1.8 (5)	100	2.411(-25)	6.603(-2)	1.86 (0)
	233	140	5.566	2.209(8)	100	3.833(-29)	3.385(-1)	2.76(-1)
	235	142	5.097	1.009(12)	100	5.992(-32)	4.740(-2)	2.01 (0)
	237	144	4.909	6.94 (13)	3.1	3.612(-33)	3.545(-4)	2.555(2)
			4.853		3.5	1.481(-33)	9.761(-4)	9.278(1)
			4.824		53	9.274(-34)	2.360(-2)	3.837(0)
			4.803		29	6.698(-34)	1.788(-2)	5.064(0)
			4.750		1.7	2.754(-34)	2.549(-3)	3.552(1)
			4.710		3.3	1.435(-34)	9.497(-3)	9.535(0)
			4.681		6	8.643(-35)	2.867(-2)	3.159(0)
			4.626		.5	3.367(-35)	6.132(-3)	1.477(1)
			4.557		.02	1.007(-35)	8.204(-4)	1.104(2)
94	233	139	6.337	1.20 (6)	100	1.101(-25)	2.169(-2)	5.16 (0)
	235	141	5.887	7.89 (7)	100	7.130(-28)	5.095(-2)	1.97 (0)
	237	143	5.687	1.19(11)	21	6.446(-29)	7.847(-5)	1.059(3)
			5.397		79	1.388(-30)	1.371(-2)	6.061(0)
	239	145	5.184	7.67(11)	72.5	7.223(-32)	3.751(-2)	2.476(0)
			5.171		16.8	5.967(-32)	1.052(-2)	8.827(0)
			5.133		10.7	3.398(-32)	1.177(-2)	7.890(0)
			5.101		.037	2.105(-32)	6.569(-5)	1.414(3)
			5.036		.013	7.838(-33)	6.198(-5)	1.498(3)
			4.954		.005	2.192(-33)	8.525(-5)	1.089(3)

Table I (cont'd.)

Alpha Decay Data[a] Assumed for Computation						Calculated Results		
Atomic no.	Mass no.	Neutron number	$E_\alpha+E_{sc}$ (Mev)[b]	Partial α half life (sec)	α group intensity (%)	Barrier penetration factor P	Reduced width δ^2(Mev)	Hindrance factor F
	241	147	4.930	9.47 (12)	75	1.621(-33)	1.400(-1)	7.14(-1)
			4.885		25	7.901(-34)	9.576(-2)	1.04 (0)
95	237	142	6.048	9.47 (7)	100	1.670(-27)	1.813(-2)	4.84(0)
	239	144	5.788	1.041 (9)	100	7.766(-29)	3.545(-2)	2.345(0)
	241	146	5.573	1.455(10)	.42	5.240(-30)	1.579(-4)	5.598(2)
			5.541		.24	3.421(-30)	1.387(-4)	6.395(2)
			5.514		84.3	2.380(-30)	6.978(-2)	1.267(0)
			5.471		13.6	1.328(-30)	2.017(-2)	4.381(0)
			5.417		1.4	6.317(-31)	4.365(-3)	2.025(1)
			5.352		.015	2.543(-31)	1.162(-4)	7.608(2)
			5.308		.004	1.361(-31)	5.791(-5)	1.526(3)
	243	148	5.377	2.493(11)	.17	3.891(-31)	5.023(-5)	1.57 (3)
			5.346		.16	2.514(-31)	7.316(-5)	1.08 (3)
			5.304		86.9	1.383(-31)	7.226(-2)	1.09 (0)
			5.262		11.5	7.547(-32)	1.752(-2)	4.51 (0)
			5.207		1.3	3.377(-32)	4.426(-3)	1.78 (1)
96	241	145	5.988	3.156 (8)	100	3.118(-28)	2.913(-2)	2.701(1)
	243	147	6.099	1.105 (9)	1	1.241(-27)	2.090(-5)	3.220(3)
			6.092		5	1.144(-27)	1.134(-4)	5.935(2)
			6.043		.9	6.430(-28)	3.630(-5)	1.854(3)
			6.025		6	5.194(-28)	2.996(-4)	2.246(2)
			5.938		.1	1.826(-28)	1.420(-5)	4.739(3)

Table I (cont'd.)

Alpha Decay Data[a] Assumed for Computation						Calculated Results		
Atomic no.	Mass no.	Neutron number	$E_\alpha + E_{sc}$ (Mev)[b]	Partial α half life (sec)	α group intensity (%)	Barrier penetration factor P	Reduced width δ^2(Mev)	Hindrance factor F
			5.910		.5	1.300(-28)	9.995(-5)	6.733(2)
			5.818		73	4.149(-29)	4.564(-2)	1.475(0)
			5.774		11.5	2.381(-29)	1.253(-2)	5.371(0)
			5.718		1.6	1.163(-29)	3.569(-3)	1.886(1)
			5.714		.2	1.104(-29)	4.700(-4)	1.432(2)
			5.672		.15	6.401(-30)	6.078(-4)	1.107(2)
	245	149	5.488	4.418(11)	15	5.893(-31)	1.651(-3)	3.923(1)
			5.398		77	1.684(-31)	2.966(-2)	2.189(0)
			5.348		8	8.281(-32)	6.267(-3)	1.036(1)
97	243	146	6.759	1.578 (7)	30	5.337(-25)	1.021(-4)	6.824(2)
			6.589		53	9.362(-26)	1.028(-3)	6.775(1)
			6.239		17	2.067(-27)	1.498(-2)	4.652(0)
	245	148	6.409	3.58 (8)	33	1.476(-26)	1.790(-4)	3.425(2)
			6.209		41	1.576(-27)	2.083(-3)	2.943(1)
			5.929		26	5.653(-29)	3.682(-2)	1.665(0)
	247	150	5.709	3.156(11)	37	3.729(-30)	9.010(-4)	4.96 (1)
			5.549		58	4.398(-31)	1.198(-2)	3.73 (0)
			5.339		5	2.289(-32)	1.984(-2)	2.25 (0)
	249	152	5.459	1.23 (12)	94	1.362(-31)	1.608(-2)	3.694(0)
			5.069		6	4.183(-34)	3.342(-1)	1.777(-1)
98	245	147	7.150	7.92 (3)	100	9.046(-24)	4.001(-2)	1.44 (0)
	249	151	6.234	1.14 (10)	1.9	8.100(-28)	5.898(-6)	8.833(3)
			6.179		1.1	4.279(-28)	6.463(-6)	8.061(3)
			6.112		.4	1.943(-28)	5.175(-6)	1.007(4)

Table I (cont'd.)

Alpha Decay Data[a] Assumed for Computation						Calculated Results		
Atomic no.	Mass no.	Neutron number	$E_\alpha + E_{sc}$ (Mev)[b]	Partial α half life (sec)	α group intensity (%)	Barrier penetration factor P	Reduced width δ^2(Mev)	Hindrance factor F
			6.030		.08	7.260(-29)	2.770(-6)	1.881(4)
			5.981		3.3	3.991(-29)	2.079(-4)	2.506(2)
			5.938		1.2	2.345(-29)	1.286(-4)	4.051(2)
			5.882		3.0	1.163(-29)	6.485(-4)	8.033(1)
			5.846		84	7.369(-30)	2.866(-2)	1.818(0)
			5.818		.5	5.152(-30)	2.440(-4)	2.135(2)
			5.789		4.4	3.546(-30)	3.119(-3)	1.670(1)
			5.727		.4	1.580(-30)	6.366(-4)	8.184(1)
99	249	150	6.800	5.44(6)	100	1.275(-25)	4.131(-3)	1.08 (1)
	251	152	6.520	2.52(7)	100	7.243(-27)	1.570(-2)	3.78 (0)
	253	154	6.673	1.73(6)	90.2	3.982(-26)	3.753(-2)	1.98(0)
			6.632		7.7	2.587(-26)	4.931(-3)	1.50 (1)
			6.592		.8	1.689(-26)	7.845(-4)	9.440(1)
			6.580		.9	1.486(-26)	1.004(-3)	7.376(1)
			6.537		.25	9.355(-27)	4.427(-4)	1.672(2)
			6.519		.1	7.697(-27)	2.152(-4)	3.441(2)
			6.469		.1	4.458(-27)	3.716(-4)	1.993(2)
			6.289		.04	5.895(-28)	1.124(-3)	6.589(1)
			6.249		.05	3.715(-28)	2.230(-3)	3.321(1)
			6.216		.02	2.529(-28)	1.310(-3)	5.653(1)
100	251	151	6.931	2.52(6)	100	1.834(-25)	6.200(-3)	8.31 (0)
	253	153	6.891	3.54(6)	100	1.322(-25)	6.123(-3)	8.251(0)
	255	155	7.071	7.74(4)	100	8.391(-25)	4.413(-2)	1.145(0)

a. It is to be most strongly emphasized that this table is not to be used as a reference for energy and half life data. In many cases only the crudest partial alpha half life or energy values are to be found in the Table of Isotopes. We give in Table I numbers used in the computer, simply in order to facilitate correcting the table for revised experimental measurements.

b. Alpha particle energy plus orbital electron screening correction.

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, express or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission to the extent that such employee or contractor prepares, handles or distributes, or provides access to, any information pursuant to his employment or contract with the Commission.