UCRL 9196

UNIVERSITY OF CALIFORNIA

Ernest O. Lawrence Radiation Laboratory

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UCRL-9196

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

EFFECT OF THE PION-PION RESONANCE ON K^--p SCATTERING

Fabio Ferrari, Graham Frye, and Modesto Pusterla

May 4, 1960

EFFECT OF THE PION-PION RESONANCE ON K^--p SCATTERING*

Fabio Ferrari, Graham Frye, and Modesto Pusterla

Lawrence Radiation Laboratory
University of California
Berkeley, California

May 4, 1960

Considerable theoretical interest has been shown in the interpretation of the low-energy K^--p scattering data. The major effort has been directed toward a convenient parameterization that takes into account the kinematic features of the many competing channels ($K^- + p \rightarrow \bar{K}^0 + n$, πY) and the "standard" Coulomb and mass-difference corrections.[1,2,3,4] In attempts to draw conclusions about the nature of the basic K-meson nucleon interaction, the assumption has been made that $k \cot \delta$ is essentially constant and equal to the reciprocal (complex) scattering length. We have examined the validity of this approximation on the basis of the Chew-Mandelstam program.[5] We found that the two-pion exchange, which determines the long-range tail of the K-N interaction, gives a substantial energy dependence to $k \cot \delta$.

A qualitative determination that remains to be made is the sign of the real part of each ($I = 0,1$) scattering length.[6] Two attempts to do this

* This work was supported by the U. S. Atomic Energy Commission, a grant from the National Academy of Science (F.F.), and the U. S. Air Force, and monitored by the Air Force Office of Scientific Research of the Air Research and Development Command.

have been based on properties of the Coulomb-nuclear interference. The angular distribution at 172 Mev/c (laboratory-system momentum) favors a constructive interference[7] and implies a positive sign. The other attempt is based on the apparent leveling off and decrease of the elastic-scattering cross section with decreasing momentum ($P_L < 150$ Mev/c). Jackson and Wyld have suggested that this behavior is due to destructive interference and consequently concluded that the sign is negative.[3] The energy dependence arising from the two-pion exchange provides an alternative interpretation of the leveling off. This can be seen qualitatively from the following argument: by virtue of the small pion mass (compared with the K meson mass) the two-pion exchange determines the longest-range part of the K^--p potential. This suggests that we regard the nuclear interaction as made up of the two-pion contribution plus another part of shorter range, representing the net effect of everything else. The peculiar energy dependence is interpreted as the destructive interference between the two parts. Knowledge of the sign of the two-pion contribution then leads to a determination of the sign of the scattering length. Recent advances in the theory of the pion-pion interaction[8] and of the electromagnetic structure of the nucleon[9] make it possible to calculate the sign and estimate the magnitude of the two-pion contribution.

From their theoretical study of the nucleon electromagnetic structure, Frazer and Fulco[9] have inferred a resonance in the I = 1, J = 1 state of pion-pion scattering. Quantitative conclusions are still uncertain,[10] but it seems likely that the two-pion contribution accounts for a large fraction of the isotopic vector charge and magnetic moment. On the basis of their

results, we suggest that the charge structure of the K meson also receives a sizable contribution from the two-pion state. The consistency of this hypothesis and its consequences for πK scattering has been studied.[11] This hypothesis provides an estimate of the order of magnitude of the matrix element for emission of two pions by a K meson $(\pi\pi \mid K\bar{K})$. Using this estimate, we have calculated the contribution of the two-pion exchange to the $\bar{K}$-nucleon interaction. (The details of this calculation will be presented in a separate report.)

We denote the S-wave $\bar{K}$-nucleon elastic-scattering amplitude by g. It is an analytic function in the cut s-plane (s is the total energy square in the $\bar{K}$-N center-of-mass system). The physical branch cut starts at $s = s_0 = (M_N + m_K)^2$ and extends to $+\infty$. There are other "right-hand" cuts starting from the thresholds of the dynamically coupled channels, $(m_\pi + M_\Lambda)^2$ and $(m_\pi + M_\Sigma)^2$.[12] The unitarity condition, which establishes certain nonlinear relations between the amplitudes g_{ij} of various coupled processes, is expressed by

$$\mathrm{Im}\,(g^{-1})_{ij} = -\frac{k_i\,\Theta_i\,\delta_{ij}}{\sqrt{s}\,(E_i + M_i)}\,, \qquad (1)$$

$$\Theta_i = \begin{cases} 1 & \text{if} \quad W_i < \sqrt{s}\,, \\ 0 & \text{if} \quad W_i > \sqrt{s}\,, \end{cases}$$

where k_i, E_i, and W_i are the center-of-mass momentum, baryon energy, and threshold energy of the ith channel. The dynamical singularity arising from the two-pion exchange is a branch cut extending from the left up to the point $s = s_2 = \sqrt{M_N^2 - m_\pi^2} + \sqrt{m_K^2 - m_\pi^2}$. Relative to the

other dynamical singularities, this cut is very close to the physical region $s \geq s_0$, and should produce the strongest energy dependence in $k \cot \delta$.

We denote the elastic $\bar{K}N$ amplitude in the state of isotopic spin I by g_I, where $I = 0,1$. These two amplitudes are related by crossing symmetry to amplitudes $g^{(+)}$ and $g^{(-)}$ that depend on the isotopic spin $I' = 0, 1$ of the exchanged pion pair, namely

$$g^0 = g^{(+)} - 3g^{(-)} ,$$
$$g^1 = g^{(+)} + g^{(-)} . \tag{2}$$

The magnitude of the discontinuity across the two-pion cut is expressed for our purpose as

$$\int_{s_1}^{s_2} ds \, \mathrm{Im}\, g^{(-)}(s) = R_1 \approx 4 M_N^4 \text{ fermi} , \tag{3}$$

where s_1 is approximately equal to $80\, m_\pi^2$. Assuming that only the resonant $g^{(-)}$ amplitude is important, we have three specific predictions: (a) the isotopic spin ratio R_0/R_1 is -3, (b) the sign of R_1 is positive, and (c) the expected magnitude of R_1 is rather large.

The normalization of our amplitude g is given by its relation to the $\bar{K}$-N eleastic scattering phase shift,

$$g(s) = \sqrt{s}\,(E + M)/(k \cot \delta - ik) . \tag{4}$$

In this note we use only a very simple model in which the effect of the two-pion cut is represented by a δ function at the position $s = a = 93\, m_\pi^2$

in the amplitude g. The dynamical singularities in all other channels are represented by subtraction constants.

This model, combined with the many-channel unitarity condition, Eq. (1), leads to the following expression for $k \cot \delta$:

$$k \cot \delta_I = \left\{ \ell(s) + \frac{1}{f_I(s) + z_I} \right\} \sqrt{s}\,(E + M) , \qquad (5)$$

where

$$\ell(s) = \frac{s - s_0}{\pi} \; P \int_{s_0}^{\infty} \frac{ds' \, \mathrm{Im}(g^{-1})_{11}}{(s' - s)(s' - s_0)} - \frac{a - s_0}{\pi} \int_{s_0}^{\infty} \frac{ds' \, \mathrm{Im}(g^{-1})_{11}}{(s' - s_0)(s' - a)} ,$$

$$f_I(s) = \frac{1}{\pi} \left(\frac{1}{s - a} - \frac{1}{s_0 - a} \right) \frac{R_I}{1 + \beta R_I} ,$$

$$\beta = \frac{1}{\pi^2} \int_{s_0}^{\infty} \frac{ds' \, \mathrm{Im}(g^{-1})_{11}}{(s' - a)^2} ,$$

and z_I is a complex quantity which depends on the subtraction constants and on the slowly varying $\pi - Y$ center-of-mass momenta. For each isotopic spin channel, we approximate z_I by a complex constant.[1,12] The calculations were done by adjusting the parameters R_I and z_I to fit the at-rest branching ratio

$$\lambda^{-1} = \frac{\sigma_{abs}\,(I = 1)}{\sigma_{abs}\,(I = 0)} = 0.18 \pm 0.09 ,$$ [7]

the total cross section at 172 Mev/c ($k\,\sigma_{tot}/4\pi = 0.7$ fermi) and the

elastic and charge-exchange cross sections at 100 Mev/c and 172 Mev/c.

Two sets of solutions were found, corresponding to constructive or destructive Coulomb-nuclear interference in the angular distribution at 172 Mev/c (Table II). The elastic and charge-exchange cross sections resulting from the "constructive" solution are shown in Fig. 1. The energy dependence of $(k \cot \delta_I)^{-1}$ is shown in Fig. 2, where the (a+) solution of Dalitz and Twan is also reported for comparison. Note that only the values of R_I and the ratio R_0/R_1 given by the "constructive" solution agree with those calculated theoretically. From this consideration we reject the "destructive" solution. Our principal result is that $k \cot \delta_I$ has a rather substantial energy dependence due to the two-pion exchange. Figure 2 shows that $\mathrm{Re}\,(k \cot \delta_I)$ is positive for each isotopic spin state in the momentum range above 100 Mev/c. However, the real parts of the scattering lengths appear to have opposite signs, that of the $I = 0$ state being small and negative.

Finally, we would like to remark that, by crossing symmetry, the two-pion contribution to the K^+-proton amplitude can be determined from the parameters R_0 and R_1. In this case there is a destructive interference and flattening of the cross section for a repulsive K^+ proton short range interaction.

We are deeply indebted to Professors Geoffrey F. Chew and Robert Karplus for many illuminating discussions.

FIGURE CAPTIONS

Fig. 1. Cross sections for elastic and charge-exchange K^--proton scattering. The (a+) solution of Dalitz and Tuan (dotted lines) is included for comparison.

Fig. 2. Momentum dependence of the real and imaginary parts of $(k \cot \delta)^{-1}$ (a) for isotopic spin $I = 0$, (b) for isotopic spin $I = 1$. The dotted lines are the (a+) solution of Dalitz and Tuan.

TABLE I

Two sets of parameters that fit the available experimental data. The solutions are characterized by the constructive or destructive nature of the Coulomb-nuclear interference in the angular distribution at 172 Mev/c.

Solution	Constructive	Destructive
R_1	$1.0\ M_N^4$ fermi	$2.0\ M_N^4$ fermi
R_0/R_1	-5.8	+ 1.50
z_0	$(1.34 + i\ 0.74)M_N^2$ fermi	$(0.93 + i\ 1.46)M_N^2$ fermi
z_1	$(0.67 + i\ 0.12)M_N^2$ fermi	$(-0.22 + i\ 1.00)M_N^2$ fermi
λ^{-1}	0.13	0.29

REFERENCES

1. Richard H. Dalitz and San F. Tuan, Ann. Phys. 8, 100 (1959).

2. Paul T. Matthews and Abdus Salam, Phys. Rev. Letters 2, 226 (1959).

3. J. D. Jackson and H. W. Wyld, Phys. Rev. Letters 2, 355 (1959).

4. M. Ross and C. Shaw, Phys. Rev. 115, 1773 (1959).

5. Geoffrey F. Chew, Ann. Rev. Nuclear Sci. 9, 29 (1960); Stanley Mandelstam, Phys. Rev. 112, 1344 (1958).

6. We would like to reiterate that the attractive nature of the K-nucleon interaction is already determined by the large magnitude of at least one of the scattering lengths; Robert Karplus, L. Kerth, and T. Kycia, Phys. Rev. Letters 2, 510 (1959).

7. KIEV Report 1959; Data of Rosenfeld, Solmitz, Tripp, Ross reported by L. W. Alvarez at 9th Annual Conference on High Energy Nuclear Physics.

8. Geoffrey F. Chew and Stanley Mandelstam, Theory of the Low-Energy Pion-Pion Interaction, Lawrence Radiation Laboratory Report UCRL-8728.

9. William R. Frazer and Jose R. Fulco, Effect of a Pion-Pion Scattering Resonance on Nucleon Structure, submitted to Phys. Rev.

10. James Ball and David Wong, Lawrence Radiation Laboratory (private communication).

11. Graham Frye, Theory of the Low-Energy Pion K-Meson Interaction (to be submitted to Phys. Rev.).

12. S. W. MacDowell, Phys. Rev. 116, 774 (1959); Fabio Ferrari, Michael Nauenberg, and Modesto Pusterla, Low Energy Dependence of π - Y and $\bar{K}$-Nucleon S-Wave Amplitude, UCRL-8985, Nov. 1959 (unpublished). Michael Nauenberg, The Inelastic Scattering of Mesons and Hyperons (Ph.D. dissertation) Cornell University. Fabio Ferrari, Michael Nauenberg and

12. Cont.

Modesto Pusterla, Partial Waves Amplitudes for $\bar{K}$-Nuclear Scattering (to be published) Lawrence Radiation Laboratory Report.



MU-20295

Fig. 1. Cross sections for elastic and charge-exchange K^--proton scattering. The (a+) solution of Dalitz and Tuan (dotted lines) is included for comparison.



MU-20294

Fig. 2. Momentum dependence of the real and imaginary parts of $(k \cot \delta)^{-1}$ (a) for isotopic spin I = 0, (b) for isotopic spin I = 1. The dotted lines are the (a+) solution of Dalitz and Tuan.

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.