UCRL-9333 Rev

UNIVERSITY OF CALIFORNIA

Ernest O. Lawrence Radiation Laboratory

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

To be published in Physical Review

UCRL-9333 Rev.

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

EFFECTS OF THE PION-PION RESONANCE AND THE THREE-PION RESONANCE OR BOUND STATE ON NEUTRAL PION DECAY

How-sen Wong

September 14, 1960

UCRL-9333 (Rev.)

EFFECTS OF THE PION-PION RESONANCE AND THE THREE-PION RESONANCE OR BOUND STATE ON NEUTRAL PION DECAY

How-sen Wong

Lawrence Radiation Laboratory
University of California
Berkeley, California

September 14, 1960

ABSTRACT

We have applied the dispersion method to the problem of neutral pion decay. It is shown that a pion-pion P-wave resonance can produce a large effect in the decay matrix element. The contribution is related to the semifundamental constant which determines the rate of photopion production from pions. The contribution of a strong three-pion state is also considered.

EFFECTS OF THE PION-PION RESONANCE AND THE THREE-PION RESONANCE OR BOUND STATE ON NEUTRAL PION DECAY*†

How-sen Wong §

Lawrence Radiation Laboratory
University of California
Berkeley, California

September 14, 1960

Recently Frazer and Fulco[1] and Chew[2] have proposed that a two-pion P-wave resonance and a three-pion resonance or bound state may account for the isotopic vector and scalar components of nucleon electromagnetic structure respectively. The purpose of this note is to investigate the effects of such resonances on neutral-pion decay.

The dispersion analysis of neutral-pion decay was first considered by Goldberger and Treiman,[3] but they assumed nucleon-antinucleon pairs to be the most important intermediate states and neglected multipion states. Here we adopt a different approach and consider the contributions of the least massive states. This can be done if we extend a photon variable q^2 into the complex plane instead of the meson variable p^2 used by Goldberger and Treiman.

---

* Work done under the auspices of the U. S. Atomic Energy Commission.

† A preliminary account of this work was given at the meeting of the American Physical Society, November 27-28, 1959, Cleveland, Ohio, (How-sen Wong, Bull. Am. Phys. Soc. 4, 407 (1959)).

§ Now at Laboratory of Nuclear Studies, Cornell University, Ithaca, New York.

Following the standard method, one has[4] (see Fig. 1)

$$(q(\mu),\ k(\nu)\ |T|\ p(3)) = \frac{i(2\pi)^4\ \delta^4\ (p - q - k)\ F_\nu\ (-q^2;\ -k^2;\ -p^2)\epsilon'_\nu}{(8\ q_0\ k_0\ p_0)^{1/2}}, \tag{1}$$

where we have

$$F_\nu = (4\ p_0\ q_0)^{1/2}\ (\ q(\mu)\ |\ J_\nu\ (0)\ |\ p(3)\ ),$$

and p is the pion four-momentum. The indices μ and ν refer to the polarization of the photons of momenta q and k, respectively. The number "3" inside the matrix element represents a neutral pion in the initial state.

From general invariance arguments, the F function can be written in the form

$$F(-q^2;\ -k^2;\ -p^2) = \epsilon_{\alpha\beta\mu\nu}\ q_\alpha\ k_\beta\ \epsilon_\mu\ \epsilon'_\nu\ f(-q^2;\ -k^2;\ -p^2). \tag{2}$$

We assume that, with both p^2 and k^2 on the mass shell, the scalar function $f(-q^2)$ satisfies the following dispersion relation without subtraction:

$$f(-q^2) = \frac{1}{\pi} \int_4^\infty \frac{\mathrm{Im}\ f(t)\ dt}{t + q^2}. \tag{3}$$

The lifetime of π^0 is then given by

$$\tau = 64\ \pi\ (f\ (0)\ )^{-2}. \tag{4}$$

Using the unitary condition, we can express the absorptive part of F as

$$\mathrm{Im}\, F = \pi\, \epsilon_\mu\, \epsilon'_\nu \sqrt{2p_0} \sum_n \delta^4 (q - p_n) \langle 0 \mid J_\mu (0) \mid n \rangle \langle n \mid J_\nu (0) \mid p(3) \rangle.$$

Since our approach is to assume that the function is determined by near-by singularities, no intermediate state except the 2π and 3π states are considered here. Let us first investigate the effect from the two-pion P-wave resonance alone and leave the 3π contribution to be discussed later; then we can show

$$\mathrm{Im}\, F(-q^2) = \frac{e_r}{48\pi} \epsilon_{\alpha\beta\mu\nu}\, q_\alpha\, k_\beta\, \epsilon_\mu\, \epsilon'_\nu$$

$$\times \quad \frac{(-q^2 - 4)^{3/2}}{(-q^2)^{1/2}} F_\pi^{\dagger} (-q^2)\, M_1 (-q^2), \tag{5}$$

where $F_\pi^{\dagger}$ is the Hermitian conjugate of F_π, the pion form factor, and M_1 is the P-wave amplitude for photopion production from pions. It has been shown by the author that the M_1 function is linearly related to a real constant A and the F_π function.[5] Comparing Eqs. (2) and (5), we obtain

$$f_{2\pi}(-q^2) = \frac{e_r}{48\pi^2} \int_4^\infty \frac{(t - 4)^{3/2}}{\sqrt{t}\,(t + q^2)} F_\pi^{\dagger}(t)\, M_1(t)\, dt. \tag{6}$$

James Ball[6] has applied the Mandelstam representation to the $\gamma + N \to \pi + N$ problem, and finds that $|A|$ must be less than 1.8e in order to make his calculated cross section compatible with experimental data. If $|A| = 1.8e$ and Frazer-Fulco form factor is used for a resonance at $t_{2\pi} = 10$, we find $\tau = 2 \times 10^{-16}$ sec.[7] Since the experimental upper limit for τ is 4×10^{-16} sec.,[7] we see that the

contribution of a two pion resonance is capable of producing a large effect in the neutral pion decay. If the resonance is sharp it follows from (6) that $f_{2\pi}(-q^2) \sim (t_{2\pi} + q^2)^{-1}$.

We note that the f function in Eq. (6) depends on the virtual-photon mass variable. Thus $-x \equiv q^2 = (P_+ + P_-)^2$ represents the square of the total four momentum of the electron-positron pair in the process $\pi^0 \to \gamma + e^+ + e^-$. Since x is less than 1, we can write

$$f(x) = f_{2\pi}(0)(1 + \alpha x) \cong f_{2\pi}(0)\left(1 + \frac{x}{t_{2\pi}}\right) \tag{7}$$

for small x.[8] Thus, we see that α is always positive in this approximation.

The calculation so far is based on the assumption that only the 2π state contributes to the dispersion integral, but there is no good reason to expect the 3π contribution to be negligible. Since no one has succeeded in treating the matrix element $(\gamma\pi \mid 3\pi)$, we are not able to handle this part. However, we may observe that if the three-pion $I = 0$ and $J = 1$ state is resonant or bound at energy $\sqrt{t_{3\pi}}$, as suggested by Chew, then

$$f_{3\pi}(-q^2) = \frac{\text{constant}}{q^2 + t_{3\pi}} ,$$

so that

$$f(-q^2) = f_{2\pi}(-q^2) + f_{3\pi}(-q^2) = \frac{f_{2\pi}(0)}{1 + \frac{q^2}{t_{2\pi}}} + \frac{f_{3\pi}(0)}{1 + \frac{q^2}{t_{3\pi}}} .$$

Thus, if we define

$$\beta = \frac{f_{3\pi}(0)}{f_{2\pi}(0)} ,$$

then for small $x(\equiv -q^2)$,

$$f(x) \cong f(0)\{1 + (\frac{1}{1+\beta})(\frac{1}{t_{2\pi}} + \frac{\beta}{t_{3\pi}})x\} . \tag{8}$$

Thus, if the 2π and 3π states make comparable contributions with opposite signs (as they do in the neutron charge structure) and $t_{2\pi} \neq t_{3\pi}$, it is possible that the parameter α defined in Eq. (7) may be negative.[9] An attempt is in progress to determine β from other experimental data involving the 3π state.

The author wishes to thank Professor Geoffrey F. Chew for his guidance, constant advice, and encouragement throughout this work. He also thanks Professor Richard J. Eden, Professor Norman M. Kroll, Dr. James S. Ball, Dr. William R. Frazer, and Dr. David Y. Wong for helpful discussions.

REFERENCES

1. W. R. Frazer and J. R. Fulco, Phys. Rev. Letters 2, 365 (1959); Phys. Rev. 117, 1609 (1960).
2. G. F. Chew, Phys. Rev. Letter 4, 142 (1960).
3. M. L. Goldberger and S. Treiman, Nuovo cimento 9, 451 (1958).
4. We use the fundamental metric tensor such that $-p^2 = M^2$. Units are used in which $\hbar = c = \mu = 1$, where μ is the mass of the pion.
5. How-sen Wong, Phys. Rev. Letters 5, 70 (1960).
6. James S. Ball, The Application of the Mandelstam Representation to Photoproduction of Pions from Nucleons, UCRL-9172, April 11, 1960.
7. G. Harris, J. Orear, and S. Taylor, Phys. Rev. 106, 327 (1957).
8. S. M. Berman and D. A. Geffen, Electromagnetic Form Factor of the π^0 and the Decay Mode $\pi^0 \rightarrow \gamma + e^+ + e^-$, preprint (no date). These authors have also studied the f function from a different approach and obtained $\alpha \simeq 0.06$.
9. The author is indebted to Dr. N. Samios of Brookhaven National Laboratory for advance communication of his preliminary experimental result that $\alpha = -0.24 \pm 0.16$.

FIGURE CAPTION

Fig. 1. Neutral pion decay. Wavy lines are photons; broken line, pion.



57831-1

57831-1