

**Reply to Comment on "Coherent p
0 photoproduction in bulk matter at high energies"
by T. C. Rogers and M. Strikman**

E. Couderc, S. Klein

This manuscript has been authored by an author at Lawrence Berkeley National Laboratory under Contract No. DE-AC02-05CH11231 with the U.S. Department of Energy. The U.S. Government retains, and the publisher, by accepting the article for publication, acknowledges, that the U.S. Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for U.S. Government purposes.

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor The Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or The Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or The Regents of the University of California.

Reply to Comment on “Coherent ρ 0 photoproduction in bulk matter at high energies” by T. C. Rogers and M. Strikman

Elsa Couderc

*Lawrence Berkeley National Laboratory, Berkeley CA 94720 USA,
and Departement de physique, Ecole Normale Supérieure, Paris, France.*

Spencer Klein

*Lawrence Berkeley National Laboratory, Berkeley CA, 94720, USA
(Dated: January 27, 2010)*

In their interesting comment [1] on “Coherent ρ 0 photoproduction in bulk matter at high energies” [2], Rogers and Strikman point out that, at high energies, $q\bar{q}$ dipoles with small separations (d) become more important, and that most of the growth of the cross-section is “driven by the increasingly large contributions from small size (high mass) configurations” [3]; at photon energies of 10^{20} eV, over half of the total cross-section is due to dipoles smaller than 0.25 fm. They state that charm production will increase, and may be as much as 30% of the cross-section.

The coherent photoproduction of heavier states requires higher energies than coherent ρ photoproduction, because the formation length scales as $1/M^2$. For the J/ψ , the required photon energy is 14 times higher than for the ρ . We agree that higher-mass states become important at higher energies. However, at this point, additional factors come into play; as we note after Eq. (7) [2], our calculation is only properly normalized when the conversion probability is relatively small. At the energies where coherent production of high mass states is possible, the coherent ρ production probability is large, and it is necessary to consider reverse reactions such as vector meson ‘back-propagation’ into real photons. The diagonal transitions found in generalized vector meson dominance,

such as from a ρ to a ϕ or J/ψ may also become important, and a recursive technique appears needed to solve the problem [4].

To the extent that the higher mass states limit the growth of the low-mass cross-section, coherent ρ photoproduction will be smaller than we calculated. However, the high-mass fractions quoted by Rogers and Strikman are considerably smaller than the fractional difference between the two hadronic models we considered: the Glauber model [5] and the ERR calculations [6]. The presence of higher mass states in our Glauber calculation would not significantly alter the incoherent photonuclear cross-section in our Fig. 1. However, the subsequent ρ coherent cross-section would be lower, between the Glauber and ERR curves.

In the near future, it should be possible to measure the ρ and J/ψ photoproduction cross-section up to quite high energies (PeV photons in the target frame) using ultra-peripheral ion collisions at the LHC [7], thus reducing these uncertainties, and, more generally, search for signs of a rise in the coherent charm cross-section.

This work was funded by the U.S. National Science Foundation and the U.S. Department of Energy under contract number DE-AC-76SF00098.

[1] T. C. Rogers and M. Strikman, preceding comment.

[2] E. Couderc and S. Klein, Phys. Rev. Lett. **103**, 062504 (2009).

[3] T. Rogers and M. Strikman, J. Phys. **G32**, 2041 (2006).

[4] W. Fetscher *et al.*, Z. Phys. **C72**, 543 (1996).

[5] S. Klein and J. Nystrand, Phys. Rev. **C60** 014903 (1999).

[6] R. Engel, J. Ranft and S. Roesler, Phys. Rev. **D55**, 6957 (1997).

[7] K. Hencken *et al.*, Phys. Rept. **458**, 1 (2008); C. Bertulani *et al.*, Ann. Rev. Nucl. Part. Sci. **55**, 271 (2005).