

Toward an implicit Drift-Lorentz mover*

R.H. Cohen, A. Friedman, D. P. Grote, LLNL, Livermore, CA, USA, and J.-L. Vay, LBNL, Berkeley, CA USA

In order to efficiently perform particle simulations in systems with widely varying magnetization, we have developed a “drift-Lorentz mover”, which interpolates between full particle dynamics and drift kinetics in such a way as to preserve a physically correct gyroradius and particle drifts for both large and small ratios of the timestep to the cyclotron period.[1]. We are now adding implicitness to the mover and the associated field solver in order to extend the mover’s applicability to systems with plasma frequency exceeding the cyclotron frequency. A first step was adding the polarization charge to the field solver and a two-time-level predictor corrector procedure [2]. We outline here two approaches to adding further implicitness. In both, we add a direct-implicit algorithm to the Lorentz portion of the mover; the drift portion can then be treated explicitly, retaining the predictor-corrector procedure from [2], or implicitly, with a modified predictor-corrector procedure. We describe the algorithms, stability analyses, and status of implementation.

[1] R.H. Cohen, A. Friedman, M. Kireeff Covo, *et. al.*, Phys. Plasmas **12**, 056708 (2005).

[2] R.H. Cohen, A. Friedman, D.P. Grote and J.-L. Vay, Nucl. Inst. Methods A **577**, 52 (2007).

* Work performed for the U.S. Department of Energy by U.C. LLNL and U.C. LBNL under contracts W7405-ENG-48 and DE-AC02-05CH11231.