Reversible Sigma C-C Bond Formation Between Phenanthroline Ligands Activated by $(C_5Me_5)_2Yb$

Grégory Nocton,*[† ‡] Wayne W. Lukens,[§] Corwin H. Booth,[§] Sergio S. Rozenel,[‡ §] Scott A. Medling, [§] Laurent Maron,[⊥] and Richard A. Andersen*[‡,§]

[†] Laboratoire de Chimie Moléculaire, CNRS, Ecole Polytechnique, Palaiseau, France.
[‡] Department of Chemistry, University of California, Berkeley, California, 94720, USA.
[§] Chemical Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, California, 94720, USA.
[⊥] LPCNO, UMR 5215, Université de Toulouse-CNRS, INSA, UPS, Toulouse, France.

Abstract. The electronic structure and associated magnetic properties of the 1,10-phenanthroline adducts of Cp^*_2Yb are dramatically different from those of the 2,2'-bipyridine adducts. The monomeric phenanthroline adducts are ground state triplets that are based upon trivalent Yb(III), f^{13} and (phen·-) that are only weakly exchange coupled, which is in contrast to the bipyridine adducts whose ground states are multiconfigurational, open-shell singlets in which ytterbium is intermediate valent (*J. Am. Chem. Soc*, **2009**, *131*, 6480; *J. Am. Chem. Soc*, **2010**, *132*, 17537). The origin of these different physical properties is traced to the number and symmetry of the LUMO and LUMO + 1 of the heterocyclic amine ligands. The bipy·- has only one π^*_1 orbital of b_1 symmetry of accessible energy but phen·- has two π^* orbitals of b_1 and a_2 symmetry that are energetically accessible. The carbon p_π-orbitals have different nodal properties and coefficients and their energies and therefore populations change depending on the position and number of methyl substitutions on the ring. A chemical ramification of the change in electronic structure is that $Cp^*_2Yb(phen)$ is a dimer, when crystallized from toluene solution, but a monomer when sublimed at 180-190 °C. When 3,8-Me_2phenanthroline is used, the adduct $Cp^*_2Yb(3,8\text{-}Me_2phen)$ exist in the solution in a dimer-monomer equilibrium in which ΔG is near zero. The adducts with 3-Me, 4-Me, 5-Me, 3,8-Me_2 and 5,6-Me_2-phenanthroline are isolated and characterized by solid state X-ray crystallography, magnetic susceptibility and L_{III}-edge XANES spectroscopy as a function of temperature and variable temperature ^1H NMR spectroscopy.

Introduction.

The concept of a ligand in a metal compound acting as a single-electron acceptor is a topic of much recent interest [*eg.* see "Forum" in volume 50 (issue 20) of *Inorganic Chemistry]*[1]. The accessibility of an empty orbital on a ligand in a coordination complex was originally referred to as a "non-innocent" ligand but this terminology does not clearly distinguish between metal ligand back-bonding in which a pair of electrons is transferred to an empty orbital and metal-to-ligand charge transfer (MLCT) where a single electron is transferred to an empty ligand orbital, and the ligand is referred to as a "redox non-innocent" or "redox active" ligand. The latter process generates an electron-transfer complex in which an electron resides in the ligand LUMO and an electron hole remains on the metal-based orbital. The ground state electronic structure is then determined by how the biradical correlates the two electrons forming either a triplet state ($M_S = 1$) or an open-shell singlet state ($M_S = 0$).

Complexes of d-transition metals with redox active ligands have been extensively and intensively studied.[1] In contrast, complexes of the f-block metals, although they are known, are not as well studied, with most of the work only appearing recently.[2-13] The 2,2'-bipyridine adducts of Cp^*_2Yb, in particular, have been shown by experimental and computational methodologies to have multiconfigurational open-shell singlet ground states in which ytterbium is intermediate valent.[4,5] In this context, an article by Scarborough and Wieghardt[14] is particularly informative as they systematize and classify the often confusing and/or contradictory literature of the 2,2'-bipyridine and related adducts of d-transition metal metallocenes using a density functional theory (DFT) broken-symmetry (BS) methodology. A comparison between the electronic ground state of $(C_5H_5)_2Ti(bipy)$[15] and $(C_5Me_5)_2Yb(bipy)$ is enlightening. Both adducts have an open-shell singlet ground state (S)[5,14] but the triplet state (T) in $Cp_2Ti(bipy)$ lies close enough to the ground state ($-2J = 600$ cm^{-1}) that it is a spin

equilibrium molecule, $S(M_S = 0) \rightleftharpoons T(M_S = 1)$, whereas the triplet in $Cp^*_2Yb(bipy)$ lies 0.28 eV (calculated) or $-2J = 0.11$ eV (920 cm^{-1}, experimental)[11] above the open-shell singlet state and the triplet is not significantly populated at 300 K. These physical properties show that strong exchange coupling does indeed occur in these 4f-block metal compounds.

Although bipyridine and related ligands, such as diazadienes, attached to d- and f-block metallocences have attracted the most attention, adducts with 1,10-phenanthroline have been largely ignored. In this article, it is shown that the ground state of $Cp^*_2Yb(phen)$ is a triplet (T), in contrast to the open-shell singlet ground state of $Cp^*_2Yb(bipy)$. One chemical ramification of the triplet electronic configuration is that the phenanthroline ligands in the individual monomer units are coupled by formation of C-C sigma bond at the 4,4'-positions resulting in a dimer. The related adduct, $Cp^*_2Yb(3,8\text{-}Me_2\text{-phen})$ exists in solution as a dimer $\rightleftharpoons$ monomer equilibrium, and analysis of solid state structure and ^1H NMR spectra show that C-C bond is long (1.592(16) Å) and weak ($\Delta H = -8$ kcal.mol^{-1}).

The thermochemistry for a dimer $\rightleftharpoons$ monomer equilibrium, $D \rightleftharpoons 2M$, where M is an organic σ-radical, σ-R, and D is the dimer , $\sigma\text{-}R_2$, is of fundamental interest since the value of ΔH is the bond dissociation enthalpy, BDE, for the $\sigma\text{-}R_2$ single bond. Although BED's for organic compounds are well known, only a few examples of BDE's for a specific σ-carbon-carbon single bond and the associated bond distance in the dimer are known. The oldest dimer-monomer equilibrium is that of Gomberg's dimer, for which the value of ΔH of 11 kcal.mol^{-1} has been measured,[16-18] is not a simple $\sigma\text{-}R_2 \rightleftharpoons 2\sigma\text{-}R$ dissociation due to the structure of the dimer. Recently, the thermochemistry of the σ-dimerization of the phenalenyl σ-dimer and the related aza-analogue have been measured.[19-21] The ΔH values of $D \rightleftharpoons 2M$ for **I** and **II** in CCl_4 are 10 kcal.mol^{-1} and 11 kcal.mol^{-1}, respectively, and the associated ΔS

values are 15 and 18 cal.mol^{-1}.K^{-1} respectively. The C-C bond length in the copper bis(trifluoroactetylacetonate) complex of the dimer of **II** is 1.58 Å. This value is identical to that calculated for the σ-C-C distance in the σ-dimer of **I**, for which the calculated value of the BDE is 16 kcal.mol^{-1}. More recently, the ΔH value for the D ⇌ 2M, M is 2,6-di-tert-butyl-4-methoxyphenoxy radical of 6 kcal.mol^{-1} has been obtained along with the σ-C-C distance in the dimer of 1.605(2) Å.[22] The reversible coupling of two pyridine ligands in a β-diketiminate iron complex has recently been published in which the C-C distance of 1.563(6) Å was measured and a ΔH value of 11 kcal.mol^{-1} was estimated.[23]



I **II**

The dimer of the phenalenyl radical also forms π-dimers when Me_3C groups are attached to the arene rings.[24-26] Although the ΔH values are similar to the σ-dimers, the π-C-C distances are much longer, as they range from 3.201(8) Å to 3.323(6) Å in the π-dimer of 1,4,7-$(Me_3C)_3C_{12}H_6$ in D_{3d} symmetry.

This article shows that single electron transfer (SET) to a π-symmetry LUMO of a close-shell ligand results in a stretched and weakened C-C bonds, σ-R_2 for which ΔG ~ 0.

Results.

Synthesis. The syntheses of $Cp^*_2Yb(phen)$ and $[Cp^*_2Yb(phen)]I$ were reported in an earlier paper,[27] and the new neutral adducts are prepared in a similar manner, Eq. 1. Some physical properties of the adducts are shown in Table 1.

$$Cp^*_2Yb(OEt_2) + \text{1,10-phenanthroline} \rightarrow Cp^*_2Yb(phen) + OEt_2 \qquad \textbf{(1)}$$

The neutral phen adducts of Cp^*_2Yb, Cp^*_2Yb(x-phen), where x is H, 3-Me, 4-Me and 5-Me (The atom numbering system is shown in the legend to Table 1) are sparingly soluble in toluene and tetrahydrofuran, they decompose in dichloromethane, and may be crystallized from a dilute solution of warm toluene. The 3,8-Me_2phen adduct is somewhat more soluble in toluene, but all of the neutral adducts are much less soluble than the 2,2'-bipyridine adducts described in earlier papers.[4,28] The solid-state and solution-state physical properties of the adducts are quite different and these properties are described in the separate sections that follow.

Table 1. Solid state properties of the Cp^*_2Yb adducts **1-7**. The legend shows the numbering scheme for the carbon positions on phenanthroline.

Compound	color	m.p (°C)	IR (cm^{-1})	μ_{eff} (300K)
Cp^*_2Yb(phen) (**1**)	deep blue	297-300	1610, 1590, 1550, 859	4.00[(a)]
$[Cp^*_2Yb(phen)]^+I^-$ (**2**)	red-brown	175-180	1622, 1518, 855	4.54
Cp^*_2Yb(3,8-Me_2phen) (**3**)	dark red	286-288	1625, 1573, 1461, 799	4.10
Cp^*_2Yb(3-Mephen) (**4**)	dark purple	270-272	1612, 1554, 880	3.92
Cp^*_2Yb(4-Mephen) (**5**)	dark purple	254-256	1618, 1512, 1445, 800	3.92
Cp^*_2Yb(5-Mephen) (**6**)	dark purple	280-283	1626, 1578, 1504, 878	3.95
Cp^*_2Yb(5,6-Me_2phen) (**7**)	deep purple	285-287	1605, 1584, 1480, 804	3.68

a) crystallized, the value for the sublimed compound is ~ 4.5 μ_B (300 K)



Physical Properties, Solid State.

Magnetism.

Plots of the effective magnetic moment, μ_{eff}, as a function of temperature for the six neutral adducts **1**, **3-7**, each obtained by crystallization from toluene and the cation $[Cp^*_2Yb(phen)]^+I$ (**2**), are shown in Figure 1 (Plots of χ, χT, $1/\chi$ and χT as a function of temperature are available in SI). The striking feature of the data in Figure 1 is that the curves have a similar shape that differs mainly by a scaling factor for the neutral and cationic adducts, although the overall magnitude of the μ_{eff} value for **7** is noticeably smaller over the entire temperature range. This similarity is in contrast to what was observed for the various bipyridine adducts of Cp^*_2Yb described in earlier work in which the neutral bipy adducts have substantially lower μ_{eff} values relative to their cationic derivatives.[4,5,28]



Figure 1a. Plot of the effective magnetic moment, μ_{eff}, as a function of temperature for **1-6** in the 2-300 K temperature range. These adducts are obtained by crystallization.



Figure 1b. Plot of the effective magnetic moment, μ_{eff}, as a function of temperature for **1-monomer** after sublimation (red unflilled dots), **1-dimer** after crystallization (blue filled dots) and **7** after crystallization (green squares) in the 2-300 K temperature range.

The effective magnetic moments of most of the neutral adducts have a slight temperature dependence as μ_{eff} decreases from about 4 μ_B to 3.2 μ_B as the temperature decreases from 300 K to 5 K. The value of μ_{eff} is somewhat lower than expected for two uncorrelated spin carriers Yb(III), $^2F_{7/2}$, and phen radical anion, $^2S_{½}$, for which a value of 4.83 μ_B is expected at 300 K. The value of 4.54 μ_B is expected for an isolated Yb(III) ion, $^2F_{7/2}$, in agreement with the value found for the cationic adduct, $[Cp^*_2Yb(phen)]^+I^-$ (**2**), at 300 K. The similarity of the magnetic moments of the neutral adducts in Figure 1a with that of the cation begets the question of the identity of the anion in these neutral adducts. Although it has similar magnetic behavior, μ_{eff} of **7** decreases from 3.5 μ_B to below 2.5 μ_B as the temperature decreases from 300 K to 5 K.

Yb L_{III}-Edge XANES Spectra.

Yb L_{III}-edge XANES spectra of the six neutral adducts, **1-dimer**, **3-7**, are shown in Figure 2 for data collected at both 30 K and 300 K. No significant change is observed over this

temperature range. All the spectra are characterized by a single white-line feature at about 8946 eV. This feature is indicative of the f^{13} configuration. Another peak at 8939 eV, indicating the f^{14} configuration as shown in Figure 2 by data on the intermediate valent Cp^*_2Yb(4-Me-bipy) compound,[4] **8**, is not clearly visible in the phen adduct spectra. These spectra were fit with methods described previously,[5] giving estimates of n_f as shown in Table 2. The Yb is these samples is found to be close to trivalent, Yb(III), with an f-hole occupancy $n_f \approx 1$.



Figure 2. Yb L_{III}-edge XANES spectra for **1-dimer**,**3-7** at 30 K (solid) and at 300 K(dotted). Also shown are previous data[4] on Cp^*_2Yb(4-Me-bipy) ($n_f = 0.79$) for comparison. The shoulder at 8939 eV below the main peak at 8946 eV is indicative of the Yb(II) contribution, which is clearly seen in the bipy adduct data. The strong overlap of all the measured phen adduct data emphasizes the overall similarity in f-orbital occupancy. The small shoulder for each of the phen adducts indicates these samples are close to trivalent Yb.

Table 2: The estimated f-hole occupancy, n_f, determined by Yb L_{III}-edge XANES measurements. No temperature dependence was observed between 30 K and 300 K. The estimated absolute error in the last digit n_f is shown in parentheses; the random error between separate traces is much smaller.

Compound	$\mathbf{n_f}$
$Cp^*_2Yb(phen)$ (**1**)	0.99(3)
$Cp^*_2Yb(3,8\text{-}Me_2phen)$ (**3**)	0.96(3)
$Cp^*_2Yb(3\text{-}Mephen)$ (**4**)	0.95(3)
$Cp^*_2Yb(4\text{-}Mephen)$ (**5**)	0.98(3)
$Cp^*_2Yb(5\text{-}Mephen)$ (**6**)	0.97(3)
$Cp^*_2Yb(5,6\text{-}Me_2phen)$ (**7**)	0.96(3)
$Cp^*_2Yb(4\text{-}Me\text{-}bipy)$ (**8**)	0.79[4]

It is clear from Figure 2 that the Cp^*_2Yb fragments are based upon Yb(III), f^{13}, which again begets the question raised from the magnetic data about the identity of the anion in the neutral adducts. The genesis of an answer is indicated by the EPR spectra.

EPR Spectra.

The EPR Spectra at 2 K of $Cp^*_2Yb(phen)$ (**1-dimer**) and $Cp^*_2Yb(3,8\text{-}Me_2phen)$ (**3**) are shown in Figure 3(a) and that of $[Cp^*_2Yb(phen)]I$ (**2**) is shown in Figure 3(b). EPR spectra of the monomethyl adducts of Cp^*_2Yb, **4-6** are shown in Supporting Information. The g-values are given in Table 3. Because of the high sensitivity of EPR, it is important to compare the EPR and magnetic susceptibility results to determine whether they are consistent. At the temperature at which the EPR spectra are obtained (~2 K), only the ground state is occupied in most cases. The effective magnetic moment of the ground state is determined from the EPR g-values using $\mu_{eff} = 0.5\ (g_1^2+g_2^2+g_3^2)^{1/2}$, which may be compared to the magnetic susceptibility data by extrapolating χT to 0 K then determining μ_{eff} (0 K). As shown in Table 3, the effective magnetic moments determined from the EPR g-values are consistent with

those determined by magnetic susceptibility, so the EPR spectra can be assigned to the Yb complexes rather than to impurities.



Figure 3: EPR spectra recorded in the solid state (powder) at 2 K for a) $Cp^*_2Yb(phen)$ (**1,** solid black line) and $Cp^*_2Yb(3,8\text{-}Me_2phen)$ (**3,** dashed red line) and b) **[**$Cp^*_2Yb(phen)]I$ (**2**).

The nature of the ligand coordinated to the $[Cp^*_2Yb]^+$ fragment has a significant effect on the EPR spectrum as previously illustrated by $[Cp^*_2Yb(bipy)]I$ and $Cp^*_2Yb(bipy)$.[29] In $[Cp^*_2Yb(bipy)]I$, $[Cp^*_2Yb]^+$ is coordinated by a neutral, closed-shell bipy ligand, and the complex has EPR parameters similar to those of **2**. On the other hand, in $Cp^*_2Yb(bipy)$, $[Cp^*_2Yb]^+$ is coordinated by the bipy radical anion, and $Cp^*_2Yb(bipy)$ is EPR silent. Based on these observations, the EPR spectra of **1-dimer-6** are consistent with trivalent, $[Cp^*_2Yb]^+$ fragments coordinated by closed-shell ligands, and the lack of an EPR spectrum for **7** is consistent with the presence of the ligand that is a radical anion. The fact that **[**$Cp^*_2Yb(phen)]I$ (**2**), is EPR active is unsurprising – the $[Cp^*_2Yb]^+$ fragment is obviously coordinated by a neutral phenanthroline ligand, and the charge is balanced by iodide. However, the fact that **1-dimer** and **3-6** are EPR-active is surprising since it means that these complexes do not contain radical anions in sharp contrast to $Cp^*_2Yb(bipy)$.

Table 3. EPR data for **1-7**.

	EPR data	μ_{eff} (EPR)[a]	μ_{eff} (0 K)[b]
$Cp^*_2Yb(phen)$ (**1-dimer**)	$g_1 = 6.85$, $g_2 = 1.47$, $g_3 = 1.40$	3.57 μ_B	3.21 μ_B
$[Cp^*_2Yb(phen)]^+I^-$ (**2**)	$g_1 = 6.70$, $g_2 = 1.92$, $g_3 = 1.21$	3.54 μ_B	3.49 μ_B
$Cp^*_2Yb(3,8\text{-}Me_2phen)$ (**3**)	$g_1 = 7.01$, $g_2 = 1.41$, $g_3 = 1.33$	3.64 μ_B	3.46 μ_B
$Cp^*_2Yb(3\text{-}Mephen)$ (**4**)	$g_1 = 7.02$, $g_2 = 1.21$, $g_3 = 1.21$	3.61 μ_B	3.29 μ_B
$Cp^*_2Yb(4\text{-}Mephen)$ (**5**)	$g_1 = 6.47$, $g_2 = 1.31$, $g_3 = 1.31$	3.37 μ_B	3.28 μ_B
$Cp^*_2Yb(5\text{-}Mephen)$ (**6**)	$g_1 = 6.45$, $g_2 = 1.42$, $g_3 = 1.21$	3.35 μ_B	3.19 μ_B
$Cp^*_2Yb(5,6\text{-}Me_2phen)$ (**7**)	EPR silent	-	2.33 μ_B

a) $\mu_{eff}(EPR) = (1/2)(g_1^2+g_2^2+g_3^2)^{1/2}$
b) μ_{eff} (0 K) was determined by using a linear fit of χT from 12 K to 45 K, and determining μ_{eff} from χT extrapolated to 0 K.

X-Ray Crystal Structures.

The nature of the bonding in these complexes and the reason why **1-dimer**, **3-6** are EPR active in contrast to **7** is clarified by their crystal structures. Although the phen adduct and substituted phen adducts are sparingly soluble in hydrocarbons and they have high melting points, Table 1, the phen adduct sublimes at 180 – 190 °C in an ampoule sealed under reduced pressure. The sublimation temperature must be maintained in this 10 °C range, since heating to a higher temperature results in substantial decomposition. In the 180 – 190 °C range, a small number of well formed crystals grow during a week, which are suitable for X-ray diffraction. The ORTEP in Figure 4 shows that the sublimed crystals are well separated monomers of Cp*Yb(phen) (**1-monomer** is used to distinguish the sublimed compound from the crystallized compound, labeled as **1-dimer**). The ORTEP of $Cp^*Yb(5,6\text{-}Me_2phen)$ (**7**) in Figure 5 shows the three independent molecules in the unit cell of the monomeric adduct obtained by crystallization from cyclohexane. The crystal structure of $Cp^*_2Yb(5,6\text{-}Me_2phen)$, obtained by sublimation in a sealed ampoule under reduced pressure at 195 °C over two months, labeled **7, sublimed** is shown in Figure 6a, along with a crystal packing diagram of

two molecules in the unit cell is shown in Figure 6b. These results contrast with the X-ray crystals structures of crystallized Cp*Yb(phen) (**1-dimer, crystallized**) and Cp*Yb(3,8-Me_2phen) (**3**), which are dimers. Crystals of the latter two adducts, obtained from toluene solution are deep blue and deep purple in color, respectively. ORTEP's of **1-dimer** and **3** are shown in Figure 7 and Figure 8, respectively. It is clear that the anionic partner is derived by dimerization of two phenantholine radical anions by formation of a C-C bond at the 4,4'-positions, forming the diamagnetic dianionic partner.



Figure 4: ORTEP for sublimed $Cp^*_2Yb(phen)$ (**1-monomer, sublimed**) (thermal ellipsoids at 50% level). Ytterbium atom is in green, nitrogen atoms in blue and carbon atoms in grey. All non-hydrogen atoms are refined anisotropically and the hydrogen atoms are placed in calculated positions but not refined. Hydrogen atoms have been omitted for clarity.



Figure 5: ORTEP for crystallized $Cp^*_2Yb(5,6\text{-}Me_2phen)$ (**7, crystallized**) (thermal ellipsoids at 50% level) showing the three independent molecules in the unit cell. Ytterbium atom is in green, nitrogen atoms in blue and carbon atoms in grey. All non-hydrogen atoms are refined anisotropically and the hydrogen atoms are placed in calculated positions but not refined. Hydrogen atoms have been omitted for clarity.



Figure 6a. ORTEP for crystallized $Cp^*_2Yb(5,6\text{-}Me_2phen)$ (**7, sublimed**) (thermal ellipsoids at 50% level). Ytterbium atom is in green, nitrogen atoms in blue and carbon atoms in grey. All non-hydrogen atoms are refined anisotropically and the hydrogen atoms are placed in calculated positions and refined isotropically. Hydrogen atoms have been omitted for clarity. **6b.** A portion of the packing diagram showing two molecules in the unit cell, showing the shortest C….C contact distances are between C(8) and C(9) methyl groups on the Cp*-ring and C(34) and C(35) methyl groups on the $5,6\text{-}Me_2phen$ ligand of 3.634 Å and 3.792 Å, respectively.



Figure 7: ORTEP for crystallized $Cp^*_2Yb(phen)$ (**1**) (thermal ellipsoids at 50% level). All non-hydrogen atoms are refined anisotropically and the hydrogen atoms are placed in calculated positions but not refined. Hydrogen atoms have been omitted for clarity.



Figure 8: ORTEP for $Cp^*_2Yb(3,8\text{-}Me_2phen)$ (**3**). The carbon atom C28 (represented in pink) is refined in two positions C28 (2/3 of occupancy) and C(28B) is located in the plane of the phenanthroline closest to it. Details are in Supporting Information. Toluene molecules have been omitted for clarity.

Bond distances and angles in the known phenanthroline adducts of ytterbocenes are shown in Table 4. The Yb-C(Cp*) distances in the neutral and cationic adducts of Cp^*_2Yb are identical, given the large range in the individual values. The Yb-C(Cp*) distances are

approximately 0.1 Å shorter than in $Cp^*_2Yb(py)_2$[30] and $[1,3\text{-}(Me_3Si)_2C_5H_3]Yb(phen)$[27] consistent with the higher oxidation number of ytterbium in the phenanthroline adducts of Cp^*_2Yb. It is particulary noteworthy that the Yb-C(Cp*) distances are identical in the monomeric and dimeric forms of $Cp^*_2Yb(phen)$.

The Yb-N distances, however, show significant differences in the neutral adducts depending upon whether they are monomers or dimers. In monomeric $Cp^*_2Yb(phen)$, the average Yb-N distance is 2.311±0.002 Å, identical to that in $Cp^*_2Yb(5,6\text{-}Me_2phen)$ of 2.318±0.007 Å. In the dimeric forms of $Cp^*_2Yb(phen)$ and $Cp^*_2Yb(3,8\text{-}Me_2phen)$, the average Yb-N distances of 2.322±0.018 Å and 2.324±0.016 Å are the same as those found in the monomers but the individual distances differ by 0.08 to 0.06 Å, respectively. Thus, the Yb-N(1) distances of 2.358(5) Å and 2.366(4) Å in **1-dimer** and **3**, respectively, are similar to those found in $[Cp^*_2Yb(phen)]I$ of 2.339(8) Å and 2.382(8) Å,[27] but the Yb-N(2) distances of 2.285(4) Å and 2.301(5) Å, respectively, are shorter and indicative of an amide-nitrogen to Yb(III) bond. This conjecture, *viz.*, that longer Yb-N bond lengths in the dimers are due to a Yb(III)-N (dative) and the shorter distances are due to a Yb(III)-N (anionic) bond is supported by comparison between the C-N and C-C bond distances in the individual pyridyl rings in **1-dimer** and **3** shown in Table 7 (see below). The trends in these bond lengths in the phenanthroline rings in both dimers are consistent with the formulation of the N(1) pyridine ring as a neutral pyridine ring and the N(2) pyridine ring as a quinoid resonance structure in which N(2) carries a negative charge.

Table 4. Bond lengths (Å) and Angles (deg) for the phenanthroline adduct of the Ytterbocenes.

Compound	Yb-C(Cp*)[(a)] ave, Å	Yb-C_t ave, Å	Yb-N ave, Å	Refs
$Cp^*_2Yb(phen)$, crystallized, **1-dimer**	2.617 ± 0.016	2.33	2.285(4) 2.358(5)	This work
$Cp^*_2Yb(phen)$, sublimed, **1-monomer**	2.610 ± 0.008	2.33	2.311 ± 0.002	This work
$Cp^*_2Yb(3,8\text{-}Me_2phen)$, crystallized (dimer**)**	2.63 ± 0.02	2.33	2.301(5) 2.366(4)	This work
$Cp^*_2Yb(5,6\text{-}Me_2phen)$, crystallized (monomer) Molecule 1 Molecule 2 Molecule 3	 2.62 ± 0.02 2.63 ± 0.01 2.63 ± 0.01 ave 2.63	 2.33 2.33 2.33	 2.330 ± 0.005 2.322 ± 0.005 2.313 ± 0.005 ave 2.322	This work
$Cp^*_2Yb(5,6\text{-}Me_2phen)$, sublimed (monomer)	2.620 ± 0.005	2.33	2.310 ± 0.009	This work
$[Cp^*_2Yb(phen)]I$	2.61 ± 0.01	2.31	2.360 ± 0.011	ref[27]
$[1,3\text{-}(Me_3Si)_2C_5H_3]Yb(phen)$	2.72 ± 0.02	2.43	2.501 ± 0.007	ref[27]
$Cp^*_2Yb(py)_2$	2.74 ± 0.04		2.565 ± 0.005	ref[30]

(a) the ± values are average deviation from the mean values.

These bond lengths are in sharp contrast to those observed in the monomeric adducts, $Cp^*_2Yb(phen)$ and $Cp^*_2Yb(5,6\text{-}Me_2phen)$, as shown in Table 8 (see below). In these two adducts, the small differences between C-N and C-C distances in the pyridine rings containing N(1) and N(2) are consistent with their formulation as delocalized radical anions.

In the $Cp^*_2Yb(x,x'\text{-bipy})$ adducts, the changes in the C(2)-C(2') provide qualitative insights into the ground state electronic structure of these adducts.[4,5] In these charge-transfer complexes, the SOMO of the bipyridine radical-anion has b_1 symmetry (in C_{2v} symmetry) and the C(2)-C(2') distance, represented by A in Scheme 1, shortens relative to the equivalent distance in the free bipyridine ligand, since these C-pπ-orbitals are a bonding combination. A related analysis of the C-N and C-C distances in the phenanthroline adducts is not as straightforward since (i) the distance represented by A is part of a rigid-ring system and (ii) the LUMO and LUMO+1 orbitals of b_1 and a_2 symmetry (in C_{2v} symmetry), respectively, are close in energy, Figure 9, and population of these bonding and antibonding orbitals results in

a complex pattern of bond length alterations since these pπ-orbitals have different nodal properties and coefficients. However, a systematic examination of all the anticipated changes when either b_1 or an a_2-symmetry orbitals are singly occupied generates an informative pattern. The four pair of distances labeled as C and O; E and M; D and K; G and N in Scheme 1 change in identical ways when either b_1 or a_2 is singly occupied. In contrast the distances labeled A; I; F and L; B and P; H and J change in opposite directions as shown in Tables 5 and 6.



Figure 9. Relative energy diagram of the b_1 and a_2 symmetry orbitals in bipy radical anion and phen radical anion as a function of the Coulomb integral on N, h_N.[39]

Scheme 1. b_1 and a_2 representation and bond labeling.



Table 5. Anticipated bond lengths changes in LUMO and LUMO+1 of phenanthroline radical anion.

Bond	Orbital[(a)]	
	LUMO, b_1	LUMO+1, a_2
A	-	+
I	-	+
F,L	-	+
B,P	+	-
H,J	+	-

(a) + means the distance increase, - means the distance decreases, when these orbitals are occupied.

Table 6. Bond lengths (Å) changes in **1-monomer** and **7**.

Bond	$\Delta^{(a,b)}$	
	$Cp^*_2Yb(phen)$ monomer	$Cp^*_2Yb(5,6\text{-}Me_2phen)$ Monomers[(c)]
A	-0.020	-0.036
I	+0.014	-0.010
F,L	+0.041	-0.011
B,P	+0.003	+0.014
H,J	-0.026	-0.015

(a) Δ is the bond length distance in the adduct minus that in the free ligand in Å.
(b) Free phen, ref[31] and free 5,6-Me_2phen, ref[32]
(c) The average change in the four individual molecules.

The pattern of bond length changes in monomeric $Cp^*_2Yb(phen)$ is inconsistent with population of either b_1 or a_2-orbitals but consistent with population of both orbitals. The pattern of bond length alteration in $Cp^*_2Yb(5,6\text{-}Me_2phen)$ is somewhat different from that found in $Cp^*_2Yb(phen)$. This difference implies that the b_1/a_2 ratio is higher in the former adduct and that methyl groups in the 5,6-positions stabilize the b_1 orbital. These inferences are consistent with the calculational results described latter in this manuscript.

The geometry of the phenanthroline ligands in the two dimers, $Cp^*_2Yb(phen)$, **1**, and $Cp^*_2Yb(3,8\text{-}Me_2phen)$, **3**, is similar but the crystallographic details are different (Figure 7 and 8). In **1**, the C(28)-C(28') atoms have well-behaved thermal parameters, see Supporting Information, and these two carbon atoms are refined anisotropically although the hydrogen

atoms attached to them are not included in the refinement. The geometry of the N(1)-ring in the $Cp^*_2Yb(phen)$ is planar while that of the N(2)-ring is non-planar. In the N(2)-ring, the dihedral angle formed by intersection of the two planes defined N(2)C(30)C(29)C(27)C(31) and C(27)C(28)C(29) is 25°, in accord with C(28) being a sp^3-carbon atom. The C(28)-C(27,29) distances in Table 7 are in the range given for Csp^3-Csp^3 bond lengths of 1.507 Å (σ = 0.015 Å).[33] The C – N distances, Table 7, are also in the range of Csp^3-N distances of 1.358 Å (σ = 0.015 Å).[33] These distances and angles in the N(2)-ring indicate that the pyridyl ring is represented by a quinoid distortion and the nitrogen atom carries a negative charge.

Scheme 2.



The orientation of the two phenanthrolyl rings in the crystal structure is shown by the Newman projection down the C(28)C(28') bond, Scheme 2, left-hand drawing. The molecule has C_i symmetry and the inversion center is the mid-point of C(28)C(28'). The Newman projection of another rotomer of C_2 symmetry in which the C(28)-H's have a trans disposition is shown in the right-hand drawing.

Table 7. C-N and C-C bond distances in **1, dimer** and **3**

Cp*$_2$Yb(phen),[c] **1, dimer**				
Ring 1[a]		Ring 2[b]		Δ[e]
Bond[c]	Distance, Å	Bond	Distance, Å	
N(1)C(21)	1.331(7)	N(2)C(30)	1.380(7)	-0.049
N(1)C(32)	1.396(6)	N(2)C(31)	1.374(7)	+0.022
C(21)C(22)	1.393(8)	C(30)C(29)	1.336(8)	+0.067
C(22)C(23)	1.375(9)	C(29)C(28)	1.504(8)	-0.129
C(23)C(24)	1.417(9)	C(28)C(27)	1.500(8)	-0.083
C(24)C(32)	1.387(7)	C(27)C(31)	1.413(7)	-0.024
Cp*$_2$Yb(3,8-Me$_2$phen),[c] **3**				
Ring 1[a]		Ring 2[b]		Δ[e]
Bond[c]	Distance, Å	Bond	Distance, Å	
N(1)C(21)	1.339(6)	N(2)C(30)	1.381(6)	-0.042
N(1)C(32)	1.384(6)	N(2)C(31)	1.374(6)	+0.010
C(21)C(22)	1.399(7)	C(30)C(29)	1.352(7)	+0.047
C(22)C(23)	1.388(7)	C(29)C(28)	1.529(9)	-0.141
C(23)C(24)	1.394(7)	C(28)C(27)	1.547(9)	-0.153
C(24)C(32)	1.419(7)	C(27)C(31)	1.403(7)	+0.016

Table 8. C-N and C-C bond distances in **1, monomer** and **7**.

Cp*$_2$Yb(phen),[d] **1, monomer**				
Ring 1[a]		Ring 2[b]		Δ[e]
Bond	Distance, Å	Bond	Distance, Å	
N(1)C(21)	1.383(6)	N(2)C(30)	1.381(7)	+0.002
N(1)C(32)	1.367(5)	N(2)C(31)	1.365(6)	+0.002
C(21)C(22)	1.371(9)	C(30)C(29)	1.382(8)	-0.011
C(22)C(23)	1.388(9)	C(29)C(28)	1.392(9)	-0.004
C(23)C(24)	1.459(8)	C(28)C(27)	1.445(8)	+0.013
C(24)C(32)	1.414(7)	C(27)C(31)	1.426(6)	-0.012
Cp*$_2$Yb(5,6-Me$_2$phen),[d] **7, monomers** (average)				
Ring 1[a]		Ring 2[b]		Δ[e]
Bond[c]	Distance, Å	Bond	Distance, Å	
N(1)C(21)	1.350(11)	N(2)C(30)	1.352(7)	-0.002
N(1)C(32)	1.377(7)	N(2)C(31)	1.371(15)	+0.006
C(21)C(22)	1.369(12)	C(30)C(29)	1.370(8)	-0.001
C(22)C(23)	1.388(16)	C(29)C(28)	1.380(7)	+0.008
C(23)C(24)	1.40(3)	C(28)C(27)	1.40(2)	0.0
C(24)C(32)	1.423(8)	C(27)C(31)	1.428(10)	-0.005

a) Ring 1 are the atoms in the pyridyl-N(1) ring.
b) Ring 2 are the atoms in the pyridyl-N(2) ring.
c) See Figure 7 and 8 for the atom numbering scheme.
d) See Figure 4 and 6 for the atom numbering scheme.
e) The differences in Å between the distances in ring 1 and ring 2.

The description of the geometry around C(28) in the $Cp^*_2Yb(3,8\text{-}Me_2phen)$, **3** is less straightforward since two positions for the C(28) atom are occupied (C(28) and C(28B)). While solving the structure, a singularity appeared at atom C28. The problem was addressed in two ways: (i) the C28 atom is forced to remain in the mean plane of ring 2 and (ii) a positional disorder model in which C28 and C28B are assigned an occupancy ratio of 0.67:0.33, respectively. Solution (i) led to elongated thermal ellipsoid perpendicular to the mean plane of ring 2, Figure S22 representing this tentative solution is shown in the Supporting Information. In this representation the C(28)C(28') distance is 3.00 Å. Solution (ii) led to well-behaved thermal ellipsoids for C(28) and C(28B) but their position differs; C(28) is comparable to that of the C(28) atom in **1-dimer** with elongated C(28)-C(29,27) distances (see Table 7) of 1.529(9) Å and 1.547(9) Å, respectively, and a C(28)-C(28') distance of 1.592(16) Å. This is compatible with the presence of a σ-dimer, that is the bond between C(28)C(28') is classified as a σ-bond between two sp^3 carbons. On the other hand, C(28B) is found close to the mean plane of ring 2 with C(28B)C(29,27) distances of 1.478(18) Å and 1.417(19) Å and a C(28B)C(28B') distance of 3.39 Å (calculated), compatible with its classification as a π-dimer, that is, a bond formed by interaction between the p_π-orbitals on the sp^2 hybridized carbon atoms. The disorder in $Cp^*_2Yb(3,8\text{-}Me_2phen)$, **3,** may be viewed as the average between these two forms (σ-dimer and π-dimer) in which the energy difference between them is small.

Vis-NIR Spectra.

The Vis-NIR spectra in the 400-950 nm range in toluene solution at 20 °C for the crystallized adducts of 4-Mephen (**5**) and 5-Mephen (**6**) are similar to the spectrum of $Cp^*_2Yb(phen)$ (**1**) reported in an earlier paper.[27] The spectra are available in SI. Morris and co-workers have given a detailed analysis of the solution spectra from 400 nm to 2500 nm of $[Cp^*_2Yb(phen)]^{0,+}$.[6] The key point that emerges from these spectroscopic studies is that the

spectra of the neutral adducts contain features associated with the phenanthroline radical anion, an absorption around 500 nm, along with f-f transitions at longer wavelengths.

^{1}H NMR Spectra.

The chemical shifts in C_6D_6 or C_7D_8 at 300 K for the neutral adducts are given and assigned in Table 9. The $Cp^*_2Yb(phen)$, $Cp^*_2Yb(3,8\text{-}Me_2phen)$ and $Cp^*_2Yb(5,6\text{-}Me_2phen)$ adducts have four resonances due to the phenanthroline ligands in the general region of δH ~ 100, ~50, ~15 and ~0 ppm, in addition to the Cp* resonance at δH ~ 4 ppm. The resonances that can be assigned with certainty are those at δH ~ 15 ppm since these are replaced by a resonance due to the Me-groups at δH ~ -10 ppm in **3**, and therefore the δH ~ 15 ppm resonance is due to $\delta_{3,8}$. The resonances at δH ~ 0.5 ppm are replaced by a resonance due to the Me-groups at δH ~ 0.03 ppm in **7**, and therefore the δH ~ 0.5 ppm is due to $\delta_{5,6}$. The most deshielded resonances are assigned to $\delta_{2,9}$ since these are closest to the paramagnetic center, and the remaining resonances at δH ~ 50 ppm are due to $\delta_{4,7}$. The appearance of four phen resonances shows that the adducts have C_{2v} symmetry in solution at 300 K. The chemical shifts of $Cp^*_2Yb(3,8\text{-}Me_2phen)$ depend upon the solvent; in THF the most downfield resonance in C_7D_8 moves upfield by about 20 ppm while the other resonances shift by a lesser amount (see Experimental Section). The eight resonances in the 5-Mephen adduct are consistent with a single isomer of C_s symmetry at 300 K but those in the 4-Mephen adduct are not observed at 300 K, while only some of the resonances for the 3-Mephen adduct are observed. The low solubility of the neutral adduct precludes a more detailed study with exception of the $3,8\text{-}Me_2phen$ and $5,6\text{-}Me_2phen$ adducts that are somewhat more soluble in THF and toluene.

Table 9. ^{1}H NMR chemical shift in C_6D_6 or C_7D_8 at 300 K for neutral adducts **1**, **3**-**7**.

Compound	2,9	4,7	3,8	5,6	Cp*
$Cp^*_2Yb(phen)$ (**1**)	139.94	47.87	14.02	0.47	4.14
$Cp^*_2Yb(3,8\text{-}Me_2phen)$ (**3**)	95.54	51.07	-10.03 (Me)	3.83	3.36
$Cp^*_2Yb(3\text{-}Mephen)$ (**4**)	121.47 118.38	59.15 57.17	18.69 -9.51 (Me)	- -	3.79
$Cp^*_2Yb(4\text{-}Mephen)$ (**5**)	-	-	-	-	4.03
$Cp^*_2Yb(5\text{-}Mephen)$ (**6**)	138.72 138.59	47.92 39.33	14.18 11.40	0.06 -0.58 (Me)	4.09
$Cp^*_2Yb(5,6\text{-}Me_2phen)$ (**7**)	137.44	44.10	14.66	0.03 (Me)	3.95

Variable temperature ^{1}H NMR of $Cp^*_2Yb(3,8\text{-}Me_2phen)$ (3).

Dissolution of the crystals of the complex **3** in toluene and THF is kinetically slow, in agreement with strong packing forces in the solid state, but gently warming (60°C) the solution over a period of one or two days gives saturated solutions that allowed ^{1}H NMR spectroscopic measurements at variable temperatures. At room temperature, both toluene-d_8 and THF-d_8 solution of **3** are deep red and show one major set of 5 resonances in a 2:2:2:30:6 ratio. This is in agreement with the presence of monomeric $Cp^*_2Yb(3,8\text{-}Me_2phen)$ with C_{2v} symmetry in which the phenanthroline ligand is symmetrically disposed relative to the Cp^*_2Yb fragment and these resonances are designated by the letter S for "symmetric". Small resonances are also present, contributing less than 5% of the peak intensity. When the toluene-d_8 and THF-d_8 solution are cooled, these low-intensity resonances grow at the expense of the resonances assigned to the monomeric $Cp^*_2Yb(3,8\text{-}Me_2phen)$ complex (S). The solutions change color from deep red at room temperature to purple at 250 K and blue at 200 K. The ^{1}H NMR spectra at low temperature in both solvents show three different sets of resonances; one set of 5 resonances attributed to the S isomer, the monomeric form of **3**, and in two other sets of resonances, labeled A_1 and A_2 (A for asymmetric), in which the methyl resonances are not equivalent, in agreement with the formation of a dimer. Ten resonances are

Eq. 2

expected for each isomer in a ratio 2:2:2:2:2:2:6:6:30, although some resonances were not located in a -100 to 100 ppm window. The ratio of the two asymmetric isomers A_1 and A_2 is approximately 60:40 in toluene and 55:45 in THF and the ratio is only slightly dependent on temperature, given the errors of the integration.

Two pairs of A resonances are attributed to the methyl groups based on the integration ratio and are highlighted by the red dots in Figure 10. These resonances are integrated and related to the S-methyl resonance that is highlighted by a blue dot in Figure 10. The relative change in population of these methyl group resonances is used to obtain the van't Hoff plot in Figure 11; the details are provided in the experimental section. The thermodynamic parameters for the reaction shown in Eq. 2, where M is the symmetric (S), monomer and D the asymmetric ($A_1 + A_2$), dimer, set of resonances, are determined from this plot.

$$2M \underset{}{\overset{K_1}{\rightleftharpoons}} D$$

The resulting ΔH values are -5.8 kcal/mol and -8.1 kcal/mol in THF and toluene, respectively, and the ΔS values are -26 kcal/mol/K and -31 kcal/mol/K in THF and toluene, respectively. At 298 K, the value of the dimerization constant (K_1) is 0.05 ± 0.005 M^{-1} in THF and 0.48 ± 0.01 M^{-1} in toluene. Similar pattern of ΔH and ΔS values are reported by Kochi and co-workers for the π-radical tricyclic phenalenyl.[34]



Figure 10. Stacked plot of ^{1}H NMR spectra in function of the temperature in THF. Red dots are the resonances used for integration of the asymmetric species and the blue dot is the resonance used for integration of the symmetrical species.



Figure 11. van't Hoff plot of the reaction 2M = D in toluene (M is **3** as a monomer, D corresponds to the the two dimeric isomers). The plot of Rln(K) (K_1 = D/M^2) *vs.* 1/T yields ΔH^0= -8.1(2) kcal/mol, ΔS^0 = -31(1) cal/mol/K and K_1 (25°C) = 0.48 M^{-1}.

Calculations.

The CASSCF methodology used in previous papers for the bipyridine adducts of Cp*$_2$Yb is extended to the monomeric phenanthroline adducts, Cp*$_2$Yb(phen) (**1**), Cp*$_2$Yb(3,8-

Me_2phen) (**3**) and Cp^*_2Yb(5,6-Me_2phen) (**7**). The ground state of Cp^*_2Yb(phen) is calculated to be comprised of two nearly degenerate triplet states, T_1 and T_2, which are 2.12 eV lower in energy than an open-shell singlet state. The state configuration for the f-orbitals are therefore pure (100 %) f^{13} and the T_1 and T_2 configuration for the π*-orbitals are 0.72 π^*_1 + 0.28 π^*_2 and 0.28 π^*_1+ 0.72 π^*_2, respectively. The calculated charge-transfer ground state is in accord with the observation of two LMCT bands near 500 cm^{-1} in the Vis-NIR spectrum in toluene solution.[6] The calculated ground states for the 3,8-Me_2phen and 5,6-Me_2phen adducts are similar to each other but somewhat different than that of the unsubstituted phenanthroline adduct. Thus, the calculated ground states are spin triplets (pure f^{13}) but the open-shell singlet states are only 0.08 eV and 0.09 eV higher in energy, respectively. The excited-state open-shell singlets are multiconfigurational in which the dominant configuration is f^{13}; in 3,8-Me_2phen, the f^{13}:f^{14} contributions are 0.75:0.25 and the π^*_1 is the only configuration that contributes. In the 5,6-Me_2phen adduct, the f^{13}:f^{14} contributions in the excited open-shell singlet state are 0.85:0.15 and the π^*_1 and π^*_2 contributions are 0.95 and 0.05, respectively.
The calculated spin triplet ground states in these three phenanthroline adducts are in dramatic contrast with the open-shell singlet ground states obtained in all the bipyridine adducts of Cp^*_2Yb. The calculated singlet-triplet separation is 0.28 eV in Cp^*_2Yb(bipy), singlet lowest and 2.12 eV in Cp^*_2Yb(phen) with triplet lowest. Thus the triplet energies change by 2.4 eV, about 60 $kcal.mol^{-1}$, just by changing the ligands.

The dimerization reaction is also studied by DFT calculations. The Cp* rings are replaced by Cp in these calculations since the full system for the dimer is prohibitively large. A transition state is calculated to be 15.4 $kcal.mol^{-1}$ (in Gibbs energy) above the monomers and the dimerization reaction is exoergic by 3.1 $kcal.mol^{-1}$ (Figure 12), consistent with the experiental observation that Cp^*_2Yb(phen) is a dimer. The calculated distance of the C-C bond formed in 4,4'-positions (1.596 Å) is in resonable agreement with the elongated C-C

bond found in the solid state structure (1.619 Å). The transition state involves two molecules of $Cp_2Yb(phen)$ with similar bond distances as calculated for the dimer, but with a C-C bond distance of 1.800 Å, the structure of the transition state is shown in Figure S26.



Figure 12. Reaction coordinate diagram for dimerization of $Cp_2Yb(phen)$.

Discussion.

Although the molecular geometry of monomeric $Cp^*_2Yb(phen)$ is similar to that found in the wide range of bipyridine adducts, their electronic structures are different.[4,5,27] The ground state electronic structure of the bipyridine adducts are open-shell singlets that are multiconfigurational in which the ground state wave function, Ψ, is $C_1|f^{13}, bipy^{\cdot-}> + C_2|f^{14}, bipy>$, where C_1 and C_2 are coefficients of the two configurations. For $Cp^*_2Yb(bipy)$, $C_1^2 = 0.83$.[4] This results in the ytterbium atom being intermediate valent, that is, it is neither Yb(III), f^{13}, nor Yb(II), f^{14} but in between these extreme values in which the f13 configuration is dominant. The open-shell singlet ground state (or states) determines the magnetic properties of these adducts, and in the case of the 4,5-diazafluorene adduct, is postulated to be the origin of the chemical reactivity.[35]

In contrast, the monomeric phenanthroline adducts $Cp^*_2Yb(phen)$ and $Cp^*_2Yb(5,6\text{-}Me_2phen)$ have open-shell triplet ground states, and the valence of ytterbium is fully trivalent. The

CASSCF computational studies indicate that two open-shell triplets are nearly degenerate and are some 2 eV lower in energy that the open-shell singlet state, consistent with the magnetic studies and the L-$_{III}$ edge XANES . A model that accounts for the different electronic ground states in the bipy and phen adducts is outlined next; the model is offered as a qualitative guide for what is known and as a guide for future experimental studies.

Whether a monomeric ytterbocene diimine complex has a triplet or singlet ground state is largely governed by kinetic exchange, that is, by mixing of excited state configurations into the ground state.[36,37] The interaction between the half-occupied ligand orbital and a half-occupied 4f-orbital stabilizes the singlet state while interactions between the half-occupied ligand orbital and the empty metal based orbitals on the Cp^*_2Yb fragment, especially the 5d-orbitals, stabilize the triplet state. To a first approximation, the strength of the interactions between the half-occupied ligand orbital and the metal orbitals are proportional to the square of the overlap and inversely proportional to the difference in energy between the ligand and metal orbitals. Since the overlap between the ligand orbitals and the Yb 5d orbitals is anticipated to be significantly larger than the overlap with the Yb 4f orbitals, whether the ground state is a singlet or triplet depends in large part on the energies of the ligand orbitals.[38] If the half-occupied ligand orbital is close in energy to the 4f-orbitals and has the proper symmetry to overlap with the lone half-filled Yb 4f-orbital, the singlet state is likely to be lowest in energy. However, if the half-occupied ligand orbital is not close in energy to the 4f orbitals or does not have the proper symmetry to overlap with the half-occupied Yb 4f orbital, the interaction between the half-occupied ligand orbital and the Yb 5d orbitals will be stronger, and the triplet state will be stabilized. If the ligand has empty orbitals close in energy to the half-occupied orbital, the ground state could be either a singlet or triplet depending on whether the interaction between the ligand orbitals and the singly occupied 4f-orbital is greater or weaker than the interactions with the empty 5d orbitals.

In $Cp^*_2Yb(bipy)$, the open-shell singlet is calculated to be 0.28 eV below the triplet and the experimental value of the singlet-triplet energy difference is about 0.1 eV by comparing to the Hubbard model.[11] When an f-electron is transferred to the LUMO, only one of the four possible π*-orbitals, the b_1-orbital, is of sufficiently low energy to be populated and the unpaired spin density is distributed among the pπ-orbitals on the C and N atoms of the bipyridine ligand. In this case, the half-occupied ligand orbital and the half-occupied Yb 4f orbital are close in energy and have the same symmetry, b_1, so the kinetic exchange configuration interaction stabilizes the open-shell singlet. This model fits all of the experimental and computational studies associated with the bipy adducts.[4,5]

In contrast, the LUMO and LUMO+1 of phenanthroline are close in energy (Figure 9, Chart 1) so that when electron transfer occurs, the electron occupies either the π^*_1 and/or π^*_2 orbitals, or both, which have b_1 and a_2 symmetry, respectively, (in C_{2v} symmetry). The ordering of these orbitals can be inverted by methyl group substituents in the solvent separated radical-anions as shown by EPR studies. Thus $phen^{\cdot-}$,[39] 2,9-$Me_2phen^{\cdot-}$,[40] 4,7-$Me_2phen^{\cdot-}$[39] and 5,6-$Me_2phen^{\cdot-}$[40] have $^2B_1$ ground states but 3,4,7,8-$Me_4phen^{\cdot-}$ has a $^2A_2$ ground state.[41] As in $Cp^*_2Yb(bipy)$, the b_1 orbital will be stabilized by interaction with the half-filled Yb 4f-orbital, which stabilizes the singlet state. This assumes that the orbital from which the electron on the close-shell Cp^*_2Yb metallocene is removed does not undergo reorganization, that is, the hole remains in a b_1 symmetry orbital. The a_2 orbital will not be stabilized by Yb 4f-orbitals since the single half-occupied orbital has b_1 symmetry. The ligand a_2 orbital can be stabilized by interaction with the empty Yb 5d orbitals. If the a_2 orbital is half-occupied, the triplet ground state will be stabilized, which is the case for all of the monomeric ytterbocene phenanthroline complexes reported here.



Chart 1.

The difference between the bipyridine and phenanthroline adducts can be illustrated using a MO diagram as illustrated in Figure 13. While the MO model does not capture the stabilization of the triplet or singlet state due to configuration interaction, the stable spin state is indicated by the relative spins of the electrons as indicated by the arrows in Figure 13. Figure 13 illustrates two extreme cases that are possible for two spins. This diagram may be extended to the specific examples of Cp^*_2Yb(diimine) since the symmetry orbitals of bent sandwich metallocenes are well known.[42,43] The d-orbitals that are empty, once the diimine σ-bonds are created, are the non-bonding $a_1(d_{x2-y2})$ metal-based orbital and the higher lying Cp*-Yb antibonding $d_{xy,yz}$ ortbitals of b_1 and a_2 symmetry. The seven f-orbitals occupied by 13 electrons are considered to be non-bonding and much lower in energy than the d-parentage orbitals. In the bipy adducts, Figure 13, left-hand side, the hole in the f-orbitals has b_1-symmetry as does the electron in the ligand-based orbital. As these two electrons have the same symmetry, they can mix to give a singlet state, following the Pauli principle. This model accounts for the electronic ground states of Cp^*_2Yb(bipy).[4,5]

Extending this MO model to the phenanthroline adducts is complicated by the fact that either the b_1 or a_2 orbitals or both are populated depending upon their relative energies, Figure 9. Thus, three idealized cases may be considered, (i) b_1 lies lower than a_2, (ii) a_2 lies lower than b_1, and (iii) b_1 and a_2 are of similar energy. Case (i) results in an orbital pattern found in

bipy, Figure 13, left-hand side. Case (ii) results in a similar orbital pattern, except that the b_1 orbital is replaced by the a_2 orbital. Both of these cases can result in singlet ground states. Case (iii), Figure 13, left-hand side, is applicable to the phen adducts described above. Thus, an electron in the a_2-ligand-based orbital is stabilized by interaction with a d_π orbital of a_2 symmetry on the Cp^*_2Yb metallocene resulting in the a_2-MO below the b_1-MO, resulting in a spin-triplet ground state, since the hole in the f-manifold is in a b_1 symmetry orbital. Case (iii) is more versatile since the methyl substitutents change the relative energies of the b_1 and a_2-orbitals generating the expectation that the magnetic properties can be controlled by the b_1-a_2 separation.



Figure 13. Qualitative MO diagram comparing bonding in $Cp^*_2Yb(bipy)$ and $Cp^*_2Yb(phen)$ monomer. Only the unpaired 4f electron is illustrated; the orbitals below those with arrows are filled. The direction of the arrows indicate the ground state: $Cp^*_2Yb(bipy)$ has a singlet ground state while $Cp^*_2Yb(phen)$ has a triplet ground state.

Inspection of the b_1-orbital in $phen^{\bullet-}$ shows that the spin density is more likely to reside on N, C(2,9), C(3,8), and C(4,7) whereas in the a_2-orbital the spin density is likely to be found on C(2,9), C(4,7) and C(5,6), chart 1. Thus, the unpaired spin density on N is greater in the b_1-orbital than in the a_2-orbital. Population of the a_2-orbital increases spin density on $p\pi$-

orbitals of C(4,7), which is the site of C-C bond formation in the dimer. Thus, substituents on the phenanthroline ring in the ytterbocene adducts modulates the unpaired spin density of the pπ-orbitals and the radical character at a given site and therefore the site at which chemical reactions occurs.

Dimerization of two σ-carbon radicals forming a σ-C_2 single bond involves two enthalpy changes with opposite signs. The exothermic term involves C-C bond formation, about 80-85 kcal.mol^{-1}, and the endothermic term is due to loss of resonance stabilization in the σ-radical, estimated to be about 35 kcal.mol^{-1} per radical.[34] The net enthalpy changes favor dimerization but loss of entropy results in ΔG being close to zero.

Conclusion.

The key conclusion in this article is that the electronic ground states for Cp*$_2$Yb(phen), the 3,8-, and the 5,6-dimethyl adducts are different from those of Cp*$_2$Yb(bipy) and its methyl and dimethyl-substituted adducts; the ground state of monomeric Cp*$_2$Yb(phen) and its adducts is a spin triplet and the valence of ytterbium is trivalent. In contrast, the ground state of Cp*$_2$Yb(bipy) is a multiconfigurational open-shell singlet and ytterbium is intermediate valent. The chemical manifestation of this difference is that the solid state structure of the sublimed phenanthroline adduct is a monomer but that of the crystallized adduct is a dimer in which the phenanthroline ligands are coupled by formation of a C-C bond between the carbons at the 4-position. The 3,8-Me$_2$phenanthroline adduct is similar but in this case the monomer and dimer are in equilibrium in toluene-d$_8$ or in tetrahydrofuran-d$_8$ with values of ΔH of -8.1 kcal/mol and -5.8 kcal/mol, respectively, and ΔS values of -31 kcal/mol/K and -26 kcal/mol/K, respectively. The origin of the different physical and chemical properties is postulated to arise from the different symmetry orbitals available in the phen$^{\bullet-}$ and bipy$^{\bullet-}$; the former has two accessible orbitals of b_1 and a_2 symmetry (in C_{2v}) while the later has only one of b_1 symmetry.

Experimental.

General considerations. All reactions were performed using standard Schlenk-line techniques or in a drybox (MBraun). All glassware was dried at 150 °C for at least 12 h prior to use. Toluene, pentane and diethyl ether were dried over sodium and distilled while CH_2Cl_2 was purified by passage through a column of activated alumina. Toluene-d_8 was dried over sodium and CH_2Cl_2-d_2 was dried over calcium hydride. All the solvents were degassed prior to use. ^{1}H NMR spectra were recorded on Bruker AVB-400 MHz, DRX-500 MHz, AVB-600 MHz and Avance 300 MHz spectrometers. ^{1}H chemical shifts are in δ units relative to TMS, and coupling constants (*J*) are given in Hz. Infrared spectra were recorded as Nujol mulls between KBr plates on a Thermo Scientific Nicolet IS10 spectrometer. Samples for UV-Vis-NIR spectroscopy were contained in a Schlenk-adapted quartz cuvette and obtained on a Varian Cary 50 scanning spectrometer. Melting points were determined in sealed capillaries prepared under nitrogen and are uncorrected. Elemental analyses were determined at the Microanalytical Laboratory of the College of Chemistry, University of California, Berkeley. X-ray structural determinations were performed at CHEXRAY, University of California, Berkeley. Magnetic susceptibility measurements were made for all samples at 1, 5 and 40 kOe in a 7 T Quantum Design Magnetic Properties Measurement System that utilizes a superconducting quantum interference device (SQUID). Sample containment and other experimental details have been described previously.[44] It is important to note that the susceptibility values obtained when the samples were contained in Kef-F containers and quartz tube are identical within experimental errors.[27] Diamagnetic corrections were made using Pascal's constants. The samples were prepared for X-ray absorption experiments as described previously, and the same methods were used to protect these air-sensitive compounds from oxygen and water contamination.[5] The samples were loaded into a LHe-flow cryostat, and X-ray absorption measurements performed at the Stanford Synchrotron

Radiation Lightsource on beamline 11-2. Data were collected at temperatures ranging from 30 to 300 K, using a Si(220) double-crystal monochromator. Fit methods were the same as described previously.[5] Reported spectra were energy calibrated by setting the first inflection point of the absorption spectrum on a Yb_2O_3 reference sample to 8943 eV. Low temperature (ca. 2 K) EPR spectra were obtained with a Varian E-12 spectrometer equipped with an EIP-547 microwave frequency counter and a Varian E-500 gaussmeter, which was calibrated using 2,2-diphenyl- 1-picrylhydrazyl (DPPH, *g*= 2.0036).

Calculations. The ytterbium atom was treated with a small-core relativistic pseudopotential (RECP) ([Ar] + 3d)[45] in combination with its adapted basis set (segmented basis set that includes up to g functions). The carbon, nitrogen, oxygen, and hydrogen atoms were treated with an all-electron double-ζ, 6-31G(d,p),[46] All the calculations were carried out with the Gaussian 03 suite of programs[47] ORCA suite of program[48] either at the Density Functional Theory (DFT) level using the B3PW91[49] hybrid functional or at the CASSCF level; only one active space and inactive orbitals were used in the calculation. The geometry optimizations were performed without any symmetry constraints at either the DFT or the CASSCF level. The electrons were distributed over four 4f orbitals and the two π* orbitals of phenanthroline.

Syntheses. The ligands, 1-10-phenanthroline (phen) and 4-methylphenanthroline (4-Mephen) were purchased from Aldrich while 5-methylphenanthroline (5-Mephen) was obtained from Tokyo Kasei Kokyo Co. All ligands were purified by sublimation between 80 and 200 °C/10^{-2} mm prior to use. The ligands 3-methyl-1,10-phenanthroline (3-Mephen) and 3,8-dimethyl-1,10-phenanthroline (3,8-Me_2phen) were synthesized according to a published procedure[50] and sublimed at 140 ºC/10^{-2} mmHg prior to use. ^{1}H NMR (3-Mephen): (CD_3Cl, 295K, δ (ppm)) 9.18 (d, J = 7.8Hz, 1H), 9.03 (s, 1H), 8.23 (d, J = 7.8Hz, 1H), 8.02 (s, 1H), 7.74 (dd, J = 7.6, 3.2Hz, 1H), 7.74 (d, J = 7.2Hz, 1H), 7.61 (dd, J = 7.3, 3.2Hz, 1H), 2.61 (s, 3H). ^{1}H

NMR (3,8-Me_2phen): (CD_3Cl, 296K, δ (ppm)) 9.05 (s, 2H), 8.64 (s, 2H), 7.74 (s, 2H), 2.64 (s, 6H).

Cp^*_2Yb(phen) (1). The complex $Cp^*_2Yb(OEt_2)$ (0.217 g, 0.420 mmol) was combined with 1,10-phenanthroline (0.095g, 0.420 mmol) and toluene (30 mL) was added at room temperature. The resulting purple/blue solution was stirred for 2h at room temperature as a dark powder formed. The suspension was cooled at -20°C and the dark colored powder was collected by filtration (125 mg, 83%). The dark powder was washed with toluene (3x5mL) and was heated in toluene (20 mL, 80°C), filtered while hot and slowly cooled at room temperature and then at -20°C. The dark microcrystalline purple powder was collected by filtration and dried under reduced pressure (70 mg, 47%). An alternative method was used in order to obtain crystals suitable for X-ray diffraction data collection by crystallization from toluene. A toluene solution of 1,10-phenanthroline (0.033 g, 0.186 mmol) was carefully layered at the top of a toluene solution of $Cp^*_2Yb(OEt_2)$ (0.096 g, 0.186 mmol). Slow diffusion of the two solutions overnight (16h) resulted in formation of X-Ray quality crystals at the interface and along the walls of the Schlenk flask that were collected by filtration and dried under reduced pressure (75 mg, 68%). M.p. 295-297°C (lit 297-300°C).[27] ^{1}H NMR: (toluene-d_8, 299 K, δ (ppm)) 139.94 (2H, phen), 47.87 (2H, phen), 14.02 (2H, phen), 4.14 (30H, Cp^*), 0.47 (2H, phen). Crystals of Cp^*_2Yb(phen) are sparingly soluble in C_7D_8 or THF-d_8. The ^{1}H NMR spectrum was obtained from a warmed concentrated C_7D_8 solution measured at 299 K. Anal. Calcd for $C_{32}H_{38}N_2Yb$: C, 61.62; H, 6.14; N, 4.49. Found: C, 61.99; H, 6.04; N, 4.45. IR (cm^{-1}): 1610 (m), 1590 (w), 1550 (w), 1498 (m), 1445 (s), 1359 (s), 1308 (s), 1290 (m), 1224 (w), 1172 (w), 1117 (m), 1054 (m), 1022 (w), 859 (w), 823 (m), 798 (m), 734 (m), 689 (m). Crashed crystals of Cp^*_2Yb(phen) sublimed in a 180 °C – 190 °C temperature range in an ampoule sealed under vacuum afforded crystals of Cp^*_2Yb(phen) (34 mg) over a one month period of time.

$[Cp^*_2Yb(phen)]^+I^-$ (2).[27] The complex $Cp^*_2Yb(OEt_2)$ (0.172g, 0.333 mmol) was combined with 1,10-phenanthroline (0.060g, 0.333 mmol) and AgI (0.078g, 0.333 mmol). Toluene (40 mL) was added at room temperature and the purple solution was stirred for 16h at room temperature. The supernatant liquid was removed and the brown residue was extract with CH_2Cl_2. The solution was red and a grey residue remained. The solution was filtered, concentrated to 5 mL and cooled to -20°C. Large red crystals formed (120mg, 48%). mp: 175-180°C. ^{1}H NMR: (CD_2Cl_2, 300K, δ (ppm)) 280.87 (2H, phen) 52.43 (2H, phen), 9.52 (2H, phen), 5.31 (2H, CH_2Cl_2), 3.82 (30H, Me_5C_5), -2.48 (2H, phen). N.B. In previous paper,[27] δ at 280 ppm was not observed and $CHDCl_2$ was assigned as a phen resonance. Anal. Calcd for $C_{32}H_{38}N_2YbI_2$•$1.5CH_2Cl_2$: C, 45.83; H, 4.71; N, 3.19. Found: C, 45.98; H, 4.50; N, 3.38. IR (cm^{-1}): 1622 (w), 1518 (w), 1460 (s), 1415 (m), 1377 (s), 1273 (w), 855 (m), 728 (s).

$Cp^*_2Yb(3,8\text{-}Me_2phen)\cdot(C_7H_8)$ (3). The complex $Cp^*_2Yb(OEt_2)$ (0.100 g, 0.192 mmol) was combined with 3,8-dimethyl-1,10-phenanthroline (3,8-Me_2phen, 0.040 g, 0.192 mmol) and toluene (10mL) was added at room temperature. The deep purple solution was stirred for 2h at room temperature, concentrated to ca. 5mL, warmed to dissolve the dark residue and the resulting dark solution was filtered while warm. The filtrate was slowly cooled at -20°C. Dark purple-red crystals suitable for X-ray crystallography formed overnight. A second crop was obtained. (Combined yield, 82 mg, 66%). mp: 286-288°C. Anal. Calcd for $C_{34}H_{42}N_2Yb\cdot C_7H_8$: C, 66.20; H, 6.77; N, 3.77. Found: C, 65.76; H, 6.65; N, 3.38. IR (cm^{-1}): 1625 (w), 1573 (w), 1461 (s), 1410 (w), 1377 (s), 1261 (s), 1214 (w), 1153 (m), 1079 (s), 1020 (s), 861 (w), 799 (s), 728 (m), 692 (m).

$Cp^*_2Yb(5\text{-}Mephen)$ (4). The complex $Cp^*_2Yb(OEt_2)$ (0.160 g, 0.309 mmol) was combined with 5-methyl-1,10-phenanthroline (5-Mephen, 0.060g, 0.309 mmol). Toluene (10 mL) was added at room temperature and the purple solution was stirred for 2h at room temperature. The volume of solvent was concentrated to 5 mL, then cooled at -20°C. A dark powder

formed overnight which was crystallized from warm toluene (152 mg, 77%). NMR: (toluene-d_8, 300K, δ (ppm) 138.72 (1H, phen), 138.59 (1H, phen), 47.92 (1H, phen), 39.33 (1H, phen), 14.18 (1H, Phen), 11.40 (1H, phen), 4.09 (30H, C_5Me_5), 0.06 (1H, phen), -0.58 (3H, Me-phen). mp: 280-283°C. Anal. Calcd for $C_{33}H_{40}N_2Yb$: C, 62.15; H, 6.32; N, 4.39. Found: C, 61.74; H, 6.02; N, 4.32. IR (cm^{-1}): 1626 (m), 1605 (w), 1578 (w), 1550 (w), 1504 (m), 1444 (s), 1377 (vw), 1355 (s), 1322 (s), 1281 (m), 1221 (vw), 1161 (m), 1136 (m), 1085 (vw), 1054 (m), 994 (w), 878 (m), 807 (w), 787 (w), 773 (m), 709 (w), 696 (m).

Cp^*_2Yb(4-Mephen) (5). The complex $Cp^*_2Yb(OEt_2)$ (0.304 g, 0.588 mmol) was dissolved in diethyl ether and added dropwise over 30 min to a cold diethyl ether suspension (10 mL, -77°C) of 4-methyl-1,10-phenanthroline (4-Mephen, 0.114 g, 0.588 mmol). While adding the phenanthroline, the suspension progressively turned to deep blue. When the addition was complete, the suspension was stirred at -77°C for 2 h and filtered to afford a dark blue-green powder (210 mg, 56%) which was washed with cold diethyl ether (2x10 mL, -77°C) and dried under reduce pressure. ^1H NMR: (C_6D_6, 295K, δ (ppm), 4.03 (Cp^*), only one discernable peak. mp: 254-256°C. Anal. Calcd for $C_{33}H_{40}N_2Yb$: C, 62.15; H, 6.32; N, 4.39. Found: C, 62.06; H, 6.43; N, 4.55. IR (cm^{-1}): 3069 (w), 3021 (w), 2954 (s), 2724 (w), 1634 (m), 1618 (m), 1512 (m), 1445 (s), 1401 (w), 1376 (w), 1354 (m), 1321 (w), 1301 (s), 1261 (w), 1190 (w), 1157 (w), 1086 (m), 1062 (w), 1048 (w), 1022 (w), 898 (s), 858 (m), 824 (m), 800 (s), 779 (w), 767 (w), 737 (w), 691 (w), 665 (w).

Cp^*_2Yb(3-Mephen)·0.5(C_7H_8) (6). The complex $Cp^*_2Yb(OEt_2)$ (0.105 g, 0.203 mmol) was combined with 3-methyl-1,10-phenanthroline (3-Mephen, 0.040 g, 0.203 mmol) and toluene (10mL) was added at room temperature. The deep purple solution was stirred for 2h at room temperature and a dark precipitate formed. The suspension was warmed to dissolve the dark powder and the resulting solution was filtered while warm. The filtrate was slowly cooled to -20°C to yield a dark microcrystalline powder. Two crop were obtained (Combined yield, 85

mg, 65%). ^{1}H NMR: (toluene-d_8, 295K, δ (ppm) 121.47 (1H, phen), 118.38 (1H, phen), 59.15 (1H, phen), 57.17 (1H, phen), 55.02 (1H, Phen), 52.07 (1H, phen), 18.69 (1H, phen), 3.79 (30H, C_5Me_5), -9.51 (3H, Me-phen). mp: 270-272°C. Anal. Calcd for $C_{33}H_{40}N_2Yb\cdot0.5(C_7H_8)$: C, 64.11; H, 6.49; N, 4.10. Found: C, 64.40; H, 6.49; N, 3.96. ^{1}H NMR spectrum confirmed the presence of the toluene. MS: {Cp^*_2Yb(3-Mephen)}, m/z = 638. IR (cm^{-1}): 1612 (m), 1554 (w), 1494 (w), 1454 (s, nujol), 1377 (s), 1364 (s), 1320 (s), 1297 (s), 1229 (m), 1174 (m), 1118 (m), 1065 (m), 1022 (w), 886 (m), 880 (m), 776 (m), 731 (s), 696 (m), 675 (m).

Cp^*_2Yb(5,6-Me_2phen) (7). The complex $Cp^*_2Yb(OEt_2)$ (0.208 g, 0.403 mmol) was combined with 5,6-dimethyl-1,10-phenanthroline (5,6-Me_2phen, 0.0838 g, 0.403 mmol) and toluene (20 mL) was added at room temperature. The deep purple solution was stirred for 16 h at room temperature, concentrated to ca. 5 mL, warmed to dissolve the dark residue and filtered while hot. The filtrate was slowly cooled at -20 °C. A dark purple microcrystalline powder formed (204 mg, 78%) which was crystallized in warm cyclohexane yielding block-like purple X-ray suitable crystals (125mg, 48%). ^{1}H NMR: (toluene-d_8, 300K) δ (ppm) 137.44 (2H, phen), 44.10 (2H, phen), 14.66 (2H), 3.95 (30H, C_5Me_5), 0.03 (6H, Me-phen). mp: 285-287°C. Anal. Calcd for Anal. Calcd. for $C_{34}H_{42}N_2Yb$: C, 62.66; H, 6.50; N, 4.30. Found: C, 62.74; H, 6.43; N, 4.37. IR (cm^{-1}): 1605 (m), 1584 (w), 1480 (w), 1426 (s), 1375 (m), 1345 (w), 1305 (w), 1275 (vw), 1218 (vw), 1190 (w), 1167 (w), 1145 (w), 1073 (w), 1019 (w), 943 (w), 804 (s), 758 (w), 736 (s), 686 (m). The crystal data, Table 10, for **7, crystallized** were obtained on crystals obtained by crystallization from cyclohexane. The crystal data, Table 10, for **7, sublimed**, were obtained on crystals that were crystallized from cyclohexane then sublimed in an ampoule sealed under vaccum at 195 °C over a period of two months. The sublimate contained needles and block-like crystals that were separated manually. The needles had the same unit cell parameters as those obtained for **7, crystallized**. The block-like crystals

crystallized in the same crystal system and space group but with different cell parameters and contained only one molecule in the unit cell, Table 10.

Variable temperature ^{1}H NMR spectra of 3.

Toluene-d$_8$. ^{1}H NMR: (toluene-d$_8$, 300K) A major species, labeled S, was observed at δ (ppm) 95.54 (2H, phen), 51.07 (2H), 3.83 (2H, phen), 3.63 (30H, C_5Me_5), -10.02 (6H, Me-phen) and two minor species (labeled A_1 and A_2 accounting for less than 5% of the total) were observed. When the NMR tube was cooled, the two minor species observed at room temperature increased in intensity that represent two unsymmetrical (the position 2 and 9, 3 and 8, 4 and 7 and 5 and 6 are no more equivalent) complexes in agreement with the formation of two isomeric dimers. The three different species are labeled S, for the symmetrical monomer, A_1 and A_2 for the two asymmetric isomeric dimers. In toluene, one proton could not be observed for A_1 and A_2, presumably because it was under the toluene resonances. The amount of A_1 and A_2 is 40 % / 60 % at 210 K and this ratio is only slightly temperature dependent. ^{1}H NMR: (toluene-d$_8$, 210K) δ (ppm) 105.21 (0.13H, phen-S), **89.95** (0.6H, Me-A_1), **85.51** (1H, Me-A_2), 74.24 (0.13H, phen-S), 56.20 (0.2H, phen-A_1), 45.99 (0.33H, phen-A_2), **22.66** (1H, Me-A_2), 14.06 (0.50H, br, phen-A_1+phenA_2), 4.41 (0.13H, phen-S), 3.63 (2H, Cp*-S), -0.85 (16.5H, br, $\nu_{1/2}$=1100Hz, Cp*-A_1+A_2), **-13.44** (0.6H, Me-A_1), -15.42 (0.33H, phen-A_2), -21.20 (0.2H, phen-A_1), **-27.10** (0.4H, Me-S), -43.25 (0.2H, phen-A_1), -70.24 (0.33H, phen-A_2), -111.47 (0.33H, phen-A_2), -113.58 (0.2H, phen-A_1).

THF-d$_8$. ^{1}H NMR: (thf-d$_8$, 300K) δ (ppm) 72.72 (2H, phen), 42.33 (2H), 5.21 (2H, phen), 2.79 (30H, C_5Me_5), -7.01 (6H, Me-phen) and two minor species (less than 5% total). In THF, two protons could not be detected for each dimer (A_1 and A_2). The amount of the isomers A_1 and A_2 is 55%-45% at 198K and is only slightly temperature dependant. ^{1}H NMR: (thf-d$_8$, 198K) δ (ppm) **96.76** (1H, Me-A_1), **92.01** (0.8H, Me-A_2), 79.94 (0.27H, phen-S), 60.91 (0.22H, phen-A_2), 60.04 (0.27H, phen-S), 49.99 (0.25H, phen-A_1), **23.55** (0.8H, Me-A_2),

13.03 (0.5H, br, phen-A_1+phenA_2), 5.70 (0.27H, phen-S), 2.91 (4.1H, Cp^*-S), -1.52 (18H, br, $\nu_{1/2}$=1200 Hz, Cp*-A_1+A_2), **-15.09** (1H, Me-A_1), -16.74 (0.25H, phen-A_1), **-19.44** (0.8H, Me-S), -24.38 (0.22H, phen-A_2), -42.09 (0.25H, phen-A_1), -75.78 (0.22H, phen-A_2).

Resonances in bold were the resonances used for the integration and the calculation of equilibrium constants. They were used because they are singlets whose resonances are clearly visible over the temperature range of the study (197.5 – 315 K). These calculations assume that the reaction shown in eq. 2, where M is the symmetric set, S, and D the asymmetric sets of resonances, A_1 and A_2.