

CONF-8006217--1



# Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

## EARTH SCIENCES DIVISION

**MASTER**

Presented at the AAPG-SEPM-EMD Annual Convention,  
Denver, CO, June 8-11, 1980


A STRUCTURAL AND SEDIMENTOLOGICAL STUDY OF THE  
CERRO PRIETO GEOTHERMAL FIELD, BAJA CALIFORNIA,  
MEXICO

Stephen Vonder Haar

LBL--12187

June 1981

DE82 007773



## **DISCLAIMER**

**This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.**

## **DISCLAIMER**

**Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.**

#### **LEGAL NOTICE**

This book was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

**A STRUCTURAL AND SEDIMENTOLOGICAL STUDY  
OF THE CERRO PRIETO GEOTHERMAL FIELD,  
BAJA CALIFORNIA, MEXICO**

**Stephen Vonder Haar**

**Earth Sciences Division  
Lawrence Berkeley Laboratory  
University of California  
Berkeley, California 94720**

**DISCLAIMER**

This book was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

**June 1981**

**This work was supported by the Assistant Secretary for Conservation and Renewable Energy, Office of Renewable Technology, Division of Geothermal and Hydropower Technologies of the U. S. Department of Energy under Contract No. W-7405-ENG-48.**

## INTRODUCTION

Since 1977 the Comisión Federal de Electricidad de México (1979) and the Lawrence Berkeley Laboratory (1978) have cooperatively studied the Cerro Prieto geothermal field, located approximately 35 km south of the Mexican-American border in the Mexicali-Salton Trough.

As part of these studies, geophysical and lithologic well logs from over fifty wells have been qualitatively and quantitatively analyzed using both manual and computer interpretation techniques. These logs were studied to make stratigraphic correlations throughout the Cerro Prieto field and to interpret the deltaic depositional environment of the field's lithologic units. Dipmeter and seismic data were of great value in making stratigraphic interpretations and extrapolations. Cross sections were constructed to illustrate lithofacies variations throughout the geothermal field. In turn, these sections were used to construct a three-dimensional model of the Cerro Prieto geothermal reservoir.

Petrographic microscopy, scanning electron microscopy, and x-ray diffraction analyses of well-bore cuttings and cores were utilized to determine the degree and distribution of hydrothermal alteration by fluids at temperatures up to 350°C, the origins of dissolution porosity, and the relative degree of fracture versus dissolution porosity. The results of these analyses were confirmed by log-derived determinations of formation fluid properties, porosity, and petrophysical properties and by studies of Cerro Prieto cores conducted under in-situ conditions. The results of this research were integrated into the Cerro Prieto reservoir model.

In order for the exploitation of a geothermal resource to be economical, the reservoir porosity and permeability must be adequate and the distribution of that porosity and permeability must be well known. This study of Cerro Prieto suggests that the roles of fracture-dominated porosity and authigenic mineral plugging in the production reservoirs may have been overstated in earlier conceptualizations of geothermal systems (see DiPippo, 1980, for a worldwide summary of systems). The role of dissolution porosity is very important. As documented by Schmidt and McDonald (1979) for petroleum reservoirs, diagenetic processes determine porosity types and their distributions throughout the life span of a field. Studies of these processes will contribute to the estimation of reservoir porosity distribution and will assist in development of a management plan to optimize the utilization of the geothermal resource.

#### DISCUSSION

This report is an updated version of the poster session and oral presentation made at the 1980 annual meeting of the AAPG-SEPM-EMD in Denver, Colorado (Noble and Vonder Haar, 1980).

Figure 1 shows the location of the Cerro Prieto geothermal field in relation to the numerous faults in the Salton Trough (see Crowell and Sylvester, 1979; Elders, 1979; Vonder Haar and Puente, 1979; and Vonder Haar and Howard, 1981, for details on faulting related to geothermal development).

Figure 2 shows the well locations and a simplified version of major faults and fault zones. There are numerous tectonic analogs to the Cerro Prieto field.

One of the most useful has been the ocean-floor transform fault "A" system on the Mid-Atlantic Ridge (Figure 3).

Within the Cerro Prieto field, coring (Figure 4) has been carried out to clarify the deltaic facies of sandstones, siltstones, shales, and occasional conglomerates. Recovery rates have been variable, from 14 to 96%, and driller's reports of loose sand at depths of 1200 m or greater were the first clue that extensive dissolution of cement has occurred. Scanning electron micrographs with 200 to 5000 times magnification, as shown in Figure 5, provided evidence of dissolution of both carbonate and silicate minerals, as well as precipitation of clay minerals in pore throats.

Figures 6 and 7, derived from wireline density logs, suggest that the porosity in sandstones at depths of 5000 to 6000 ft could reach 15 to 45% in some wells. In Well M-103 (Figure 7) these zones of high porosity and low density are well below the A/B contact, a gradational contact which represents the transition from unconsolidated sediments to the hydrothermally altered and densified ones. In Well M-93 (Figure 6) the divergence of shale and sandstone density near the 7500-ft depth again suggested dissolution porosity. This interpretation is based on analogous log data found in petroleum studies (see Schmidt and McDonald, 1979). Thin-section analysis of cores confirmed that dissolution porosity results not only from the removal of cement but also of feldspar and quartz (see the report by Lyons and van de Kamp, 1980, for their interpretation).

The surface depicted in Figure 8 represents the first occurrence of 5+ authigenic epidote in the sandstones of the Cerro Prieto field. This may

indicate a zone above which there is extensive dissolution porosity. Below this surface, it is suggested that there is less dissolution porosity and a relative increase in microfracture porosity.

#### ACKNOWLEDGEMENTS

This work was supported by the Assistant Secretary for Conservation and Renewable Energy, Office of Renewable Technology, Division of Geothermal and Hydropower Technologies of the U.S. Department of Energy under Contract No. W-7405-ENG-48. Special thanks are due to colleagues at Cerro Prieto and at Lawrence Berkeley Laboratory. Grateful appreciation is due Norman Goldstein, Jack Howard, Marcelo Lippmann, and John Noble for their encouragement.

#### REFERENCES

- Choukroune, P., Francheteau, J., and Le Pichon, X., 1978. In situ structural observations along transform fault A in the FAMOUS area, Mid-Atlantic Ridge. *Geol. Soc. America Bull.*, v. 89, p. 1013-1029.
- Comisión Federal de Electricidad, 1979. Proceedings, Second symposium on the Cerro Prieto geothermal field, October 17-19, 1979, Mexicali, Baja California, Mexico. Comisión Federal de Electricidad de México, 640 p.
- Crowell, J. C., and Sylvester, A. G., 1979. Tectonics of the juncture between the San Andreas fault system and the Salton Trough, Southeastern California. Guidebook for annual G.S.A. meeting, San Diego, Nov. 1979. Dept. Geol. Sci., University of California, Santa Barbara, California, 193 p.
- DiPippo, R., 1980. Geothermal energy as a source of electricity. U. S. Department of Energy Pub., U. S. Government Printing Office, Washington, D.C., Stock Number 061-000-00390-8, 370 p.
- Elders, W. A. (Ed.), 1979. Geology and geothermics of the Salton Trough. University of California, Riverside, Campus Museum Contr. No. 5; G.S.A. Annual Meeting Field Trip Guidebook No. 7, San Diego, California, Nov. 1979, 108 p.

Elders, W. A., Hoagland, J. R., Vallette, J. N., Williams, A., Barker, C., and Collier, P., 1981. A comprehensive study of samples from geothermal reservoirs: Petrology and geochemistry of subsurface and surface samples from the Cerro Prieto geothermal field, Baja California, Mexico. Preliminary report prepared by the Institute of Geophysics and Planetary Physics, University of California at Riverside, for LBL/DOE.

Lawrence Berkeley Laboratory, 1978. Proceedings of the first symposium on the Cerro Prieto geothermal field, Baja California, Mexico, September 20-22, 1978, San Diego, California. Lawrence Berkeley Laboratory Report LBL-7098, 445 p.


Lyons, D. J., and van de Kamp, P. C., 1980. Subsurface geological and geophysical study of the Cerro Prieto geothermal field, Baja California, Mexico. Lawrence Berkeley Laboratory Report LBL-10540, 95 p.

Noble, J. E., and Vonder Haar, S. P. 1980. A study of diagenetic processes in sandstone from the Cerro Prieto Geothermal field, Baja California, Mexico. Abstracts volume of AAPG-SEPM-EMD annual convention, Denver Colorado, June 8-11, p. 99.

Schmidt, V., and McDonald, D. A., 1979. Secondary reservoir porosity in the course of sandstone diagenesis. Continuing Education Course Notes, Amer. Assoc. Petroleum Geologists, Tulsa, Oklahoma, 125 p.

Vonder Haar, S., and Puente C., I., 1979. Fault intersection and hybrid transform faults in the southern Salton Trough geothermal area, Baja California, Mexico. Geothermal Resources Council Transactions, v. 3, p. 761-764.


Vonder Haar, S., and Howard, J. H., 1981. Intersecting faults and sandstone stratigraphy at the Cerro Prieto geothermal field. Geothermics, in press.



XBL 801-6718


Figure 1. Regional geology of the Cerro Prieto geothermal field. Note the prevalence of long NW-SE trending faults intersected by much shorter faults with NE-SW trends.





XBL 797-7583A

Figure 3. Interpretive block diagram of the oceanic transform fault "A" in the Mid-Atlantic Ridge showing structural domains. The ramps (R), cross-faults, and 200-m-wide active zone of strike-slip movement (AZ) within a 4-km-wide fault trough suggest the possible complexity of faulting along the Cerro Prieto and Imperial faults and within the producing geothermal fields (after Choukroune et al., 1978).




XBL813-2721

**Figure 4.** Composite core section from the earliest developed wells at Cerro Prieto field. Note the transition from unconsolidated sand and clay to hydrothermally densified sandstones and shales between 700 and 900 m depth. These cores helped confirm ideas on deltaic facies later greatly altered by fluids of 250 to 350°C. Numbers such as 0.90 next to cores indicate the length in meters of individual core segments.



XBB 797-9450A


Figure 5. Clay minerals (A) clogging a pore throat in the deltaic sandstone at 1215 m depth in well M-38. Note also the framework mineral overgrowth (B) in the lower center of the figure. Such pore throat reduction may be present even when porosity is 25%. Field of view across the scanning electron micrograph is 0.1 mm.



Mineral data after Elders and others (in press)\*

XBL 805-7074


Figure 6. Density versus depth plot for well M-93. Note the similarity of density in the sandstones and shales to approximately 5000 ft depth and the decrease in relative density of sandstone to shale below this depth. This is interpreted as an indication of moderate dissolution porosity.



\*Mineral data after Elders and others (in press)

XBL 805-7079

Figure 7. Plot of porosity and density versus depth for the very productive well M-103 as determined from well logs and cuttings. The high porosity in the perforated interval indicates extensive dissolution porosity. Note the marked density increase at 4000 ft depth at the A/B contact. This contact is the zone reported by CFE geologists as the transition from unconsolidated sands and clays to sandstone and shale.



XBL 804-7023

Figure 8. Three-dimensional computer plot of the first occurrence of 5% epidote in well cuttings. Quantitative x-ray diffraction data are from Elders et al. (1981). The 29 wells used are indicated by "x," with wells M-5 and M-6 shown for orientation. This 5% epidote surface may be a diagnostic horizon above which dissolution porosity may reach 35%, at least in patchy or lenticular zones. See Figure 7 for the 5% prehnite and 5% epidote zone near the base of the suggested region of extensive dissolution porosity and good permeability.