LBL-13291

Lawrence Berkeley Laboratory
UNIVERSITY OF CALIFORNIA

Physics, Computer Science & Mathematics Division

RECE
LAWR
OCT 9 1981

Presented at the European Physical Society International Conference on High Energy Physics, Lisbon, Portugal, July 9-15, 1981

MULTIPION CORRELATIONS IN e^+e^- ANNIHILATION AT SPEAR

Gerson Goldhaber

September 1981

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 6782.

Prepared for the U.S. Department of Energy under Contract W-7405-ENG-48

LBL-13291

Paper presented on results from the SLAC-LBL Mark II Collaboration at SPEAR; International Conference on High Energy Physics, Lisbon, Portugal, July 9-15, 1981.

LBL-13291

MULTIPION CORRELATIONS IN e^+e^- ANNIHILATION AT SPEAR†

Gerson Goldhaber
Lawrence Berkeley Laboratory and Department of Physics
University of California, Berkeley, California 94720

A comparison between like charge and unlike charge particle distributions allow the study of both Bose-Einstein effects and of inclusive resonance production. In this paper I will present some results on the first of these topics.‡

- The data presented here is based on 1.3×10^6 events at the ψ/J and a much smaller number obtained in the $E_{cm} = 4-7$ GeV region in the Mark II detector operating at SPEAR.
- The Bose-Einstein effects are obtained in terms of ratios R^{--}_{+-} ($\equiv R^L_U$) of distributions of the Lorentz invariant quantities[1] $Q^2(12) = M^2(12) - (m_1+m_2)^2$ for two particles and the corresponding three-particle quantity: $Q^2(123) = M^2(123) - (\sum_{i=1}^{3} m_i)^2$; viz.,

$$R^{--}_{+-} = \left(\frac{1}{\sigma_{--}}\frac{d\sigma^{--}}{dQ^2(12)}\right)\Big/\left(\frac{1}{\sigma_{+-}}\frac{d\sigma^{+-}}{dQ^2(12)}\right)$$

  with an analogous definition for R^{---}_{+--} for three-particle correlations. Here M(12), M(123) are effective masses for two and three particles respectively, m_i are rest masses of the particles.
- The Bose-Einstein enhancement in R can be expressed[1] as $1+\alpha e^{-\beta Q^2}$. Here α measures the degree of chaoticity[2,3] of the production process. For two pions $\alpha \to 1$ corresponds to maximal chaoticity, while β corresponds to an effective radius (squared) of the interaction volume from which the pions originate. Alternately the three-dimensional radius r_o and the time development $c\tau_o$ are sometimes treated independently.[4]

Results at the $\psi/J(3.1)$ Resonance

- R^{--}_{+-} for identical and non-identical boson pairs.

  The high-statistics data sample available at the ψ/J allows a clear demonstration that the enhancement is present for identical particles (two pions) and not for non-identical particles ($K\pi$). Figure 1 shows R^{--}_{+-} for $Q^2(\pi\pi)$ and $Q^2(K\pi)$ distributions integrated over all other variables. These distributions are fitted with $(1+\alpha e^{-\beta Q^2})\gamma$ where γ is a normalizing constant for the local region considered. This fit is made in regions which avoid known resonances (K^o region, ρ^o region, etc.) and spurious effects (converted γ rays yield e^+e^- pairs which are partly misidentified as $\pi^+\pi^-$ pairs).* Thus only the Q^2 regions 0.02 - 0.15, 0.19 - 0.38 are used for the fits.

---

†Work supported primarily by the Director, Office of Energy Research, Division of HighEnergy and Nuclear Physics of the U.S. Department of Energy under Contracts DE-AC0-3-76SF00515 and W-7405-ENG-48.

*Furthermore Coulomb repulsion effects which also occur in the very low Q^2 region have been ignored here.

‡Results on S* resonance: G. Gidal et al., LBL-13239, submitted to Phys. Lett.

Results:	α	β (GeV/c)$^{-2}$	r (Fermi)
$\pi\pi$	0.71 ± 0.03	18.7 ± 0.8	0.85 ± 0.02
$K\pi$	0.16 ± 0.04	10.0 ± 4.1	0.62 ± 0.12

Conclusion: The observed enhancement occurs for identical particles only; it is not the result of some reflection of resonances.*

- Dependence on $\delta = ||\vec{p}_1| - |\vec{p}_2||$.

  If we define R in terms $(1/\sigma)(d^2\sigma/dQ^2(12)d\delta)$, we can study the maximal Bose-Einstein effect as $\delta \to 0$. Figure 2 shows R^{--}_{+-} for several δ intervals as shown.

  Conclusion: The corresponding fits to α and β are given in Table I and show that as $\delta \to 0$, $\alpha \to 1$ indicating maximal chaoticity at the ψ/J.

- Three-pion correlations.

  Figure 3 shows R^{---}_{+--} at the ψ/J. We obtain a very striking enhancement at low $Q^2(123)$ values. Here we must remember that ideally we would like to study R^{---}_{+-o} and that R^{---}_{+--} contains an enhancement in the denominator as well as in the numerator. The total enhancement factor expected for R^{---}_{+-o} can be approximated by the product of $(1+\alpha_2)(1+\alpha'_3)$ where α_2 is the two-body value for $\psi \to 3\pi^+ + 3\pi^- + X$ and α'_3 is the value from the fit to R^{---}_{+--} (see Table II). The total enhancement of ~ 6 at the ψ/J again indicates maximal chaoticity.

Results from the $E_{cm} = 4 - 7$ GeV Region

This is the region where jet formation begins and the e^+e^- annihilation is believed to proceed via $q\bar{q}$ rather than via ggg as at the ψ/J. Table II gives the fits to R^{--}_{+-} and R^{---}_{+--} for this region (figure not shown).

Conclusion: There is a dramatic decrease in the enhancement between ψ/J and the 4 - 7 GeV region. Whether this represents a dynamic effect[3] or a decrease in chaoticity (more coherence) is not clear at present. It will be interesting to learn whether the Υ will behave like the ψ/J or not. We also note that r differs markedly for the two- and three-body system but is not very different on and off the ψ/J resonance.[5]

References

1. G. Goldhaber, S. Goldhaber, W. Lee and A. Pais, Phys. Rev. 120, 300 (1960). This paper gives an independent derivation of Bose-Einstein enhancements for the $\pi\pi$ system. The results are related to what has become known as the "Hanbury-Brown/Twiss effect" for the two-photon system. R. Hanbury-Brown and R. Q. Twiss, Philos. Mag. 45, 663 (1954), Nature 178, 1046 (1956); E. M. Purcell, Nature 178, 1449 (1956).
2. G. N. Fowler and R. M. Weiner, Phys. Lett. 70B, 201 (1977).
3. A. Giovannini and G. Veneziano, Nucl. Phys. B130, 61 (1977) and references to experiments given in this paper.

*The small enhancement ($\alpha = 0.16$) for the $K\pi$ system may be due to feedthrough because of K-π misidentification, although a small $I = 3/2$ effect cannot be ruled out.

4. G. I. Kopylov, Phys. Lett. 50B, 472 (1974); G. I. Kopylov and M. I. Podgoretskii, Sov. J. Nucl. Phys. 19, 215 (1974); M. Gyulassy, S. K. Kaufmann and L. W. Wilson, Phys. Rev. C20, 2267 (1979).

5. G. Cocconi, Phys. Lett. 49B, 459 (1974) and private communication.

Table I. Results from fits of $(1+\alpha e^{-\beta Q^2})^\gamma$ to the $\pi\pi$ data at the $\psi/J(3.1)$ for various δ regions.

δ (GeV/c)	α	β (GeV/c)$^{-2}$	r (Fermi)
< 0.1	0.94 ± 0.01	22.2 ± 0.2	0.93 ± 0.01
0.1, 0.2	0.71 ± 0.03	16.0 ± 1.2	0.79 ± 0.03
0.2, 0.3	0.55 ± 0.04	12.6 ± 1.7	0.70 ± 0.05
> 0.3	0.44 ± 0.05	10.5 ± 2.5	0.64 ± 0.07

Table II. Parameters from Bose-Einstein correlation fits as a function of E_{cm} for $3\pi^+ + 3\pi^- + X$ final states.

	α_2	r_2 (Fermi)	α'_3	r_3 (Fermi)	$(1+\alpha_2)(1+\alpha'_3)$
$\psi/J(3.1)$	0.89 ± 0.03	0.78 ± 0.02	2.33 ± 0.06	0.49 ± 0.003	6.2 ± 0.14
4 - 7 GeV	0.52 ± 0.06	0.77 ± 0.08	1.09 ± 0.11	0.39 ± 0.04	3.3 ± 0.3



Fig. 1. The R_U^L distribution for $\pi\pi$ and $K\pi$ systems as function of Q^2 at the ψ/J. The shaded regions along the x-axis in (a) represent the regions left out in the fits both here and in Fig. 2. The curves are the result of fits to $(1 + \alpha e^{-\beta Q^2})^\gamma$.



Fig. 2. The $R_U^L(Q^2,\delta)$ distribution for $\pi\pi$ at the ψ/J.



Fig. 3. The $R^{---}_{+--}(Q^2)$ distribution for $\pi\pi\pi$ at the ψ/J.