Submitted to Biochimica et
Biophysica Acta

LBL-1591
Preprint c.1

AN ELECTRON PARAMAGNETIC RESONANCE STUDY OF THE BINDING OF MANGANESE TO RIBULOSE 1, 5-DIPHOSPHATE CARBOXYLASE--INHIBITION BY MAGNESIUM

D.K. Chu, R. Chang, and L.E. Vickery

RECEIVED
LAWRENCE
RADIATION LABORATORY

February 1973

LIBRARY AND
DOCUMENTS SECTION

Prepared for the U.S. Atomic Energy
Commission under Contract W-7405-ENG-48

For Reference

Not to be taken from this room

LBL-1591
c.1

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

0 0 0 0 3 9 0 0 9 3 8

An electron paramagnetic resonance study of the binding of manganese to ribulose 1,5-diphosphate carboxylase--inhibition by magnesium

D. K. CHU, R. CHANG*, and L. E. VICKERY**

Laboratory of Chemical Biodynamics, Lawrence Berkeley Laboratory, University of California, Berkeley, California 94720 (U.S.A.)

SUMMARY

Electron paramagnetic resonance (EPR) was used to measure the extent of binding of Mn^{2+} to ribulose 1,5-diphosphate carboxylase purified from spinach leaves. In the Mn^{2+} concentration range 0.60 mM to 0.04 mM only a single Mn^{2+} binding site having K_d = 0.06 mM was observed. Mg^{2+} was found to competitively inhibit the binding of Mn^{2+}, and K_d = 0.54 mM was calculated for Mg^{2+}.

D-Ribulose 1,5-diphosphate carboxylase (3-phospho-D-glycerate carboxylase [dimerizing] EC 4.1.1.39) catalyzes the carboxylation of RuDP to yield two moles of 3-D-phosphoglycerate in the primary carbon fixation step of photosynthesis. The reaction has an absolute requirement for a divalent metal cation[1], and it has been suggested that control of metal ion activation might play a role in the regulation of the enzyme's activity *in vivo*[2]. Little is known, however, about the stoichiometry of

---

*On leave, permanent address: Department of Chemistry, Williams College, Williamstown, Mass. 01267.

**Postdoctoral Fellow of the National Institute of General Medical Sciences, National Institutes of Health.

the metal ion-enzyme interaction, the nature of the groups on the protein involved in the interaction, or the mechanism of the activation. In fact, the only cases where binding of metal ions to the enzyme has been studied involved inactive ternary complexes of the enzyme with a substrate analogue or complexes with cyanide[3]. Putative evidence for the existence of an enzyme-CO_2-Mg^{2+} has also been reported[4].

We have used electron paramagnetic resonance techniques to study the binding of Mn^{2+} to the enzyme. Free Mn^{2+} ion in solution has an intense isotropic EPR spectrum, but the EPR of Mn^{2+} bound to small molecules and proteins is often broadened and undetectable providing a convenient means of measuring binding constants[5-7]. Electrophoretically pure RuDP carboxylase was isolated from spinach leaves as previously described[8] with the following modifications: the heating step was no longer used and 0.02 M sodium phosphate, pH 7.4, replaced tris-HCl and Mg^{2+} during purification. Freshly prepared enzyme was concentrated by precipitation with ammonium sulfate, resuspended in 0.05 M Tris, pH 7.8, and exhaustively dialyzed against this buffer. Protein concentrations were determined spectrophotometrically, taking the absorbancy at 280 nm to be 0.61 $(cm\text{-}mg/ml)^{-1}$ and assuming a molecular weight of 557,000 [9]. X-band EPR spectra were recorded on a Varian E-3 spectrometer.

When RuDP carboxylase was added to solutions of Mn^{2+} in buffer a quantitative reduction in the intensity of the Mn^{2+} EPR signal was observed. No changes in the EPR spectrum of the aquo-complex appeared, and no evidence for any new signal was obtained. A titration of the

---

Abbreviations: EPR, electron paramagnetic resonance; RuDP, D-ribulose 1,5-diphosphate.

enzyme was made assuming that the heights of the hyperfine components in the first derivative EPR spectrum are proportional to the concentration of free Mn^{2+}. The results of three experiments, using different enzyme preparations and concentrations, are presented in the form of a Hughes-Klotz plot[7,10] in Fig. 1. In the absence of competing divalent cations the data fit a single straight line reasonably well. The x and y intercepts are equal to the reciprocals of the dissociation constant and the number of binding sites, respectively. Within the Mn^{2+} concentration range used, 0.80 mM to 0.04 mM, $K_d \approx 0.06$ mM and n = 1. When Mg^{2+} was added to solutons of enzyme and Mn^{2+}, an increase in the free Mn^{2+} EPR signal was observed (Fig. 1). In the presence of 0.6 and 1.2 mM Mg^{2+} the K_d for Mn^{2+} increased to 0.12 mM and 0.18 mM, respectively, while n remained equal to 1. Assuming competitive binding of the two cations, K_d for Mg^{2+} can be determined[11,12]. The results of separate calculations at each Mn^{2+} and Mg^{2+} concentration are shown in Table I. The average value for K_d (Mg^{2+}) = 0.54 mM.

The dissociation constant for Mn^{2+} in the absence of Mg^{2+} measured in this work is close to the Michaelis constant for Mn^{2+}, K_m = 0.04 mM, determined by others[9]. This suggests that the binding observed might yield the active enzyme-metal complex required for catalysis. The stoichiometry of the Mn^{2+}-enzyme interaction observed here, however, is unexpected. RuDP carboxylase is composed of sixteen subunits of molecular weight of approximatley 12,000 and 56,000 daltons in a 1:1 molar ratio[13]. Moreover, there seem to be eight binding sites for RuDP[3]. The presence of a single Mn^{2+} binding site could be explained by a unique site linking several subunits, by an undetected difference in one subunit,

or by the existence of an additional subunit. This Mn^{2+} binding site may be related to the single Cu^{2+} binding site which has been reported[3] since our preparations of the enzyme do not appear to contain copper (Vickery, Chang, and Chu, unpublished).

The dissociation constant determined for Mg^{2+} competition with Mn^{2+} binding is significantly lower than the Michaelis constant for Mg^{2+}, K_m = 1.1 mM [9]. Thus, this binding may not be limiting for activation of the enzyme. The ability of Mg^{2+} to compete with Mn^{2+} in this concentration range might compensate for the lower K_m for Mn^{2+} and rule out the possibility of Mn^{2+} activation of the enzyme *in vivo*. Further experiments utilizing the Mn^{2+} EPR probe can be designed to determine the effects of temperature, pH, CO_2, RuDP and competing ions on the metal-enzyme interaction.

This work was supported, in part, by the U. S. Atomic Energy Commission.

REFERENCES

1 Weissbach, A., Horecker, B. L. and Hurwitz, J. (1956) *J. Biol. Chem.* 218, 795-810.

2 Bassham, J. A. and Kirk, M. (1968) in *Comparative Biochemistry and Biophysics of Photosynthesis* (Shibata, K. *et al.*, eds), pp. 365-378, Univ. Tokyo Press.

3 Wishnick, M., Lane, M. D. and Scrutton, M.C. (1970) *J. Biol. Chem.* 245, 4939-4947.

4 Akoyunoglou, G. and Calvin, M. (1963) *Biochem. Z.* 338, 20-30.

5 Cohn, M. and Townsend, J. (1954) *Nature* 173, 1090-1091.

6 Malmstrom, B. G., Vanngard, T. and Larsson, M. (1958) *Biochim. Biophys. Acta* 30, 1-5.

7 Mildvan, A. S. and Cohn, M. (1963) Biochemistry 2, 910-919.

8 Chu, D. K. and Bassham, J. A. (1972) Plant Physiol. 50, 224-227.

9 Paulsen, J. M. and Lane, M. D. (1966) Biochemistry 5, 2350-2357.

10 Hughes, T. R. and Klotz, I. M. (1956) Meth. Biochem. Anal. 3, 265-299.

11 Hastings, A. B., McLean, F. L., Eichelberger, L., Hall, J. L. and Da Costs, E. (1934) J. Biol. Chem. 107, 351-370.

12 Mildvan, A. S. and Cohn, M. (1964) J. Biol. Chem. 240, 238-246.

13 Rutner, A. C. (1970) Biochem. Biophys. Res. Commun. 39, 923-929.

TABLE I

DISSOCIATION CONSTANTS OF MANGANESE AND MAGNESIUM RuDP CARBOXYLASE

Experimental conditions are given in Fig. 1 and text. K_d^{Mn} was determined from best straight line fit to data; K_d^{Mg} was calculated for each data according to reference 11, assuming competition of Mg^{2+} with Mn^{2+}.

Mn^{2+} (mM)	Mg^{2+} (mM)	K_d^{Mn} (mM)	K_d^{Mg} (mM)
.04 to .60	0	.06	--
.06	0.6	.12	.54
.12	0.6	.12	.53
.18	0.6	.12	.57
.06	1.2	.18	.53
.12	1.2	.18	.48
.18	1.2	.18	.58
		average	.54

FIGURE LEGENDS

Fig. 1. Titration of RuDP carboxylase with Mn^{2+}. Hughes-Klotz plot of the concentrations of free and bound Mn^{2+} determined by EPR (see text). Experimental conditions: buffer, 0.05 M Tris, pH 7.8; temperature, 22°; sample volume, 0.15 ml; enzyme concentrations Δ Δ Δ 148 mg/ml, o o o 89 mg/ml, ● ● ● 48 mg/ml.


+1.2 mM Mg^{2+}
+0.6 mM Mg^{2+}
$[RuDPCase]/[Mn]_{bound}$
8
6
4
2
-2
0
2
4
6
8
10
12
$\frac{1}{[Mn]_{free}} \times 10^{-4}$ M

XBL731-4633

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720